

## **COELIAC DISEASE STUDY GROUP**

**University of Tampere  
Tampere University Hospital**

### **LIST OF PUBLICATIONS**

	<b>Page</b>
<b>Completed academic dissertations (MD and PhD thesis programs).....</b>	<b>2</b>
<b>Advanced studies.....</b>	<b>3</b>
<b>Original articles.....</b>	<b>4</b>
<b>Submitted manuscripts.....</b>	<b>18</b>
<b>Patents.....</b>	<b>18</b>
<b>Films.....</b>	<b>18</b>
<b>Abstracts.....</b>	<b>19</b>

## Completed academic dissertations (MD and PhD thesis programs)

1. Aine L. Dental enamel defects and dental maturity in children and adolescents with coeliac disease. University of Turku. Proceedings of the Finnish Dental Society 1986;82(Suppl 3):1-77.
2. Hällström O. IgA-class reticulin antibody test in coeliac disease and dermatitis herpetiformis. Acta Universitatis Tamperensis, University of Tampere. Tampere 1989;281(ser A):1-166.
3. Holm K. Latent and silent coeliac disease. A family study. Acta Universitatis Tamperensis, Tampere 1993;371(ser A):1-141.
4. Collin P. Associated diseases and survival in coeliac disease. Acta Universitatis Tamperensis, University of Tampere. Tampere 1994;405(ser A):1-96.
5. Lähdeaho M-L. Homology between adenovirus E1B protein and A-gliadin. Adenovirus and A-gliadin specific B cell responses in coeliac disease. Acta Universitatis Tamperensis, University of Tampere. Vammala 1994;430(ser A):1-124.
6. Polvi A. Genetic susceptibility to celiac disease: Major histocompatibility complex genes in celiac disease and in its animal model. Finnish Red Cross Blood Transfusion Service, No. 37, University of Helsinki. Helsinki 1998;1-106.
7. Kerttula T. Circulating T-cell subsets in coeliac disease and HLA DQ2-positive individuals. Acta Universitatis Tamperensis 666, University of Tampere. Vammala 1999;1-102.
8. Iltanen S. Markers of coeliac disease latency. Acta Universitatis Tamperensis 682, University of Tampere. Vammala 1999;1-140.
9. Lähteenoja H. Coeliac disease and oral mucosa. Annales Universitatis Turkuensis. Ser. D Medica-Odontologica. University of Turku. Turku 1999; 1-182.
10. Sulkanen S. Serum antibodies in coeliac disease. Acta Universitatis Tamperensis; 728, University of Tampere, Tampere 2000;1-124. Acta Electronica Universitatis Tamperensis, 15, <http://acta.uta.fi>. Tampereen yliopisto 2000.
11. Kaukinen K. Small bowel mucosa in suspected and confirmed coeliac disease. Acta Universitatis Tamperensis; 770, University of Tampere. Tampere 2000;1-122. Acta Electronica Universitatis Tamperensis, 62, <http://acta.uta.fi>. Tampereen yliopisto 2000.
12. Halttunen T. Biological functions of coeliac disease autoantibodies. Acta Universitatis Tamperensis, 865, University of Tampere, Tampere 2002; 1-126. Acta Electronica Universitatis Tamperensis; 171, <http://acta.uta.fi>. Tampereen yliopisto 2002.
13. Holopainen P. Genetic susceptibility to celiac disease: HLA-unlinked candidate genes. Academic dissertations from the Finnish Red Cross Blood Transfusion Service, No. 45; 2002;1-108. <http://ethesis.helsinki.fi>. University of Helsinki.
14. Karell Kati. Dissecting genetic susceptibility to gluten sensitivity: HLA-linked risk factors in coeliac disease and dermatitis herpetiformis. Academic dissertations from the Finnish Red Cross Blood Transfusion Service, No. 47; 2003;1-105. <http://ethesis.helsinki.fi>. University of Helsinki.
15. Luostarinen Liisa. Neurological manifestations in coeliac disease. Acta Universitatis Tamperensis;938, Tampereen yliopisto, Tampere 2003. Acta Electronica Universitatis Tamperensis;258, Tampereen yliopisto 2003, <http://acta.uta.fi>.
16. Mustalahti Kirsi. Silent coeliac disease. Acta Universitatis Tamperensis, 946. Tampereen yliopisto, Tampere 2003;1-134. Acta Electronica Universitatis Tamperensis 269, <http://acta.uta.fi>.
17. Lindfors Katri. Differential gene expression in the intestinal crypt-villus axis. Klingendahl Paino Oy, Tampere 2004;1-130. Acta Electronica Universitatis Tamperensis 357, <http://acta.uta.fi>. Tampereen yliopisto 2004.
18. Hervonen Kaisa. Familial occurrence of dermatitis herpetiformis and associated conditions. Acta Universitatis Tamperensis, 1026, Tampereen yliopisto, Tampere 2004. Acta Electronica Universitatis Tamperensis, 371, Tampereen yliopisto 2004, <http://acta.uta.fi>.
19. Viljamaa Mervi. Evaluation of the results of active diagnostics and treatment of coeliac disease in a high-prevalence area - a special focus on dietary compliance, quality of life and associated complications. Acta Universitatis Tamperensis, 1171, Tampere 2006. Acta Electronica Universitatis Tamperensis, 550, <http://acta.uta.fi>.
20. Salmi Teea. Diagnosing coeliac disease: beyond the conventional histology. Acta Universitatis Tamperensis, 1180, Tampere 2006. Acta Electronica Universitatis Tamperensis, 559, <http://acta.uta.fi>.
21. Peräaho Markku. Dietary aspects in the diagnosis and treatment of coeliac disease. Acta Universitatis Tamperensis, 1220, Tampere 2007. Acta Electronica Universitatis Tamperensis 606, <http://acta.uta.fi>.
22. Raivio Tiina. Whole blood self-tissue transglutaminase-based antibody testing in coeliac disease. Acta Universitatis Tamperensis 1310, Tampere 2008. Acta Electronica Universitatis Tamperensis, 718, <http://acta.uta.fi>.
23. Juuti-Uusitalo Kati. Functional genomics of a three-dimensional epithelial cell culture and coeliac small intestinal mucosal biopsy samples. Acta Universitatis Tamperensis 1319, Tampere 2008. Acta Electronica Universitatis Tamperensis, 730, <http://acta.uta.fi>.
24. Haimila Katri. Genetics of T cell co-stimulatory receptors –CD28, CTLA4, ICOS and PDCD1 in immunity and transplantation. Academic dissertations from the Finnish Red Cross Blood Service, Number 54; 2009:1-107. <http://ethesis.helsinki.fi>. University of Helsinki.

25. Viiri K. The Sin3-Associated Protein 30 (SAP30) Family of Transcriptional Regulators. Acta Universitatis Tamperensis; 1396, Tampere University Press, Tampere 2009. Acta Electronica Universitatis Tamperensis, 824. <http://acta.uta.fi>.

## Advanced studies

1. Kallonen K. Muuttuva lasten keliakiatauti. (in Finnish). Advanced studies. University of Tampere, Medical School, 1985:1-48.
2. Lähdeaho M-L. Retikuliinivasta-aineet lasten keliakian seulontakokeena. (Reticulin antibodies as a screening test in childhood coeliac disease, in Finnish). Advanced studies. University of Tampere, Medical School, Study of childhood coeliac disease. 1985:1-46.
3. Hartikainen M. Keliakia. Ohutsuolikoepalojen morfometrinen tutkimus. (in Finnish). Advanced studies. University of Tampere, Medical School, Coeliac Disease Study Project. 1994:1-32.
4. Lindfors K. Ohutsuolen krypta-villusakseli. (Intestinal crypt-villus axis, in Finnish). Bachelor's degree. University of Turku, Faculty of Mathematics and Natural Sciences, Department of Biology, Laboratory of Genetics. 1997:1-40.
5. Leivo E-L. Coeliac Disease: Human umbilical cord-derived fibroblasts as antigen for detecting coeliac disease autoantibodies. (in Finnish). Advanced studies. University of Tampere, Medical School. 1997:1-26.
6. Lindfors K. Erilaistuneiden ja erilaistumattomien epiteelisolujen geeniekspressioerojen tutkiminen differential display reverse transcription PCR:llä. (Differential gene expression between differentiated and undifferentiated epithelial cell by differential display reverse transcription PCR, in Finnish). Master's thesis. University of Turku, Faculty of Mathematics and Natural Sciences, Department of Biology, Laboratory of Genetics 1998:1-57.
7. Holopainen P. Kromosomin 5q ehdokasgeenialueiden kytkentä- ja assosiaatioanalyysi suomalaisissa keliakiaperheissä. (Linkage and association analysis of candidate loci at chromosome 5q in Finnish families with coeliac disease.) Master's thesis. University of Jyväskylä, Department of Biological and Environmental Science. 1999:1-60.
8. Vähänen M. Lasten keliakian taudinkuva kaura- ja vehnäaltistuksen aikana. Advanced studies. University of Tampere, Medical School, 1999:1-35.
9. Huotari P. Ihmisen AATF:n karakterisointi ja rekombinanttiproteiinin tuotto (Characterization of human AATF and production of recombinant human AATF protein, in Finnish). Master's thesis. University of Jyväskylä, Department of Biological and Environmental Science. 2001:1-69.
10. Rannila T. Piilevän keliakian yleisyys suomalaisessa "koilliskaira"-materiaalissa. Advanced studies. University of Tampere, Medical School, 2003: 1-23.
11. Kärki S. Keliakialasten sukulaisten syöpätapaukset. Advanced studies. University of Tampere, Medical School, 2005: 1-29.
12. Stenman S. Ihmisen ohutsuolen kudosiseljelymalli: vehnän ja gliadiinin vaikutukset normaalissa ja keliakikon ohutsuolessa *ex vivo*. (The organ culture of human small intestine: the effects of wheat gliadin on the small intestine in normal mucosa and in coeliac disease *ex vivo*.) Master's thesis. University of Jyväskylä, 2006.
13. Syrjänen Mari. Keliakian vaikutus verisuonten endoteeli- ja sileälihassoluihin ohutsuolen limakalvolla ja soluviljelymallissa. (The effect of celiac disease on the vascular endothelial and smooth muscle cells in the mucosa of the small bowel and in a cell culture model.) Master's thesis. University of Jyväskylä, 2006:1-56.
14. Wegelius A. Characterization of the binding specificities exhibited by the periferal autoantibody response in coeliac disease. Advanced studies. University of Tampere, Medical School 2006: 1-45.
15. Ylilehto Elina. Kudostransglutaminaasin (TG2) inhibiittorin vaikutukset hiirellä. Advanced studies. University of Tampere, Medical School 2006:1-22.
16. Jokinen S. Ohutsuolikoepalan kudosiseljely ja keliakia. Advanced studies. University of Tampere, Medical School 2007.
17. Raivio T. Uusi ja nopea kokoveritesti keliakian diagnostiikassa. (Self transglutaminase-based rapid coeliac disease antibody detection by a lateral flow method). Advanced studies. University of Tampere, Medical School, 2007.
18. Blomqvist T. Vehnän gliadiinin haittavaikutusten estäminen elävien probioottibakteerien avulla suolen epiteelisoluviljelmissä. Master's thesis. University of Tampere, Institute of Medical Technology, 2007.
19. Koskinen O. Gluten-dependent small-bowel mucosal transglutaminase-2 specific IgA deposits in overt and mild enteropathy coeliac disease. Advanced studies. University of Tampere, Medical School, 2007.
20. Villanen M. Ohutsuolen limakalvon transglutaminaasi-2 autovasta-ainekertymät keliakiaa sairastavilla lapsilla kaura- ja gluteenialtistuksen aikana. (Oats does not induce systemic or mucosal autoantibody response in children with coeliac disease.) Advanced studies. University of Tampere, Medical School, 2008.
21. Pesonen J. SAP30L-kandidaattigeenin sekvensointi ja mutaatioiden kartoitus keliakiapotilailta. Advanced studies. University of Tampere, Medical School, 2008.
22. Ränsi S. Dlx-geeniperhe. Master's thesis. University of Tampere, Institute of Medical Technology, 2008.
23. Hautala A. Farmakologisen hoidon mahdollisuudet keliakiassa (Possibilities of pharmacological treatments in coeliac disease). Master's thesis. University of Helsinki, Department of Pharmacology and Toxicology, 2008: 1-74.

24. Räsänen T. Keliakian vasta-ainetiittereiden korrelaatio oireisiin ja ohutsuolen limakalvovaurioihin. (Correlation of celiac autoantibody levels between symptoms and small intestinal mucosal villous damage). Advanced studies. University of Tampere, Medical School, 2008; 1-17.
25. Siltanen A. Genetic variation in the Calreticulin 3 gene among celiac disease patients. Master's thesis. University of Jyväskylä, 2008.
26. Virtanen S.  $\beta$ 1,3-Glukosyylitransferaasin tuoton optimointi ja puhdistus. Master's thesis. University of Tampere, Institute of Medical Technology, 2008; 1-86.

## Original articles

1. Mäki M, Hällström O, Vesikari T, Visakorpi JK. Evaluation of a serum IgA class reticulin antibody test for the detection of childhood celiac disease. *J Pediatr* 1984; 105:901-5.
2. Mäki M, Hällström O, Huupponen T, Vesikari T, Visakorpi JK. Increased prevalence of coeliac disease in diabetes. *Arch Dis Child* 1984;59:739-42.
3. Mäki M. Guide wire manipulation of Crosby jejunal biopsy capsule. *Br Med J* 1984; 288:862-3.
4. Hällström O, Reunala T. IgA class reticulin antibodies in dermatitis herpetiformis: a good indicator of jejunal damage. *Acta Derm Venereol (Stockh)* 1985;65:330-65.
5. Rantala I, Mäki M, Laasonen A, Visakorpi JK. Periodate-lysine-paraformaldehyde as fixative for the study of duodenal mucosa. Morphologic and immunohistochemical results at light and electron microscopic levels. *Acta Path Microb Immunol Scand Sect A* 1985;93:165-73.
6. Vainio E, Collin P, Lehtonen O-P. Avidity of antigliadin IgA and IgG antibodies in gluten-sensitive enteropathy and dermatitis herpetiformis. *Clin Immunol Immunopathol* 1986;41:295-300.
7. Vainio E, Kosnai I, Hällström O, Karpati S, Mäki M, Reunala T. Antigliadin and antireticulin antibodies in children with dermatitis herpetiformis. *J Pediatr Gastroent Nutr* 1986;5:735-9.
8. Teppo A-M, Mäki M, Hällström O, Maury CPJ. Antibodies to 90 kilodalton glycoprotein in childhood and adolescent celiac disease: Relationship to reticulin antibodies. *J Pediatr Gastroent Nutr* 1987;6:908-14.
9. Auricchio S, Mazzacca G, Tosi R, Visakorpi J, Mäki M, Polanco I. Coeliac disease as a familial condition: Identification of asymptomatic coeliac patients within family groups. *Gastroenterol Int* 1988; 1:25-31.
10. Mäki M, Hällström O, Verronen P et al. Reticulin antibody, arthritis, and coeliac disease in children. *Lancet* 1988;I:479-80.
11. Mäki M, Kallonen K, Lähdeaho M-L, Visakorpi JK. Changing pattern of childhood coeliac disease in Finland. *Acta Paediatr Scand* 1988;77:408-12.
12. Ståhlberg M-R, Hietanen E, Mäki M. Musosal biotransformation rates in the small intestine of children. *Gut* 1988;29:1058-63.
13. Collin P, Salmi J, Hällström O, Oksa H, Oksala H, Mäki M, Reunala T. High frequency of coeliac disease in adult patients with type-I diabetes. *Scand J Gastroenterol* 1989; 24:81-4.
14. Hällström O. Comparison of IgA-class reticulin and endomysium antibodies in coeliac disease and dermatitis herpetiformis. *Gut* 1989;30:1225-32.
15. Mäki M, Lähdeaho M-L, Hällström O, Viander M, Visakorpi JK. Postpubertal gluten challenge in coeliac disease. *Arch Dis Child* 1989;64:1604-07.
16. Aine L, Mäki M, Collin P, Keyriläinen O. Dental enamel defects in celiac disease. *J Oral Pathol Med* 1990;19:241-45.
17. Collin P, Hällström O, Mäki M, Viander M, Keyriläinen O. Atypical coeliac disease found with serologic screening. *Scand J Gastroenterol* 1990;25:245-50.
18. Mäki M, Holm K, Koskimies S, Hällström O, Visakorpi JK. Normal small bowel biopsy followed by coeliac disease. *Arch Dis Child* 1990;65:1137-41.
19. Mäki M, Holm K. Incidence and prevalence of coeliac disease in Tampere. Coeliac disease is not disappearing. *Acta Paediatr Scand* 1990;79:980-82.
20. Mäki M. Keliakiatauti (Coeliac disease, in Finnish). *Suomen Hammlääkl* 1990;37:544-47.
21. Pasternack A, Collin P, Mustonen J, Reunala T, Rantala I, Laurila K, Teppo A-M. Glomerular IgA deposits in patients with celiac disease. *Clin Nephrol* 1990;34:56-60.
22. Walker-Smith JA, Guandalini S, Schmitz J, Shmerling DH, Visakorpi JK. Revised criteria for diagnosis of coeliac disease. *Arch Dis Child* 1990;65:909-911.
23. Aine L, Reunala T, Mäki M. Dental enamel defects in children with dermatitis herpetiformis. *J Pediatr* 1991;118:572-4.
24. Collin P, Pirttilä T, Nurmikko T, Somer H, Erilä T, Keyriläinen O. Coeliac disease, brain atrophy and dementia. *Neurology* 1991;41:372-5.
25. Koger O, Uibo R, Mäki M. Coeliac disease in children. (In Russian.). *Pediatrics* 1991;10:79-85.
26. Koskimies S, Lipsanen V, Mäki M, Visakorpi JK. HLA and coeliac disease, a Finnish family study. In: Mearin ML, Mulder CJJ, eds. *Coeliac Disease, 40 years gluten-free*. Kluwer Academic Publishers, Dordrecht, 1991:9-13.
27. Koskimies S, Mäki M, Lipsanen V, Visakorpi J. HLA haplotypes in Finnish coeliac patients and their family members, possible protecting haplotypes. In: Kumar PJ, Walker-Smith JA, eds. *Coeliac Disease: One Hundred Years*. Univ. Printing Service, Univ. Leeds, Leeds, 1991:51-53.

28. Mäki M, Aine L, Lipsanen V, Koskimies S. Dental enamel defects in first-degree relatives of coeliac disease patients. *Lancet* 1991;337:763-4.
29. Mäki M, Holm K, Lipsanen V, Hällström O, Viander M, Collin P, Savilahti E, Koskimies S. Serological markers and HLA genes among healthy first-degree relatives of patients with coeliac disease. *Lancet* 1991;338:1350-53.
30. Mäki M, Holm K, Collin P, Savilahti E. Increase in gamma/delta T cell receptor bearing lymphocytes in normal small bowel mucosa in latent coeliac disease. *Gut* 1991;32:1412-4.
31. Mäki M, Holm K, Koskimies S, Visakorpi JK. Latent coeliac disease. In: Mearin ML, Mulder CJJ, eds. *Coeliac Disease, 40 years gluten-free*. Kluwer Academic Publishers, Dordrecht, 1991:153-6.
32. Mäki M, Hällström O, Marttinen A. Reaction of human non-collagenous polypeptides with coeliac disease autoantibodies. *Lancet* 1991;338:724-25.
33. Aine L, Mäki M, Reunala T. Coeliac-type dental enamel defects in patients with dermatitis herpetiformis. *Acta Derm Venereol* 1992;72:25-27.
34. Collin P, Korpela M, Keyriläinen O, Hällström O, Viander M, Mäki M. Rheumatic complaints as a presenting symptom in patients with celiac disease. *Scand J Rheumatol* 1992;21:20-3.
35. Collin P, Mäki M, Keyriläinen O, Hällström O, Reunala T, Pasternack A. Selective IgA-deficiency and coeliac disease. *Scand J Gastroenterol* 1992;27:367-71.
36. Greco L, Mäki M, Di Donato F, Visakorpi JK. Epidemiology of coeliac disease in Europe and the mediterranean area. A summary report on the multicentre study by the European Society of Paediatric Gastroenterology and Nutrition. In: Auricchio S, Visakorpi JK, eds. *Common Food Intolerances 1: Epidemiology of Coeliac Disease*. Dyn Nutr Res, Basel, Karger 1992;2:25-44.
37. Holm K, Mäki M, Savilahti E, Lipsanen V, Laippala P, Koskimies S. Intraepithelial gamma/delta T-cell-receptor lymphocytes and genetic susceptibility to coeliac disease. *Lancet* 1992;339:1500-03.
38. Lähdeho M-L, Mäki M, Parkkonen P, Kantonen T, Lehtinen M. Determination of antibodies to non-structural and structural adenovirus antigens by enzyme-linked immunosorbent assay. *Acta Virol* 1992;36:524-32.
39. Mäki M, Hällström O, Marttinen A, Lipsanen V, Holm K, Collin P, Koskimies S. Screening tools for use in coeliac disease. In: Auricchio S, Visakorpi JK, eds. *Common Food Intolerances 1: Epidemiology of Coeliac Disease*. Dyn Nutr Res, Basel, Karger 1992;2:93-104.
40. Mäki M, Holm K, Ascher H, Greco L. Factors affecting epidemiological trends in coeliac disease. Type and amount of cereals in the diet. In: Auricchio S, Visakorpi JK, eds. *Common Food Intolerances 1: Epidemiology of Coeliac Disease*. Dyn Nutr Res, Basel, Karger 1992;2:76-82.
41. Mäki M. Use of serological antibody tests in coeliac disease. In: Branski D, Rozen P, Kagnoff M, eds. *Gluten-Sensitive enteropathy*. Front Gastrointest Res. Basel, Karger 1992;19:108-29. (review, includes original material)
42. Savilahti E, Mäki M, Holm K, Koskimies S, Lipsanen V, Reunala T, Klemola T. Immunological markers of latent coeliac disease. In: Auricchio S, Visakorpi JK, eds. *Common Food Intolerances 1: Epidemiology of Coeliac Disease*. Dyn Nutr Res, Basel, Karger 1992;2:128-41.
43. Savilahti E, Reunala T, Mäki M. Increase of lymphocytes bearing the gamma/delta T-cell receptor in the jejunum of patients with dermatitis herpetiformis. *Gut* 1992;33:206-11.
44. Visakorpi JK. Silent coeliac disease: The risk groups to be screened. In: Auricchio S, Visakorpi JK, eds. *Common Food Intolerances 1: Epidemiology of Coeliac Disease*. Dyn Nutr Res, Basel, Karger 1992;2:84-92.
45. Ascher H, Holm K, Kristiansson B, Mäki M. Different features of coeliac disease in two neighbouring countries. *Arch Dis Child* 1993;69:375-80.
46. Collin P, Helin H, Mäki M, Hällström O, Karvonen AL. Follow-up of patients positive in reticulín and gliadin antibody tests with normal small-bowel biopsy. *Scand J Gastroenterol* 1993;28:595-98.
47. Holm K. Correlation of HLA-DR alleles to jejunal mucosal morphology in healthy first-degree relatives of coeliac disease patients. *Eur J Gastroenterol Hepatol* 1993;5:35-39.
48. Lähdeho M-L, Lehtinen M, Rissa H-R, Hyöty H, Reunala T, Mäki M. Antipeptide antibodies to adenovirus E1B protein indicate enhanced risk of coeliac disease and dermatitis herpetiformis. *Int Arch Allergy Immunol* 1993;101:272-76.
49. Lähdeho M-L, Parkkonen P, Reunala T, Mäki M, Lehtinen M. Antipeptide antibodies to the enteric adenovirus type 40 indicate enhanced risk of celiac disease and dermatitis herpetiformis. *Clin Immunol Immunopathol* 1993;69:300-5.
50. Mäki M, Marttinen A, Koskimies S. Molecular pathogenesis of coeliac disease. Autoimmune mechanisms operative? In: *Frontiers in Molecular and Clinical Gastroenterology*, Farthing MJG, Grejs GJ, eds. Syllabus of European Association for Gastroenterology and Endoscopy, Postgraduate Course-1993, 1993;4:1-12.
51. Marttinen A, Mäki M. Purification of fibroblast-derived celiac disease autoantigen molecules. *Pediatr Res* 1993;34:420-23
52. Partanen J, Milner C, Campbell RD, Mäki M, Lipsanen V, Koskimies S. HLA-linked heat shock protein 70 (HSP70-2) gene polymorphism and celiac disease. *Tissue Antigens* 1993;41:15-19.
53. Reunala T, Mäki M. Dermatitis herpetiformis: A genetic disease. *Eur J Dermatol* 1993;3:519-26.
54. Uibo O, Uibo R, Kleimola V, Jögi T, Mäki M. IgA-class antigliadin antibodies in an adult population sample: high prevalence without coeliac disease. *Dig Dis Sci* 1993;38:2034-37.
55. Visakorpi JK. The diagnosis of coeliac disease. *Annales Nestlé* 1993;51:43-49.
56. Aine L. Permanent tooth dental enamel defects leading to the diagnosis of coeliac disease. *Br Dent J* 1994;177:253-4.

57. Collin P, Mäki M. Associated disorders in coeliac disease. Clinical aspects. *Scand J Gastroenterol* 1994;29:769-75.
58. Collin P, Reunala T, Pukkala E, Laippala P, Keyriläinen O, Pasternack A. Coeliac disease-associated disorders and survival. *Gut* 1994;35:1215-1218.
59. Collin P, Salmi J, Hällström O, Reunala T, Pasternack A. Autoimmune thyroid disorders and coeliac disease. *Eur J Endocrinol* 1994;130:137-140.
60. Holm K, Savilahti E, Koskimies S, Lipsanen V, Mäki M. Immunohistochemical changes in the jejunum of first degree relatives of celiac patients. HLA class II antigen expression, interleukin 2 receptor positive cells and dividing crypt cells. *Gut* 1994;35:55-60.
61. Mäki M, Marttinen A, Koskimies S. Maladie coeliaque. Pathologie moléculaire et mécanismes auto-immuns. *Méd Chir Dig* 1994;23:215-17.
62. Mäki M. Autoantibodies as markers of autoimmunity in coeliac disease pathogenesis. In: Feighery C, O'Farrelly C, eds. *Gastrointestinal immunology and gluten-sensitive disease*. Oak Tree Press, Dublin 1994:246-52.
63. Marttinen A. Keliakiataudin autoantigeenit. *Solubiologi* 1994;2:39-40.
64. Visakorpi JK, Mäki M. Changing clinical features of coeliac disease. *Acta Paediatr* 1994;Suppl 395:10-13.
65. Collin P. Keliakian seulonta aikuisilla. *Suom lääkl* 1995;50:1729-33.
66. Kerttula TO, Hällström O, Mäki M. Phenotypical characterization of peripheral blood T cells in patients with coeliac disease - elevation of antigen-primed CD45RO+ T lymphocytes. *Immunology* 1995;86:104-9.
67. Lähdeaho ML, Vainio E, Lehtinen M, Parkkonen P, Partanen J, Koskimies S, Mäki M. Activation of coeliac disease immune system by specific  $\alpha$ -gliadin peptides. *Cereal Chem* 1995;72:475-9.
68. Mäki M, Hällström O, Sulkanen S, Marttinen A. Antibodies in coeliac disease. In: Auricchio S, Ferguson A, Troncone R, eds: *Mucosal Immunity and the Gut Epithelium: Interactions in Health and Disease*. Dyn Nutr Res, Basel, Karger 1995;4:115-126.
69. Mäki M, Huupponen T, Holm K, Hällström O. Seroconversion of reticulon autoantibodies predicts coeliac disease in insulin-dependent diabetes mellitus. *Gut* 1995;36:239-42.
70. Mäki M. Changing incidence of coeliac disease. International Symposium on the Coeliac Condition. Conference proceedings, Nutricia Dietary Products. The Governor's Hall, St. Thomas' Hospital, London, 27 October 1995; pp. 29-34.
71. Mäki M. Keliakian diagnostiikka. *Moodi* 1995;19:34-5.
72. Mäki M. The humoral immune system in coeliac disease. In: Howdle PD (ed.), *Coeliac disease*. Baillière's Clinical Gastroenterology 1995;9:231-49.
73. Visakorpi JK, Mäki M. Çöliak hastaliginin degisen klinik özellikleri. *Cocuk Sagligi ve Hastaliklari Dergisi* 1995;38:103-109.
74. Aine L. Coeliac-type permanent-tooth enamel defects. *Ann Med*, 1996;28:9-12.
75. Collin P, Pukkala E, Reunala T. Malignancies and survival in dermatitis herpetiformis: a comparison to coeliac disease. *Gut* 1996;38:528-30.
76. Collin P, Vilksa S, Heinonen PK, Hällström O, Pikkarainen P. Infertility and coeliac disease. *Gut* 1996;39:382-4.
77. Halttunen T, Mäki M. Epiteelisolujen biologisa suolen krypta-villus akselilla ja sen tutkiminen kolmiulotteisella soluviljelytekniikalla. *Solubiologi* 1996;14:51-57.
78. Halttunen T, Marttinen A, Rantala I, Kainulainen H, Mäki M. Fibroblasts and transforming growth factor  $\beta$  induce organization and differentiation of T84 human epithelial cells. *Gastroenterology* 1996;111:1252-62.
79. Jalava T, Mäki M, Marttinen A, Partanen J, Koskimies S. The in vitro response to human fibroblast-derived extracellular matrix proteins is restricted by specific HLA class II genes. Relevance for coeliac disease. *Human Immunol*, 1996;49:106-112.
80. Kerttula T, Collin P, Polvi A, Korpela M, Partanen J, Mäki M. Distinct immunological features of Finnish Sjögren's syndrome patients with the alleles HLA-DRB1\*0301, DQA1\*0501, and DQB1\*0201. Alterations in circulating T cell receptor  $\gamma\delta$  subsets. *Arthr Rheum* 1996;39:1733-9.
81. Lewis HW, Reunala TL, Garioch JS, Leonard JN, Fry JS, Collin P, Evans D, Fry L. Protective effect of gluten-free diet against development of lymphoma in dermatitis herpetiformis. *Br J Dermatol* 1996;135:363-7.
82. Mäki M, Rantala I. Jejunal epithelial cell stress protein expression in gluten-induced enteropathy (celiac disease). In: van Eden W and Young DB, eds. *Stress-proteins in medicine*. Marcel Dekker, New York, 1996;pp. 399-405.
83. Mäki M. Coeliac disease and autoimmunity due to unmasking of cryptic epitopes? *Lancet* 1996;348:1046-47.
84. Mäki M. Keliakian periytyvyys. (Inheritance of coeliac disease, in Finnish.) *Keliakia* 1996;2:5-7.
85. Mäki M. Onko suomalainen kaura vehnätöntä. (Is Finnish oats gluten-free, in Finnish.) *Keliakia* 1996;2:8.
86. Polvi A, Eland C, Koskimies S, Mäki M, Partanen J. HLA DQ and DP in Finnish families with celiac disease. *Eur J Immunogenetics* 1996;23:221-34.
87. Visakorpi JK, Mäki M. Celiakia - zmieniajacy sie obraz kliniczny. *Medycyna Praktyczna* 1996;WS 3:32-35.
88. Collin P, Eerola M, Pirttilä T. Dementia and coeliac disease. In: Gobbi G et al, eds. *Epilepsy and other neurological diseases in coeliac disease*. London: John Libbey & Company Ltd, 1997:285-9.
89. Collin P, Holm K, Kaukinen K, Laurila K, Mäki M. Latentti keliakia. (Latent coeliac disease, in Finnish.) *Suom lääkl*. 1997;52:4279-83.
90. Collin P, Karvonen A-L, Korpela M, Laippala H, Helin H. Gastritis classified according to the Sydney system in patients with primary Sjögren's syndrome. *Scand J Gastroenterol* 1997;32:108-11.
91. Collin P, Mäki M. Celiac disease - even a neurological disorder. *J Pediatr Gastroenterol Nutr* 1997;24:116-7

92. Collin P, Mäki M. Non-gastrointestinal manifestations of coeliac disease. In: Gobbi G et al, eds. Epilepsy and other neurological disorders in coeliac disease. London: John Libbey & Company, 1997:13-18.
93. Collin P, Reunala T, Mäki M. New diagnostic strategy to find coeliac disease. In: Mäki M, Collin P, Visakorpi JK, eds. Coeliac disease. Proceedings of the Seventh International Symposium on Coeliac Disease. Coeliac Disease Study Group, University of Tampere. Tampere 1997;47-51.
94. Collin P, Reunala T, Rasmussen M, Kyrönpalo S, Pehkonen E, Laippala P, Mäki M. High incidence and prevalence of adult celiac disease: Augmented diagnostic approach. *Scand J Gastroenterol.* 1997;32:1129-33.
95. Collin P, Julkunen R, Lehtola J, Mäki M, Rasmussen M, Reunala T, Savilahti E, Uusitupa M, Vuoristo M. Keliakian hoitosuositus. (Diagnosis and treatment recommendations of coeliac disease, in Finnish.) *Duodecim* 1997;113:82-7.
96. Marttinen A, Sulkanen S, Mäki M. Human fibroblast-derived molecules as antigens in enzyme-linked immunosorbent assay for celiac disease-specific IgA. *Eur J Clin Invest* 1997;27:135-40.
97. Mäki M, Collin P. Coeliac disease. *Lancet* 1997;349:1755-9.
98. Mäki M, Laurila K. Keliakia. *Moodi* 1997;21:247-50.
99. Mäki M. Autoantibodies and autoantigens. In: Mäki M, Collin P, Visakorpi JK, eds. Coeliac disease. Proceedings of the Seventh International Symposium on Coeliac Disease. Coeliac Disease Study Group, University of Tampere. Tampere 1997;155-62.
100. Mäki M. Immunopathogenesis of coeliac disease. *Int J Immunopath Pharmacol* 1997;10:81-2.
101. Mäki M. Keliakia ja lehmänmaidon sietämättömyys. (Coeliac disease and cow's milk intolerance, in Finnish). Kirjassa: Simell O, toim. Neuvolakirja. Orion, Vammala, 1997:447-53.
102. Mäki M. Keliakian hoito puhuttaa lähitulevaisuudessa - vehnätrökkelystä vai ei. Pääkirjoitus. (Leading article, in Finnish). *Keliakia* 1997;4:3.
103. Mäki M. Puheenjohtajan palsta. Pääkirjoitus. (Leading article, in Finnish.) *Keliakia* 1997;1:3.
104. Mäki M. Tissue transglutaminase as the autoantigen of coeliac disease. *Gut* 1997;41:565-6.
105. Patinen P, Savilahti E, Hietanen J, Malmström M, Mäki M, Reunala T. Intraepithelial lymphocytes bearing the  $\gamma/\delta$  receptor in the oral and jejunal mucosa in patients with dermatitis herpetiformis. *Eur J Oral Sci* 1997;105:130-5.
106. Polvi A, Garden OA, Elwood CM, Sørensen SM, Batt RM, Mäki M, Partanen J. Canine major histocompatibility complex genes DQA and DQB in Irish setter dogs. *Tissue Antigens* 1997;49:236-43.
107. Polvi A, Mäki M, Partanen J. Celiac patients predominantly inherit HLA-DPB1\*0101 positive haplotype from HLA-DQ2 homozygous parent. *Human Immunol* 1997;53:156-8.
108. Reunala T, Collin P, Lewis, HM, Fry L. Associated diseases and malignancy in dermatitis herpetiformis. In: Mäki M, Collin P, Visakorpi JK, eds. Coeliac disease. Proceedings of the Seventh International Symposium on Coeliac Disease. Tampere: Coeliac Disease Study Group, University of Tampere; Tampere 1997:75-9.
109. Reunala T, Collin P. Diseases associated with dermatitis herpetiformis. *Br J Dermatol.* 1997;136:315-8.
110. Savilahti E, Mäki M. Lasten toistuvat vatsakivut. *Duodecim* 1997;113:589-93.
111. Savilahti E, Örmälä T, Arato A, Hacesek G, Holm K, Klemola T, Nemeth A, Mäki M, Reunala T. Density of  $\gamma/\delta$ + T cells in the jejunal epithelium of patients with coeliac disease and dermatitis herpetiformis is increased with age. *Clin Exp Immunol* 1997;109:464-7.
112. Ashorn M, Mäki M, Ruuska T. Lasten toistuvat vatsavaivat. *Ferring Lääkkeet Oy: Espoo;* 1998:1-23.
113. Collin P, Kaukinen K. Associated disorders in coeliac disease. In: Lohiniemi S, Collin P, Mäki M, eds. Changing Features of Coeliac Disease. The Finnish Coeliac Society. Tampere 1998;89-91.
114. Collin P, Mäki M. Ohutsuoli. Keliakia. (Coeliac disease, in Finnish.) In: Höckerstedt K, Färkkilä M, Kivilaakso E, Pikkarainen P, eds. Gastroenterologia. *Duodecim, Jyväskylä* 1998;265-74.
115. Collin P, Pikkarainen P. Ohutsuoli. Laktaasin ja muiden disakkaridaasien puutostilat. In: Höckerstedt K, Färkkilä M, Kivilaakso E, Pikkarainen P, toim. Gastroenterologia. *Duodecim, Jyväskylä* 1998;285-88.
116. Collin P. Ohutsuoli. Eosinofiilinen gastroenteriitti. In: Höckerstedt K, Färkkilä M, Kivilaakso E, Pikkarainen P, toim. Gastroenterologia. *Duodecim, Jyväskylä* 1998;301.
117. Collin P. Ohutsuoli. Rakenne ja toiminta. In: Höckerstedt K, Färkkilä M, Kivilaakso E, Pikkarainen P, eds. Gastroenterologia. *Duodecim, Jyväskylä* 1998;254-58.
118. Collin P. Serologic screening for coeliac disease - time for tissue transglutaminase test? *Ital J Gastroenterol Hepatol* 1998;30:498-9.
119. Holm K. Markers of coeliac disease latency. In: Lohiniemi S, Collin P, Mäki M, eds. Changing Features of Coeliac Disease. The Finnish Coeliac Society. Tampere 1998;19-24.
120. Jokinen J, Peters U, Mäki M, Miettinen A, Collin P. Coeliac sprue in patients with chronic oral mucosal symptoms. *J Clin Gastroenterol* 1998;26:23-6.
121. Kaukinen K, Collin P, Holm K, Karvonen A-L, Pikkarainen P, Mäki M. Small-bowel mucosal inflammation in reticulin or gliadin antibody-positive patients without villous atrophy. *Scand J Gastroenterol* 1998;33:944-9.
122. Kerttula TO, Holm K, Partanen J, Polvi A, Mäki M. Circulating T lymphocyte subsets in coeliac disease (CoD) patients and healthy family members. *Clin Exp Immunol* 1998;111:536-40.
123. Lähteenoja H, Toivanen A, Viander M, Mäki M, Irjala K, Rähä I, Syrjänen S. Oral mucosal changes in coeliac patients on a gluten-free diet. *Eur J Oral Sci* 1998;106:899-906.
124. Lohiniemi S, Mustalahti K, Collin P, Mäki M. Measuring quality of life in coeliac disease patients. In: Lohiniemi S, Collin P, Mäki M, eds. Changing Features of Coeliac Disease. The Finnish Coeliac Society. Tampere 1998;73-7.
125. Mäki M, Collin P. Coeliac disease. Reply. *Lancet* 1998;351:63.

126. Mäki M, Collin P. Enfermedad Celiaca. (In Spanish). *Tribuna Celiaca* 1998;6-14.
127. Mäki M, Kaukinen K, Holm K, Collin P. Treatment of coeliac patients with oats and wheat starch. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;93-6.
128. Mäki M, Sulkanen S, Collin P. Antibodies in relation to gluten intake. *Digestive Diseases* 1998;16:330-2.
129. Mäki M. Autoimmunità, transglutaminasi tissutale e malattie celiaca. *Medico e Bambino* 1998;17:7.
130. Mäki M. Celiac disease. *Pediatrics Review Series. Food Allergy and Gastroenterology* 1998;2:2-5.
131. Mäki M. Changing features of coeliac disease. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;1-6.
132. Mäki M. Coeliac disease in Finland. In: Banchini G, Donadio A, eds. *La Malattia Celiaca (alla ricerca del tempo perduto)*. 8 Incontro Internazionale "Nuove Technologie in Pediatria". Divisione di Pediatria Arcispedale S. Maria Nuova, Associazione Italiana Celiachia. Reggio Emilia 1998; 43-8.
133. Mäki M. Keliakia tunnetuksi. Pääkirjoitus (Leading article, in Finnish). *Keliakia* 1998;4:3.
134. Mäki M. Keliakia yleistyy ja muuttaa muotoaan. *Keliakia* 1998;4:12.
135. Mäki M. Keliakiaseulonnan haitat ja edut punnittava. *Keliakia* 1998;4:13.
136. Mäki M. Ohutsuoli. Immunologia. (Immunology, in Finnish). In: Höckerstedt K, Färkkilä M, Kivilaakso E, Pikkarainen P, eds. *Gastroenterologia. Duodecim, Jyväskylä* 1998;263-4.
137. Mäki M. To screen or not to screen for coeliac disease. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;51-3.
138. Not T, Horvath K, Hill ID, Partanen J, Hammed A, Magazzu G, Fasano A. Celiac disease risk in the USA: High prevalence of antiendomysium antibodies in healthy blood donors. *Scand J Gastroenterol* 1998;33:494-8.
139. Polvi A, Arranz E, Fernandez-Arquero M, Collin P, Mäki M, Sanz A, Calvo C, Maluenda C, Westman P, de la Concha EG, Partanen J. HLA-DQ2-negative celiac disease in Finland and Spain. *Human Immunol* 1998;59:169-75.
140. Polvi A, Garden OA, Houlston RS, Mäki M, Batt RM, Partanen J. Genetic susceptibility to gluten sensitive enteropathy in Irish setter dogs is not linked to the major histocompatibility complex. *Tissue Antigens* 1998;52:543-9.
141. Polvi A, Mäki M, Collin P, Partanen J. TNF microsatellite alleles  $\alpha 2$  and  $\beta 3$  are not primarily associated with celiac disease in the Finnish population. *Tissue Antigens* 1998;51:553-5.
142. Reunala T, Collin P, Holm K, Pikkarainen P, Miettinen A, Vuolteenaho N, Mäki M. Tolerance to oats in dermatitis herpetiformis. *Gut* 1998;43:490-3.
143. Reunala T. Dermatitis herpetiformis: from gut to skin. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;13-17.
144. Ruuska T, Ashorn M, Mäki M. Suoliston, maksan ja haiman sairaudet. In: Raivio K, Siimes MA. *Lastentaudit. Duodecim, Jyväskylä* 1998:343-53.
145. Scott HS, Heino M, Peterson P, Mittaz L, Lalioti MD, Betterle C, Cohen A, Seri M, Lerone M, Romeo G, Collin P, Salo M, Metcalfe R, Weetman A, Papanavvas M-P, Rossier C, Nagamine K, Kudoh J, Shimizu N, Krohn KJE, Antonarakis SE. Common mutations in autoimmune polyendocrinopathy-candidiasis -ectodermal dystrophy patients of different origins. *Molec Endocrinol* 1998;12:1112-9.
146. Sulkanen S, Collin P, Laurila K, Mäki M. IgA- and IgG-class antihuman umbilical cord antibody tests in adult coeliac disease. *Scand J Gastroenterol* 1998;33:251-4.
147. Sulkanen S, Halttunen T, Laurila K, Kolho K-L, Korponay-Szabo IR, Sarnesto A, Savilahti E, Collin P, Mäki M. Tissue transglutaminase autoantibody enzyme-linked immunosorbent assay in detecting celiac disease. *Gastroenterology* 1998;115:1322-8.
148. Sulkanen S, Halttunen T, Marttinen A, Leivo E-L, Laurila K, Mäki M. Autoantibodies in celiac disease. Importance of fibroblasts. *J Pediatr Gastroenterol Nutr* 1998;27:206-13.
149. Collin P, Kaukinen K, Mäki M. Clinical features of celiac disease today. *Digestive Diseases* 1999;17:100-6.
150. Collin P. New diagnostic findings in coeliac disease. *Ann Med* 1999;31:399-405.
151. Halttunen T, Mäki M. Serum immunoglobulin A from patients with celiac disease inhibits human T84 intestinal crypt epithelial cell differentiation. *Gastroenterology* 1999;116:566-72.
152. Holopainen P, Arvas M, Sistonen P, Mustalahti K, Collin P, Mäki M, Partanen J. CD28/CTLA4 gene region on chromosome 2q33 confers genetic susceptibility to celiac disease. A linkage and family-based association study. *Tissue Antigens* 1999;53:470-5.
153. Iltanen S, Collin P, Korpela M, Holm K, Partanen J, Polvi A, Mäki M. Celiac disease and markers of celiac disease latency in patients with primary Sjögren's syndrome. *Am J Gastroenterol* 1999;94:1042-6.
154. Iltanen S, Holm K, Ashorn M, Ruuska T, Laippala P, Mäki M. Changing jejunal  $\gamma\delta$  T cell receptor (TCR)-bearing intraepithelial lymphocyte density in coeliac disease. *Clin Exp Immunol* 1999;117:51-5.
155. Iltanen S, Holm K, Partanen J, Laippala P, Mäki M. Increased density of jejunal  $\gamma\delta$  T cells in patients having normal mucosa - marker of operative autoimmune mechanisms? *Autoimmunity* 1999;29:179-87.
156. Iltanen S, Rantala I, Laippala P, Holm K, Partanen J, Mäki M. Expression of HSP-65 in jejunal epithelial cells in patients clinically suspected of coeliac disease. *Autoimmunity* 1999;31:125-32.
157. Kaukinen K, Collin P, Holm K, Rantala I, Vuolteenaho N, Reunala T, Mäki M. Wheat starch-containing gluten-free flour products in the treatment of coeliac disease and dermatitis herpetiformis. A long-term follow-up study. *Scand J Gastroenterol*, 1999;34:163-9.

158. Kaukinen K, Collin P, Mykkänen A-H, Partanen J, Mäki M, Salmi J. Celiac disease and autoimmune endocrinologic disorders. *Dig Dis Sci* 1999;44:1428-33.
159. Kaukinen K, Salmi J, Lahtela J, Siljamäki-Ojansuu U, Koivisto A-M, Oksa H, Collin P. No effect of gluten-free diet on the metabolic control of type 1 diabetes in patients with diabetes and celiac disease. *Diabetes Care* 1999; 22:1747-8.
160. Kerttula TO, Collin P, Mäki M, Hurme M. Normal T-helper 1/T-helper 2 balance in peripheral blood of coeliac disease patients. *Scand J Immunol* 1999;49:197-202.
161. Luostarinen L, Pirttilä T, Collin P. Coeliac disease presenting with neurological disorders. *Eur Neurol* 1999;42:132-5.
162. Mäki M. Is coeliac disease a syndrome? UEGW Roma'99. ESPGHAN Postgraduate Course "Gastrointestinal uncertainties. 1999, p. 20-22.
163. [Mustalahti K, Collin P, Sievänen H, Salmi J, Mäki M. Osteopenia in patients with clinically silent coeliac disease warrants screening. \*Lancet\* 1999;354:744-5.](#)
164. Rantala I, Collin P, Holm K, Kainulainen H, Mustonen J, Mäki M. Small bowel T-cells, HLA-DR antigen, and GroEL stress protein in IgA nephropathy. *Kidney Int, in press-Int* 1999;55:2274-80.
165. Collin P, Kaukinen K. Clinical news: the occurrence and treatment of coeliac disease. In: Auricchio S, Greco L, Maiuri L, Troncone R, eds. *Proceedings of the Eighth International Symposium on Coeliac Disease, April 21-24, 1999, Naples, Italy.* JGC Editions, Naples, 2000;119-23.
166. Collin P, Mäki M. Ohutsuoli. Keliakia. (Coeliac disease, in Finnish.) In: Höckerstedt K, Färkkilä M, Kivilaakso E, Pikkarainen P, eds. *Gastroenterologia. Duodecim, Jyväskylä* 2000;266-75.
167. Hervonen K, Karell K, Holopainen P, Collin P, Partanen J, Reunala T. Concordance of dermatitis herpetiformis and celiac disease in monozygous twins. *J Invest Dermatol* 2000;115:990-3.
168. Janatuinen E, Kempainen T, Pikkarainen P, Holm K, Kosma V-M, Uusitupa M, Mäki M, Julkunen R. Lack of cellular and humoral immunological responses to oats in adults with coeliac disease. *Gut* 2000;46:327-31.
169. Kaukinen K, Salmi J, Lahtela J, Oksa H, Collin P. Effect of gluten-free diet on the metabolic control of type I diabetes in patients with diabetes and celiac disease. *Diabetes Care* 2000;23:713.
170. Kaukinen K, Turjanmaa K, Mäki M, Partanen J, Venäläinen R, Reunala T, Collin P. Intolerance to cereals is not specific for coeliac disease. *Scand J Gastroenterol* 2000;35:942-6.
171. Korponay-Szabo I, Sulkanen S, Halttunen T, Maurano F, Rossi M, Mazzarella G, Laurila K, Troncone R, Mäki M. Tissue transglutaminase is the target in both rodent and primate tissues for celiac disease-specific autoantibodies. *J Pediatr Gastroenterol Nutr* 2000;31:520-7.
172. Lähteenoja H, Mäki M, Viander M, Riihinen I, Vilja P, Rantala I, Toivanen A, Syrjänen S. Local challenge on oral mucosa with an  $\alpha$ -gliadin related synthetic peptide in patients with celiac disease. *Am J Gastroenterol* 2000;95:2880-7.
173. Lähteenoja H, Mäki M, Viander M, Toivanen A, Syrjänen S. Local challenge of oral mucosa with gliadin in patients with coeliac disease. *Clin Exp Immunol* 2000;120:38-45.
174. Lähteenoja H, Toivanen A, Viander M, Riihinen I, Rantala I, Syrjänen S, Mäki M. Increase in T cell subsets of oral mucosa: a late immune response in patients with treated coeliac disease? *Scand J Immunol* 2000;52:602-8.
175. Lindfors K, Halttunen T, Huotari P, Nupponen N, Vihinen M, Visakorpi T, Mäki M, Kainulainen H. Identification of novel transcription factor-like gene from human intestinal cells. *EMBL Nucleotide Sequence Database accession number AJ249940. Biochem Biophys Res Com* 2000;276:660-6.
176. Lohiniemi S, Mäki M, Kaukinen K, Laippala P, Collin P. Gastrointestinal symptoms rating scale in coeliac disease patients on wheat starch-based gluten-free diets. *Scand J Gastroenterol* 2000;35:947-9.
177. Mäki M. Autoantibodies and autoantigens in coeliac disease. In: Auricchio S, Greco L, Maiuri L, Troncone R, eds. *Proceedings of the Eighth International Symposium on Coeliac Disease, April 21-24, 1999, Naples, Italy.* JGC Editions, Naples, 2000;57-63.
178. Mäki M. Keliakian diagnostiikka tänään. *Pharmacia Now* No. 4/2000.
179. Mäki M. Ohutsuoli. Immunologia. (Immunology, in Finnish.) In: Höckerstedt K, Färkkilä M, Kivilaakso E, Pikkarainen P, eds. *Gastroenterologia. Duodecim, Jyväskylä* 2000;264-5.
180. Partanen J. The HLA-DRB4 gene does not explain genetic susceptibility in HLA-DQ2-negative celiac disease. *Immunogenetics* 2000;51:249-50.
181. Stern M, Working Group on Serologic Screening for Celiac Disease (Andersson C, Bürgin-Wolff A, Grodzinsky E, Kapitza S, Laurila K, Lüdtke R, Mäki M, Mascart F, Neunhoffer E, Ocmant A, Savilahti E, Volta U, Wäscher I, Wruk M, Zalewski T). Comparative evaluation of serologic tests for celiac disease: a European initiative toward standardization. *J Pediatr Gastroenterol Nutr* 2000;31:513-9.
182. Bourgain C, Genin E, Holopainen P, Mustalahti K, Mäki M, Partanen J, Clerget-Darpoux F. Use of closely related affected individuals for the genetic study of complex diseases in founder populations. *Am J Hum Genet*, 2001;68:154-9.
183. Collin P, Hakanen M, Salmi J, Mäki M, Kaukinen K. Autoimmune hypopituitarism in patients with coeliac disease - symptoms confusingly similar. *Scand J Gastroenterol* 2001;36:558-60.
184. Fabiani E, Catassi C and the International Working Group on Eu-tTG (incl. Sulkanen S, Mäki M). The serum IgA class anti-tissue transglutaminase antibodies in the diagnosis and follow up of coeliac disease. Results of an international multi-centre study. *Eur J Gastroenterol Hepatol* 2001;13:659-65.
185. Hakanen M, Luotola K, Salmi J, Laippala P, Kaukinen K, Collin P. Clinical and subclinical autoimmune thyroid disease in adult celiac disease. *Dig Dis Sci* 2001;46:2631-5.

186. Halttunen T, Mäki M. The potential role of disease specific IgA autoantibodies in coeliac disease pathogenesis. In: Mäki M, Tossavainen M, eds. Proceedings of the Workshop on Transglutaminases, Protein Cross-Linking and Coeliac Disease, September 14-15, 2001, Tampere, Finland. Coeliac Disease Study Group, University of Tampere. Tampere, 2001;149-56.
187. Holopainen P, Mustalahti K, Uimari P, Collin P, Mäki M, Partanen J. Candidate gene regions and genetic heterogeneity in gluten sensitivity. *Gut* 2001;48:696-701.
188. Kaukinen K, Mäki M, Collin P. Compliance to a stright gluten-free diet : Are dietary transgressions detrimental? *Food Allergy Intolerance* 2001;15-27.
189. Kaukinen K, Mäki M, Collin P. Physiopathology of celiac disease. In: Morteau O, ed. The intestinal mucosa and dietary antigens/Oral Tolerance. Landes Bioscience, Georgetown, 2001;Chapter 10:1-13.
190. Kaukinen K, Mäki M, Partanen J, Sievänen H, Collin P. Coeliac disease without villous atrophy: revision of criteria called for. *Dig Dis Sci* 2001;46:879-87.
191. Kokkonen J, Haapalahti M, Laurila K, Karttunen T, Mäki M. Cow's milk protein-sensitive enteropathy at school age. *J Pediatr* 2001;139:797-803.
192. Korponay-Szabo I, Dahlbom I, Mäki M. Recognition of transglutaminase 2 by coeliac autoantibodies - importance of TG2 conformations and cofactors. In: Mäki M, Tossavainen M, eds. Proceedings of the Workshop on Transglutaminases, Protein Cross-Linking and Coeliac Disease, September 14-15, 2001, Tampere, Finland. Coeliac Disease Study Group, University of Tampere. Tampere, 2001;123-33.
193. Lindfors K, Halttunen T, Kainulainen H, Mäki M. Differentially expressed CC3/TIP30 and rab11 along in vivo and in vitro intestinal epithelial cell crypt-villus axis. *Life Sciences* 2001;69:1363-72.
194. Luostarinen L, Collin P, Peräaho M, Mäki M, Pirttilä T. Coeliac disease in patients with cerebellar ataxia of unknown origin. *Ann Med* 2001;33:445-9.
195. Luostarinen L, Dastidar P, Collin P, Peraaho M, Maki M, Erila T, Pirttila T. Association between coeliac disease, epilepsy and brain atrophy. *Eur Neurol* 2001;46:187-91.
196. Mäki M. Celiac disease. In: Delvin EE, Lentze MJ, eds. Gastrointestinal functions. Nestlé Nutrition Workshop Series, Pediatric Program Vol. 46. Vevey 2001;257-74.
197. Mäki M. Coeliac disease, gluten-induced disease entities, clinical aspects. In: Mäki M, Tossavainen M, eds. Proceedings of the Workshop on Transglutaminases, Protein Cross-Linking and Coeliac Disease, September 14-15, 2001, Tampere, Finland. Coeliac Disease Study Group, University of Tampere. Tampere, 2001;1-5.
198. Mäki M. Minne menet, keliakia? *Keliakia* 2001;4:17-18.
199. Susi M, Holopainen P, Mustalahti K, Mäki M, Partanen J. Candidate gene region 15q26 and genetic susceptibility to coeliac disease in Finnish families. *Scand J Gastroenterol* 2001;36:372-4.
200. Arvola T, Mustalahti K, Saha M-T, Vehmanen P, Partanen J, Ashorn M. Celiac disease, thyrotoxicosis, and autoimmune hepatitis in a child. *J Pediatr Gastroenterol Nutr* 2002;35:90-2.
201. Collin P, Kaukinen K, Mäki M, Vehmainen P, Partanen J, Färkkilä M, Halme L, Höckerstedt K. Discussion on celiac disease in patients with severe liver disease: Gluten-free diet may reverse hepatic failure. Reply. *Gastroenterology* 2002;123:2159-60.
202. Collin P, Kaukinen K, Välimäki M, Salmi J. Endocrinological disorders and celiac disease. *Endocrine Reviews* 2002;23:464-83.
203. Collin P, Rasmussen M, Kyrönpalo S, Laippala P, Kaukinen K. The hunt for coeliac disease in primary care. *Q J Med* 2002;95:75-7.
204. Collin P, Syrjänen J, Partanen J, Pasternack A, Kaukinen K, Mustonen J. Celiac disease and HLA DQ in patients with IgA nephropathy. *Am J Gastroenterol*, 2002;97:2572-6.
205. Hill I, Bhatnagar S, Cameron DJ, De Rosa S, Mäki M, Russell GJ, Troncone R. Report of the Celiac Disease: Working Group Report of the First World Congress of Pediatric Gastroenterology, Hepatology and Nutrition. *J Pediatr Gastroenterol Nutr* 2002;35,Suppl2:S78-88.
206. Janatuinen EK, Kempainen TA, Julkunen RJK, Kosma V-M, Mäki M, Heikkinen M, Uusitupa MIJ. No harm from five year ingestion of oats in coeliac disease. *Gut* 2002;50:332-5.
207. Kainulainen H, Rantala I, Collin P, Ruuska T, Päivärinne H, Halttunen T, Lindfors K, Kaukinen K, Mäki M. Blisters in the small intestinal mucosa of coeliac patients contain T cells positive for cyclooxygenase-2. *Gut* 2002;50:84-9.
208. Karell K, Holopainen P, Mustalahti K, Collin P, Mäki M, Partanen J. Not all HLA DR3 DQ2 haplotypes confer equal susceptibility to coeliac disease: transmission analysis in families. *Scand J Gastroenterol* 2002;37:56-61.
209. Karell K, Korponay-Szabo I, Szalai Zs, Holopainen P, Mustalahti K, Collin P, Mäki M, Partanen J. Genetic dissection between coeliac disease and dermatitis herpetiformis in sib pairs. *Ann Hum Genet* 2002;66:387-92.
210. Kaukinen K, Collin P. Oats and wheat starch-based gluten-free products: two contentious diets in the treatment of coeliac disease. 10th International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002, Lectures. Paris 2002.
211. Kaukinen K, Halme L, Collin P, Färkkilä M, Mäki M, Vehmanen P, Partanen J, Höckerstedt K. Celiac disease in patients with severe liver disease: gluten-free diet may reverse hepatic failure. *Gastroenterology* 2002;122:881-8.
212. Kaukinen K, Partanen J, Mäki M, Collin P. HLA-DQ typing in the diagnosis of celiac disease. *Am J Gastroenterol* 2002;97:695-9.
213. Kaukinen K, Sulkanen S, Mäki M, Collin P. IgA-class transglutaminase antibodies in evaluating the efficacy of gluten-free diet in coeliac disease. *Eur J Gastroenterol Hepatol* 2002;14:311-5.

214. Kokkonen J, Ruuska T, Karttunen TJ, Mäki M. Lymphonodular hyperplasia of the terminal ileum associated with colitis shows an increased  $\gamma\delta^+$  T-cell density in children. *Am J Gastroenterol* 2002;97:667-72.
215. Liu J, Juo S-H, Holopainen P, Terwilliger J, Tong X, Grunn A, Brito M, Green P, Mustalahti K, Mäki M, Gilliam TC, Partanen J. Genomewide linkage analysis of celiac disease in Finnish families. *Am J Hum Genet* 2002;70:51-9.
216. Mustalahti K, Holopainen P, Karell K, Mäki M, Partanen J. Genetic dissection between silent and clinically diagnosed symptomatic forms of coeliac disease in multiplex families. *Digest Liver Dis* 2002;34:842-5.
217. Mustalahti K, Lohiniemi S, Collin P, Vuolteenaho N, Laippala P, Mäki M. Gluten-free diet and quality of life in patients with screen-detected celiac disease. *Effective Clinical Practice* 2002;5:105-13.
218. Mustalahti K, Sulkanen S, Holopainen P, Laurila K, Collin P, Partanen J, Mäki M. Coeliac disease among healthy members of multiple case coeliac disease families. *Scand J Gastroenterol* 2002;37:161-5.
219. Woolley N, Holopainen P, Bourgain C, Mustalahti K, Collin P, Mäki M, Partanen J. CD80 (B7-1) and CD86 (B7-2) genes and genetic susceptibility to coeliac disease. *Eur J Immunogenet* 2002;29:331-3.
220. Woolley N, Holopainen P, Ollikainen V, Mustalahti K, Mäki M, Kere J, Partanen J. A new locus for coeliac disease mapped to chromosome 15 in a population isolate. *Human Genetics* 2002;111:40-5.
221. Babron M-C, Nilsson S, Adamovic S, Torinsson Nalwai Å, Wahlström J, Ascher H, Ciclitira PJ, Sollid LM, Partanen J, Greco L, Clerget-Darpoux F, and the European Genetics Cluster on Coeliac Disease (incl. Holopainen P, Karell K, Partanen J, Collin P, Mustalahti K, Mäki M). Meta and pooled analysis of European coeliac disease data. *Eur J Hum Genet* 2003;11:828-34.
222. Bolognesi E, Karell K, Percopo S, Coto I, Greco L, Mantovani V, Suoraniemi E, Partanen J, Mustalahti K, Mäki M, Momigliano-Richiardi P. Additional factor in some HLA DR3/DQ2 haplotypes confers a fourfold increased genetic risk of celiac disease. *Tissue Antigens* 2003;61:308-16.
223. Collin P, Kaukinen K. Keliakian diagnosointi. *Suom lääkl* 2003;58:271-5.
224. Collin P, Reunala T. Recognition and management of the cutaneous manifestations of celiac disease. A guide for dermatologists. *Am J Clin Dermatol* 2003;4:13-20.
225. Hausch F, Halttunen T, Mäki M, Khosla C. Design, synthesis and evaluation of gluten peptide analogs as selective inhibitors of human tissue transglutaminase. *Chemistry & Biology* 2003;10:225-31.
226. Heinonen TYK, Pasternack L, Lindfors K, Breton C, Gastinel LN, Mäki M, Kainulainen H. A novel human glycosyltransferase: primary structure and characterisation of the gene and transcripts. *Biochem Biophys Res Com* 2003;309:166-74.
227. Iltanen S, Mäki M, Collin P, Kaukinen K, Partanen J, Hulkkonen J, Hurme M, Aittoniemi J. The association between mannan-binding lectin gene alleles and celiac disease. *Am J Gastroenterol* 2003;98:2808-9.
228. Järvinen TT, Kaukinen K, Laurila K, Kyrönpalo S, Rasmussen M, Mäki M, Korhonen H, Reunala T, Collin P. Intraepithelial lymphocytes in celiac disease. *Am J Gastroenterol* 2003;98:1332-7.
229. Karell K, Louka AS, Moodie SJ, Ascher H, Clot F, Greco L, Ciclitira PJ, Sollid LM, Partanen J and the members of the European Genetics Cluster on Coeliac Disease (incl. Collin P, Mustalahti K, Mäki M). HLA types in coeliac disease patients not carrying the DQA1\*05-DQB1\*02 (DQ2) heterodimer: Results from the European genetics cluster on coeliac disease. *Human Immunology* 2003;64:469-477.
230. Kaukinen K, Collin P, Mäki M, Partanen J, Halme L, Färkkilä M, Höckerstedt K. Screening for celiac disease in patients with chronic liver disease. Reply. *Gastroenterology* 2003;125:1289-90.
231. Korponay-Szabo IR, Dahlbom I, Laurila K, Koskinen S, Woolley N, Partanen J, Kovacs JB, Mäki M, Hansson T. Elevation of IgG antibodies against tissue transglutaminase as a diagnostic tool for coeliac disease in selective IgA deficiency. *Gut* 2003;52:1567-71.
232. Korponay-Szabó IR, Laurila K, Szondy Z, Halttunen T, Szalai Z, Dahlbom I, Rantala I, Kovács JB, Fésüs L, Mäki M. Missing endomysial and reticulin binding of coeliac antibodies in transglutaminase 2 knockout tissues. *Gut* 2003;52:199-204.
233. Lindfors K, Viiri KM, Niittynen M, Heinonen TYK, Mäki M, Kainulainen H. TGF-beta induces the expression of SAP30L, a novel nuclear protein. *BMC Genomics* 2003;4:1-22.
- Louka AS, Moodie SJ, Karell K, Bolognesi E, Ascher H, Greco L, Momigliano-Richiardi P, Partanen J, Ciclitira PJ, Sollid LM, and the European Genetics Cluster on Celiac Disease (incl. Holopainen P, Karell K, Partanen J, Collin P, Mustalahti K, Mäki M). A collaborative European search for non-DQA1\*05-DQB1\*02 celiac disease loci on HLA-DR3 haplotypes: analysis of transmission from homozygous parents. *Human Immunol* 2003;64:350-8.
234. Mäki M, Mustalahti K, Knip M. Screening for celiac disease. *N Engl J Med* 2003;349:1673-4. The author reply.
235. Mäki M, Mustalahti K, Kokkonen J, Kulmala P, Haapalahti M, Karttunen T, Ilonen J, Laurila K, Dahlbom I, Hansson T, Höpfl P, Knip M. Prevalence of celiac disease among children in Finland. *N Engl J Med* 2003;348:2517-24.
236. Mäki M. Strategies for screening for coeliac disease. In: Cerf-Bensussan N, Brousse N, Caillat-Zucman S, Cellier C, Schmitz J, eds. *Coeliac disease. Proceedings of the Xth International Symposium on Coeliac Disease, Paris, June 2-5, 2002*. John Libbey Eurotext. Montrouge 2003;191-8.
237. Peräaho M, Kaukinen K, Paasikivi K, Sievänen H, Lohiniemi S, Mäki M, Collin P. Wheat-starch-based gluten-free products in the treatment of newly detected coeliac disease. Prospective and randomized study. *Aliment Pharmacol Ther* 2003;17:587-94.
238. Collin P, Mäki M, Kaukinen K. Complete small intestinal mucosal recovery is obtainable in the treatment of celiac disease. Letter to the Editor. *Gastrointest Endosc* 2004;59:158-9.

239. Collin P, Maki M, Kaukinen K. It is the compliance, not milligrams of gluten, that is essential in the treatment of celiac disease. Letter to the Editor. *Nutrition Reviews* 2004;62:490.
240. Collin P, Mustalahti K, Kyrönpalo S, Rasmussen M, Pehkonen E, Kaukinen K. Should we screen reflux oesophagitis patients for coeliac disease? *Eur J Gastroenterol Hepatol* 2004;16:917-20.
241. Collin P, Thorell L, Kaukinen K, Mäki M. The safe threshold for gluten contamination in gluten-free products. Can trace amounts be accepted in the treatment of coeliac disease? *Aliment Pharmacol Ther* 2004;19:1277-83.
242. Collin P. Antibodies and biopsy in the diagnosis of coeliac disease: lessons from the Celikey multicentre study. *Elia Journal*, Nov 6-8, 2004.
243. Collin P. Keliakia rasvaripulista yleiseksi autoimmuunitaudiksi. *Suom Lääkäril*, 2004; 45: 4355.
244. Dolinsek J, Urlep D, Karell K, Partanen J, Micetic-Turk D. The prevalence of celiac disease among family members of celiac disease patients. *Wiener Klinische Wochenschrift* 2004; 116: 8-12.
245. Haimila K, Smedberg T, Mustalahti K, Mäki M, Partanen J, Holopainen P. Genetic association of coeliac disease susceptibility to polymorphisms in the ICOS gene on chromosome 2q33. *Genes Immun* 2004;5:85-92.
246. Hervonen K, Viljamaa M, Collin P, Knip M, Reunala T. The occurrence of type 1 diabetes in patients with dermatitis herpetiformis and their first-degree relatives. *Br J Dermatol*, 2004;150:136-8.
247. Holopainen P, Naluai AT, Moodie S, Percopo S, Coto I, Clot F, Ascher H, Sollid L, Ciclitira P, Greco L, Clerget-Darpoux F, Partanen J, and the members of the European Genetics Cluster on Coeliac Disease (incl. Collin P, Mustalahti K, Mäki M). Candidate gene region 2q33 in European families with coeliac disease. *Tissue Antigens* 2004;63:212-22.
248. Järvinen TT, Collin P, Rasmussen M, Kyrönpalo S, Mäki M, Partanen J, Reunala T, Kaukinen K. Villous tip intraepithelial lymphocytes as markers of early-stage coeliac disease. *Scand J Gastroenterol* 2004;39:428-33.
249. Juuti-Uusitalo K, Mäki M, Kaukinen K, Collin P, Visakorpi T, Vihinen M, Kainulainen H. cDNA microarray analysis of gene expression in coeliac disease jejunal biopsy samples. *Journal of Autoimmunity* 2004;22:249-65.
250. Kokkonen J, Holm K, Karttunen TJ, Mäki M. Enhanced local immune response in children with prolonged gastrointestinal symptoms. *Acta Paediatr* 2004;93:1601-7.
251. Korponay-Szabo IR, Halttunen T, Szalai Z, Laurila K, Király R, Kovács JB, Fésüs L, Mäki M. In vivo targeting of intestinal and extraintestinal transglutaminase 2 by coeliac autoantibodies. *Gut* 2004;53:641-8.
252. Lohi O, Mäki M. Vehnää keliaakikolle? *Duodecim* 2004;120:645-6
253. Mäki M, Lohi O. Celiac disease. In: Walker et al, eds. *Pediatric Gastrointestinal Disease. Pathophysiology, diagnosis, management*. 4<sup>th</sup> Edition. BC Decker Inc. Hamilton, 2004;932-43.
254. Patinen P, Aine L, Collin P, Hietanen J, Korpela M, Enckell G, Kautiainen H, Kontinen YT, Reunala T. Oral findings in coeliac disease and Sjögren's syndrome. *Oral Diseases* 2004;10:330-4.
255. Peräaho M, Collin P, Kaukinen K, Kekkonen L, Miettinen S, Mäki M. Oats can diversify a gluten-free diet in celiac disease and dermatitis herpetiformis. *J Am Diet Assoc* 2004;104:1148-50.
256. Peräaho M, Kaukinen K, Mustalahti K, Vuolteenaho N, Mäki M, Laippala P, Collin P. Effect of an oats-containing gluten-free diet on symptoms and quality of life in coeliac disease. A randomised study. *Scand J Gastroenterol* 2004;39:27-31.
257. Peräsaari J, Koskela S, Partanen J. Characterization of a novel HLA-B40 allele with serological Bw4 motif, HLA B4047, in the Finnish population and confirmation of B\*270503 allele. *Tissue Antigens* 2004; 63: 595-7.
258. Troncone R, Bhatnagar S, Butzner D, Cameron D, Hill I, Hoffenberg E, Mäki M, Mendez V, de Jimenez MZ. Celiac disease and other immunologically mediated disorders of the gastrointestinal tract: Working Group Report of the Second World Congress of Pediatric Gastroenterology, Hepatology and Nutrition. *J Pediatr Gastroenterol Nutr* 2004;39:S601-10.
259. Viljamaa M, Kaukinen K, Sievänen H, Collin P. Keliakia myös luuston tauti. *Suom Lääkäril*, 2004; 45: 4357-60.
260. Cellier C, Green PH, Collin P, Murray J. ICCE Consensus for Celiac Disease. *Endoscopy*. 2005;37:1055-9.
261. Collin P, Kaukinen K, Vogelsang H, Korponay-Szabó I, Sommer R, Schreier E, Volta U, Granito A, Veronesi L, Mascart F, Ocmant A, Ivarsson A, Lagerqvist C, Bürgin-Wolff A, Hadziselimovic F, Furlano R I, Sidler M A, Mulder CJJ, Goerres M S, Mearin ML, Ninaber MK, Gudmand-Høyer E, Fabiani E, Catassi C, Tidlund H, Alaintalo L, Mäki M. Antiendomysial and antihuman recombinant tissue transglutaminase antibodies in the diagnosis of coeliac disease. A biopsy-proven European multicentre study. *Eur J Gastroenterol Hepatol* 2005;17:85-91.
262. Collin P. Who should be screened for celiac disease? The benefits and harms of screening. *Gastroenterology* 2005;128:S104-8.
263. Collin P, Wahab PJ, Murray JA. Intraepithelial lymphocytes and coeliac disease. *Best Pract Res Clin Gastroenterol*, 2005;19:341-50.
264. Haapalahti M, Kulmala P, Karttunen TJ, Paajanen L, Laurila K, Mäki M, Mykkänen H, Kokkonen J. Nutritional status in adolescents and young adults with screen-detected celiac disease. *J Pediatr Gastroenterol Nutr* 2005;40:566-70.
265. Hernando A, Garcia E, Llorente M, Mujico JR, Lombadia M, Mäki M, Kaukinen K, Collin P, Mendez E. Measurements of hydrolysed gliadins in malts, breakfast cereals, heated/hydrolysed foods, whiskies and beers by means of a new competitive R5 ELISA. In: Stern M, ed. *Proceedings of the 19th Meeting, Working Group on Prolamin Analysis and Toxicity*. Verl. Wissensch. Scripten. Zwickau 2005;31-7.
266. Hervonen K, Vornanen M, Kautiainen H, Collin P, Reunala T. Lymphoma in patients with dermatitis herpetiformis and their first-degree relatives. *Br J Dermatol*, 2005;152:82-6.

267. Honkanen T, Mustonen J, Kainulainen H, Myllymäki J, Collin P, Hurme M, Rantala I. Small bowel cyclooxygenase 2 (COX-2) expression in patients with IgA nephropathy. *Kidney Int*, 2005; 67:2187-95.
268. Kaukinen K. Tulisiko keliakiaa seuloa väestöstä? Pääkirjoitus. *Duodecim* 2005;121:1361-2.
269. Kaukinen K, Mäki M, Collin P. Trace amounts of gluten in the treatment of coeliac disease. In: Stern M, ed. Proceedings of the 19th Meeting Working Group on Prolamin Analysis and Toxicity. Verl. Wissensch. Scripten. Zwickau 2005;115-6.
270. Kaukinen K, Peräaho M, Collin P, Partanen J, Woolley N, Kaartinen T, Nuutinen T, Halttunen T, Mäki M, Korponay-Szabo I. Small-bowel mucosal transglutaminase 2-specific IgA deposits in coeliac disease without villous atrophy: A prospective and randomized clinical study. *Scand J Gastroenterol* 2005;40:564-72.
271. Korponay-Szabó I, Raivio T, Laurila K, Opre J, Király R, Kovács JB, Kaukinen K, Fésüs L, Mäki M. Celiac disease case finding and diet monitoring by point-of-care testing. *Aliment Pharmacol Therap* 2005;22:729-37.
272. Lähdeaho M-L, Kaukinen K, Collin P, Ruuska T, Partanen J, Haapala A-M, Mäki M. Celiac disease: From inflammation to atrophy: A long-term follow-up study. *J Pediatr Gastroenterol Nutr* 2005;41:44-8.
273. Mujico JR, Hernando A, Lombardia M, Benavides A, Silio V, Mäki M, Kaukinen K, Collin P, Thorell L, Ryöppy P, Mendez E. Quantification of wheat, barley and rye contamination in oat samples by real-time PCR. In: Stern M, ed. Proceedings of the 19th Meeting Working Group on Prolamin Analysis and Toxicity. Verl. Wissensch. Scripten. Zwickau 2005;87-94.
274. Peräaho M, Collin P, Kaukinen K, Mäki M. Contaminated oats and other gluten-free foods in the United States. Authors' response. *J Am Diet Assoc* 2005;105:348-9.
275. Ranua J, Luoma K, Auvinen A, Mäki M, Haapala AM, Peltola J, Raitanen J, Isojärvi J. Celiac disease-related antibodies in an epilepsy cohort and matched reference population. *Epilepsy Behav* 2005;6:388-92.
276. Ranua J, Luoma K, Auvinen A, Haapala AM, Mäki M, Peltola J, Raitanen J, Isojärvi JI. Antimitochondrial antibodies in patients with epilepsy. *Epilepsy Behav* 2005;7:95-7.
277. Viljamaa M, Collin P, Huhtala H, Sievänen H, Mäki M, Kaukinen K. Is coeliac disease screening in risk groups justified? A fourteen-year follow-up with special focus on compliance and quality of life. *Aliment Pharmacol Ther* 2005;22:317-24.
278. Viljamaa M, Kaukinen K, Huhtala H, Kyrönpalo S, Rasmussen M, Collin P. Coeliac disease, autoimmune diseases and gluten exposure. *Scand J Gastroenterol* 2005;40:437-43.
279. Woolley N, Mustalahti K, Mäki M, Partanen J. Cytokine gene polymorphisms and genetic association with coeliac disease in the Finnish population. *Scand J Immunol* 2005;61:51-6.
280. Collin P, Julkunen R, Lehtola J, Kaukinen K, Mäki M, Rasmussen M, Reunala T, Savilahti E, Uusitupa M, Vuoristo M, Vuorio A. Keliakia. Käypä hoito-suosituksen päivitys. *Duodecim* 2005;121:2705-7.
281. Collin P, Kaukinen K, Vogelsang H, Korponay-Szabó I, Sommer R, Schreier E, Volta U, Granito A, Veronesi L, Mascart F, Ocmant A, Ivarsson A, Lagerqvist C, Bürgin-Wolff A, Hadziselimovic F, Furlano R I, Sidler M A, Mulder C J J, Goerres M S, Mearin M L, Ninaber M K, Gudmand-Høyer E, Fabiani E, Catassi C, Tidlund H, Alaintalo L, Mäki M. Antibodies and biopsy in the diagnosis of coeliac disease: Lessons from the Celikey European Multicentre Study. In: McMillan S, Feighery C, Watson P, O'Farrelly C, eds. Coeliac disease. Proceedings of the Eleventh International Symposium on Coeliac Disease, April 28-30, 2004, Belfast, Northern Ireland. Juvela, Semper. UK & Ireland, Nordic Countries 2006;11-4.
282. Collin P, Kaukinen K. Are trace amounts of gluten acceptable in the treatment of coeliac disease? In: Stern M, ed. Proceedings of the 20th Meeting, Working Group on Prolamin Analysis and Toxicity. Verl. Wissensch. Scripten. Zwickau 2006;107-10.
283. Collin P, Reunala T. Jos suoli ei parane? Keliakia ja pahanlaatuiset sairaudet. In: Mäki M et al, eds. Keliakia. Kustannus Oy Duodecim, Jyväskylä 2006;159-63.
284. Collin P, Reunala T. Jos suoli ei parane? Keliakiaan ja ihokeliakiaan liittyvä kuolleisuus. In: Mäki M et al, eds. Keliakia. Kustannus Oy Duodecim, Jyväskylä 2006;163-4.
285. Collin P, Reunala T. Keliakia: harvinaisuus vai kansantauti? Ihokeliakian esiintyvyys. In: Mäki M et al, eds. Keliakia. Kustannus Oy Duodecim, Jyväskylä 2006;78-9.
286. Collin P. Keliakia: harvinaisuus vai kansantauti? Esiintyvyys lapsilla. In: Mäki M et al, eds. Keliakia. Kustannus Oy Duodecim, Jyväskylä 2006;78.
287. Collin P. Keliakia: harvinaisuus vai kansantauti? Esiintyvyys Suomessa. In: Mäki M et al, eds. Keliakia. Kustannus Oy Duodecim, Jyväskylä 2006;77.
288. Collin P. Keliakia: harvinaisuus vai kansantauti? Keliakian yleisyys. In: Mäki M et al, eds. Keliakia. Kustannus Oy Duodecim, Jyväskylä 2006;76.
289. Collin P. Mistä tunnistan keliakian? Liitännäissairaudet. In: Mäki M et al, eds. Keliakia. Kustannus Oy Duodecim, Jyväskylä 2006;40-2.
290. Collin P. Mistä tunnistan keliakian? Suoliston ulkopuoliset oireet. In: Mäki M et al, eds. Keliakia. Kustannus Oy Duodecim, Jyväskylä 2006;37-40.
291. Collin P. Miten keliakia todetaan? Keitä seulotaan? In: Mäki M et al, eds. Keliakia. Kustannus Oy Duodecim, Jyväskylä 2006;67-70.
292. Hadjivassiliou M, Mäki M, Sanders DS, Williamson C, Grünewald RA, Woodroffe N, Korponay-Szabó IR. Autoantibody targeting of brain and intestinal transglutaminase in gluten ataxia. *Neurology* 2006;66:373-7.

293. Heinonen TYK, Peltö-Huikko M, Pasternack L, Mäki M, Kainulainen H. Murine ortholog of the novel glycosyltransferase, B3GTL: primary structure, characterization of the gene and transcripts, and expression in tissues. *DNA Cell Biol* 2006;25:465-74.
294. Hischenhuber C, Crevel R, Jarry B, Mäki M, Moneret-Vautrin DA, Romano A, Troncone R, Ward R. Review article: Safe amounts of gluten for patients with wheat allergy or coeliac disease. *Aliment Pharmacol Ther* 2006;23:559-75.
295. Holm K, Mäki M, Vuolteenaho N, Mustalahti K, Ashorn M, Ruuska T, Kaukinen K. Oats in the treatment of childhood coeliac disease: a 2-year controlled trial and a long-term clinical follow-up study. *Aliment Pharmacol Ther* 2006;23:1463-72.
296. Juuti-Uusitalo K, Kaukinen K, Mäki M, Kainulainen H. Gene expression in TGFbeta induced epithelial cell differentiation in a three-dimensional intestinal epithelial cell differentiation model. *BMC Genomics* 2006;7:279.
297. Kaukinen K, Mäki M, Collin P. Immunohistochemical features in antiendomysium positive patients with normal villous architecture. Letter to the Editor. *Am J Gastroenterol* 2006;101:675-6.
298. Kaukinen K, Mäki M, Collin P. Intraepithelial lymphocytes in the diagnosis of coeliac disease. In: McMillan S, Feighery C, Watson P, O'Farrelly C, eds. *Coeliac disease. Proceedings of the Eleventh International Symposium on Coeliac Disease, April 28-30, 2004, Belfast, Northern Ireland. Juvola, Semper. UK & Ireland, Nordic Countries* 2006;22-5.
299. Kaukinen K, Mäki M, Kekkonen L. Uusi tutkimus etsii vastausta. Kannattaako keliakiaa seuloa väestöstä? *Keliakia* 2006;5:6
300. Kaukinen K. Gluten-free diet: The recommendation of the Codex Alimentarius. In: McMillan S, Feighery C, Watson P, O'Farrelly C, eds. *Coeliac disease. Proceedings of the Eleventh International Symposium on Coeliac Disease, April 28-30, 2004, Belfast, Northern Ireland. Juvola, Semper. UK & Ireland, Nordic Countries* 2006;100-3.
301. Kaukinen K. Hoitona gluteeniton ruokavakio. Vehnätärkkelys keliakian hoidossa. In: Mäki M et al, eds. *Keliakia. Kustannus Oy Duodecim, Jyväskylä* 2006;95-7.
302. Kaukinen K. Miten keliakia todetaan? Diagnostiikan periaatteet. In: Mäki M et al, eds. *Keliakia. Kustannus Oy Duodecim, Jyväskylä* 2006;56-7.
303. Kaukinen K. Miten keliakia todetaan? Keliakia ennen atrofiaa. In: Mäki M et al, eds. *Keliakia. Kustannus Oy Duodecim, Jyväskylä* 2006;60-2.
304. Kaukinen K. Miten keliakia todetaan? Ohutsuolen koepala ja sen tulkinta. In: Mäki M et al, eds. *Keliakia. Kustannus Oy Duodecim, Jyväskylä* 2006;57-60.
305. Korponay-Szabó I, Raivio T, Nemes É, B. Kovács J, Laurila K, Kaukinen K, Mäki M. Coeliakia autoantitestein helyszíni kimutatása Biocard<sup>TM</sup> gyorstesztel (Detection of coeliac disease antibodies by the Biocard rapid test, in Hungarian). *Gyermekgyógyászat* 2006;57:351-357.
306. Korponay-Szabó I. Keliakian pikatesti. *Keliakia* 2006;1:4-7.
307. Kurppa K, Kaukinen K, Mäki M. Kannattaako oireetonta keliakiaa tutkia ja hoitaa? *Keliakia* 2006;1:8-9.
308. Kurppa K, Kaukinen K, Mäki M. Keliakian puhkeamista ja taudinkuvaa selvitetään. Keliakian perinnöllisyystutkimus menossa. *Keliakia* 2006;5:4-5.
309. Kurppa K, Kaukinen K, Mäki M. Uusi tutkimus alkamassa. Keliakian puhkeamiseen vaikuttavia perintötekijöitä etsitään. *Keliakia* 2006;1:8-9.
310. Mäki M. Gluteeniton turvallisesti. Gluteenin mittaaminen elintarvikkeista. In: Mäki M et al, eds. *Keliakia. Kustannus Oy Duodecim, Jyväskylä* 2006;132-3.
311. Mäki M. Keliakia. In: Lehtola J et al., eds. *Suomen Gastroenterologiyhdistys 50 vuotta. Suomen Gastroenterologiyhdistys ry, Tampere* 2006:73-5.
312. Mäki M. Kuinka keliakia syntyy? Gluteeni. In: Mäki M et al, eds. *Keliakia. Kustannus Oy Duodecim, Jyväskylä* 2006;18-20.
313. Mäki M. Kuinka keliakia syntyy? In: Mäki M et al, eds. *Keliakia. Kustannus Oy Duodecim, Jyväskylä* 2006;17-24.
314. Mäki M. Kuinka keliakia syntyy? Perimä. In: Mäki M et al, eds. *Keliakia. Kustannus Oy Duodecim, Jyväskylä* 2006;20-1.
315. Mäki M. Kuinka keliakia syntyy? Suolen limakalvolla tapahtuu. In: Mäki M et al, eds. *Keliakia. Kustannus Oy Duodecim, Jyväskylä* 2006;21-34.
316. Mäki M. Minne menet keliakia? In: Suomen Gastroenterologiyhdistys ry. 50-vuotisjuhlakokous. Gastroenterologian saavutuksia ja tulevaisuuden näkymiä. Luennot, Turku 7.-8.9.2006;12-14.
317. Mäki M. Minne menet keliakia? Keliakia huomenna. In: Mäki M et al, eds. *Keliakia. Kustannus Oy Duodecim, Jyväskylä* 2006;167-70.
318. Mäki M. Miten keliakia todetaan? Vasta-ainetestit diagnostiikassa. In: Mäki M et al, eds. *Keliakia. Kustannus Oy Duodecim, Jyväskylä* 2006;62-6.
319. Mearin ML, Catassi C, Brousse N, Brand R, Collin P, Fabiani E, Schweizer JJ, Abuzakouk M, Szajewska H, Hallert C, Farre Masip C, Holmes GKT; on behalf of the Biomed Study Group on Coeliac Disease and Non-Hodgkin Lymphoma. European multi-centre study on coeliac disease and non-Hodgkin lymphoma. (Biomed Study Group on Coeliac Disease and Non-Hodgkin Lymphoma, incl. Kaukinen K). *Eur J Gastroenterol Hepatol* 2006;18:187-94.
320. Metzger M-H, Heier M, Mäki M, Bravi E, Schneider A, Loewel H, Illig T, Schuppan D, Wichmann H-E. Mortality excess in individuals with elevated IgA anti-transglutaminase antibodies: The KORA/MONICA Augsburg cohort study 1989-1998. *Eur J Epidemiol* 2006;21:359-65.

321. Raivio T, Kaukinen K, Nemes E, Laurila K, Collin P, Kovacs JB, Mäki M, Korponay-Szabo IR. Self transglutaminase-based rapid coeliac disease antibody detection by a lateral flow method. *Aliment Pharmacol Ther* 2006;24:147-54.
322. Salmi T, Collin P, Korponay-Szabo IR, Laurila K, Partanen J, Huhtala H, Kiraly R, Lorand L, Reunala T, Mäki M, Kaukinen K. Endomysial antibody-negative coeliac disease: clinical characteristics and intestinal autoantibody deposits. *Gut* 2006;55:1746-53.
323. Salmi TT, Collin P, Järvinen O, Haimila K, Partanen J, Laurila K, Korponay-Szabo IR, Huhtala H, Reunala T, Mäki M, Kaukinen K. Immunoglobulin A autoantibodies against transglutaminase 2 in the small intestinal mucosa predict forthcoming coeliac disease. *Aliment Pharmacol Ther* 2006;24:541-52.
324. Viiri KM, Korkeamäki H, Kukkonen MK, Nieminen LK, Lindfors K, Peterson P, Mäki M, Kainulainen H, Lohi O. SAP30L interacts with members of the Sin3A corepressor complex and targets Sin3A to the nucleolus. *Nucleic Acids Research* 2006;34:3288-98.
325. Viljamaa M, Kaukinen K, Pukkala E, Hervonen K, Reunala T, Collin P. Malignancies and mortality in patients with coeliac disease and dermatitis herpetiformis: 30-year population-based study. *Digest Liver Dis* 2006;38:374-80.
326. Visakorpi J. Huolellinen seuranta varmentaa tuloksen. Keliakian seuranta lapsilla. In: Mäki M et al, eds. *Keliakia*. Kustannus Oy Duodecim, Jyväskylä 2006;128-9.
327. Visakorpi J. Jo antiikin kreikkalaiset sairastivat keliakiaa! Dicke löytää vahingollisen tekijän. In: Mäki M et al, eds. *Keliakia*. Kustannus Oy Duodecim, Jyväskylä 2006;11-2.
328. Visakorpi J. Jo antiikin kreikkalaiset sairastivat keliakiaa! Keliakian historiasta. In: Mäki M et al, eds. *Keliakia*. Kustannus Oy Duodecim, Jyväskylä 2006;10-11.
329. Visakorpi J. Jo antiikin kreikkalaiset sairastivat keliakiaa! Keliakian diagnostiset kriteerit. In: Mäki M et al, eds. *Keliakia*. Kustannus Oy Duodecim, Jyväskylä 2006;14.
330. Visakorpi J. Jo antiikin kreikkalaiset sairastivat keliakiaa! Suolibiopsia paljastaa taudin peruslöydöksen. In: Mäki M et al, eds. *Keliakia*. Kustannus Oy Duodecim, Jyväskylä 2006;12-4.
331. Visakorpi J. Jo antiikin kreikkalaiset sairastivat keliakiaa! Keliakia Suomessa. In: Mäki M et al, eds. *Keliakia*. Kustannus Oy Duodecim, Jyväskylä 2006;15-6.
332. Visakorpi J. Mistä tunnistan keliakian? Lasten ja nuorten erityispiirteitä. In: Mäki M et al, eds. *Keliakia*. Kustannus Oy Duodecim, Jyväskylä 2006;42-7.
333. Collin P, Huhtala H, Virta L, Kekkonen L, Reunala T. Diagnosis of coeliac disease in clinical practice. Physician's alertness to the condition essential. *J Clin Gastroenterol* 2007;41:152-6.
334. Collin P, Kekkonen L, Virta L, Reunala T, Kaukinen K. The diagnostic criteria of coeliac disease. Letter to the editor. *Acta Paediatr* 2007;96:604-6.
335. Collin P, Mäki M, Kaukinen K. Safe gluten threshold for patients with coeliac disease: some patients are more tolerant than others. Letter to the editor. *Am J Clin Nutr* 2007;86:260.
336. Collin P, Mäki M. Ohutsuoli. Keliakia. In: Höckerstedt K, Färkkilä M, Kivilaakso E, Pikkarainen P, eds. *Gastroenterologia ja hepatologia*. Duodecim, Jyväskylä 2007;378-88.
337. Collin P. Keliakian toteaminen on Suomessa huipputasoa. *Keliakia* 2007;(3):6-7.
338. Collin P. Risk groups of coeliac disease. In: Dolinsek J, Ornik T, eds. *Proceedings of the International Coeliac Disease Meeting, Maribor 13-16 September 2007, University Medical Center, Maribor* 2007:97.
339. Garcia-Horsman JA, Venäläinen JI, Lohi O, Auriola IS, Korponay-Szabo IR, Kaukinen K, Mäki M, Männistö PT. Deficient activity of prolyl oligopeptidase on the immunoactive peptide digestion in coeliac disease. *Scand J Gastroenterol* 2007;42:562-71.
340. Juuti-Uusitalo K, Kurppa K. XII Kansainvälinen keliakiakokous: Keliakiaan kehitteillä uusia hoitoja ja diagnoosimenetelmiä. *Keliakia* 2007;(1):6-7.
341. Juuti-Uusitalo K, Mäki M, Kainulainen H, Isola J, Kaukinen K. Gluten affects epithelial differentiation-associated genes in small intestinal mucosa of coeliac patients. *Clin Exp Immunol* 2007;150:294-305.
342. Kaukinen K, Collin P, Laurila K, Kaartinen T, Partanen J, Mäki M. Resurrection of gliadin antibodies in coeliac disease. Deamidated gliadin peptide antibody test provides additional diagnostic benefit. *Scand J Gastroenterol* 2007;42:1428-33.
343. Kaukinen K, Collin P, Mäki M. Latent coeliac disease or coeliac disease beyond villous atrophy? *Gut* 2007;56:1339-40.
344. Kaukinen K, Collin P, Mäki M. Long-term consumption of oats in coeliac disease. In: Stern M, ed. *Proceedings of the 21st Meeting, Working Group on Prolamin Analysis and Toxicity*. Verl. Wissensch. Scripten, Zwickau 2007;131-5.
345. Kaukinen K, Peräaho M, Lindfors K, Partanen J, Woolley N, Pikkarainen P, Karvonen A-L, Laasanen T, Sievänen H, Mäki M, Collin P. Persistent small bowel mucosal villous atrophy without symptoms in coeliac disease. *Aliment Pharmacol Ther* 2007;25:1237-45.
346. Kaukinen K. Ohutsuoli. Ohutsuolen immunologia. In: Höckerstedt K, Färkkilä M, Kivilaakso E, Pikkarainen P, eds. *Gastroenterologia ja hepatologia*. Duodecim, Jyväskylä 2007;375-7.
347. Kaukinen K. Ohutsuoli. Rakenne ja toiminta. In: Höckerstedt K, Färkkilä M, Kivilaakso E, Pikkarainen P, eds. *Gastroenterologia ja hepatologia*. Duodecim, Jyväskylä 2007;356-60.
348. Kaukinen K. Uusi kokoveripikatesti keliakian diagnostiikassa ja hoidon seurannassa. *Moodi* 2007;(1):64-5.

349. Kempainen T, Janatuinen E, Holm K, Kosma V-M, Heikkinen M, Mäki M, Laurila K, Uusitupa M, Julkunen R. No observed local immunological response at cell level after five years of oats in adult coeliac disease. *Scand J Gastroenterol* 2007;42:54-9.
350. Korponay-Szabó IR, Szabados K, Pustai J, Uhrin K, Ludmány É, Nemes É, Kaukinen K, Kapitány A, Koskinen L, Sipka S, Imre A, Mäki M. Population screening for coeliac disease in primary care by district nurses using a rapid antibody test: diagnostic accuracy and feasibility study. *Brit Med J* 2007;335:1244-7.
351. Korponay-Szabó IR. Antibody testing in coeliac disease. In: Dolinsek J, Ornik T, eds. Proceedings of the International Coeliac Disease Meeting, Maribor 13-16 September 2007, University Medical Center, Maribor 2007:71.
352. Kurppa K, Lähdeaho M-L. Uusia tuulia keliakiatutkimuksessa. Kansainvälinen keliakiakokous Mariborissa. *Keliakia* 2007;(5)10-11.
353. Lohi S, Mustalahti K, Kaukinen K, Laurila K, Collin P, Rissanen H, Lohi O, Bravi E, Gasparin M, Reunanen A, Mäki M. Increasing prevalence of coeliac disease over time. *Aliment Pharmacol Ther* 2007;26:1217-25.
354. Mäki M. Lääkehoito keliakiaan? *Keliakia* 2007;(4):6-7.
355. Raivio T, Korponay-Szabó I, Collin P, Laurila K, Huhtala H, Kaartinen T, Partanen J, Mäki M, Kaukinen K. Performance of a new rapid whole blood coeliac test in adult patients with low prevalence of endomysial antibodies. *Dig Liver Dis* 2007;39:1057-63.
356. Saavalainen P, Kaukinen K, de Kauwe A. Tutkimusta verisuolilla: keliakiarokote ja uutta diagnostiikkaa? *Keliakia* 2007;(4):8-10.
357. Saavalainen P. Genetics of celiac disease. In: Dolinsek J, Ornik T, eds. Proceedings of the International Coeliac Disease Meeting, Maribor 13-16 September 2007, University Medical Center, Maribor 2007:61.
358. Salmi T. Keliakian diagnostiset kriteerit vaativat uudistamista. *Keliakia* 2007;(5)8-9.
359. Viljamaa M. Hoitamaton keliakia - osteoporoosin riskitekijä. *Keliakia* 2007;(2):6-7.
360. Viljamaa M. Keliakia ja elämänlaatu. *Keliakia* 2007;(3):8-9.
361. Ashorn S, Raukola H, Välineva T, Ashorn M, Wei B, Braun J, Rantala I, Kaukinen K, Luukkaala T, Collin P, Mäki M, Iltanen S. Elevated serum anti-*Saccharomyces cerevisiae*, anti-I2 and anti-OmpW antibody levels in patients with suspicion of celiac disease. *J Clin Immunol* 2008;28:486-94.
362. Collin P, Kaukinen K, Mattila AK, Joukamaa M. Psychoneurotic symptoms and alexithymia in coeliac disease. *Scand J Gastroenterol* 2008;43:1329-33.
363. Collin P, Kaukinen K. Serologic screening for coeliac disease in risk groups: is once in the lifetime enough? *Dig Liver Dis* 2008;40:101-3.
364. Collin P. Rasvaripuli vai yleistauti? In: Anttila S et al., eds. XXIX Tampereen lääkäripäivät Tampereella 13.-15.3.2008, luennot. Lege Artis Oy, Vammala 2008;18-9.
365. Collin P. Vain noin puolet keliakiasta tunnetaan. *Yleislääkärelehti* 2008;23:9-11.
366. Haveri H, Westerholm-Ormio M, Lindfors K, Mäki M, Andersson LC, Heikinheimo M. Transcription factors GATA-4 and GATA-6 in normal and neoplastic human gastrointestinal mucosa. *BMC Gastroenterology* 2008;8:9.
367. Kaukinen K, Collin P, Mäki M. Natural history of celiac disease. In: Fasano A, Troncone R, Branski D, eds. *Frontiers in Celiac Disease*. *Pediatr Adolesc Med*. Basel, Karger 2008;12:12-7.
368. Kaukinen K, Collin P, Mäki M. Vehnätärkkelyspohjaiset makeutusaineet ovat turvallisia keliakian hoidossa. *Keliakia* 2008;(4):6-7.
369. Kaukinen K, Mäki M. Self-detection of coeliac disease through a transglutaminase-based antibody: from laboratory to onsite testing. In: Stern M, ed. Proceedings of the 22nd Meeting, Working Group on Prolamin Analysis and Toxicity. Verlag Wissenschaftliche Scripten, Zwickau 2008:81-4.
370. Kaukinen K, Salmi T, Collin P, Huhtala H, Kärjä-Lahdensuu T, Mäki M. Clinical trial: gluten microchallenge with wheat-based starch hydrolysates in coeliac disease patients - a randomized, double-blind, placebo-controlled study to evaluate safety. *Aliment Pharmacol Ther* 2008;28:1240-8.
371. Kaukinen K. Keliakian hoidon ongelmat. In: Anttila S et al., eds. XXIX Tampereen lääkäripäivät Tampereella 13.-15.3.2008, luennot. Lege Artis Oy, Vammala 2008;23.
372. Kondrashova A, Mustalahti K, Kaukinen K, Viskari H, Volodicheva V, Haapala AM, Ilonen J, Knip M, Mäki M, Hyöty H, and the EPIVIR Study Group. Lower economic status and inferior hygienic environment may protect against celiac disease. *Ann Med* 2008;40:223-31.
373. Korkeamäki H, Viiri K, Kukkonen MK, Mäki M, Lohi O. Alternative mRNA splicing of SAP30L regulates its transcriptional repression activity. *FEBS Lett* 2008;582:379-84.
374. Korponay-Szabó IR, Vecsei Z, Kiraly R, Dahlbom I, Chirido R, Nemes E, Fesus L, Mäki M. Deamidated gliadin peptides form epitopes that transglutaminase antibodies recognize. *J Pediatr Gastroenterol Nutr* 2008;46:253-61.
375. Koskinen LLE, Korponay-Szabó IR, Viiri K, Juuti-Uusitalo K, Kaukinen K, Lindors K, Mustalahti K, Kurppa K, Adany R, Pocsai Z, Szeles G, Einarsdottir E, Wijmenga C, Mäki M, Partanen J, Kere J, Saavalainen P. Myosin IXB gene region and gluten intolerance: linkage to coeliac disease and a putative dermatitis herpetiformis association. *J Med Genet* 2008;45:222-7.
376. Koskinen O, Collin P, Korponay-Szabó I, Salmi T, Iltanen S, Haimila K, Partanen J, Mäki M, Kaukinen K. Gluten-dependent small bowel mucosal transglutaminase 2-specific IgA deposits in overt and mild enteropathy coeliac disease. *J Pediatr Gastroenterol Nutr* 2008;47:436-42.
377. Kurppa K, Koskinen O, Collin P, Mäki M, Reunala T, Kaukinen K. Changing phenotype of celiac disease after long-term gluten exposure. *J Pediatr Gastroenterol Nutr* 2008;47:500-3.

378. Kurppa K, Saavalainen P, Mäki M, Kaukinen K. Perinnöllisyystutkimuksessa on löydetty runsaasti keliakiaepäilyjä. *Keliakia* 2008;(4):8-9.
379. Kurppa K, Sütas Y, Lähdeaho M-L, Ashorn M. Pienen pojan vatsavaivat, kasvuhäiriö ja hypoalbuminemia. *Duodecim* 2008;124:103.
380. Lindfors K, Blomqvist T, Juuti-Uusitalo K, Stenman S, Venäläinen J, Mäki M, Kaukinen K. Live probiotic *Bifidobacterium lactis* bacteria inhibit the toxic effects induced by wheat gliadin in epithelial cell culture. *Clin Exp Immunol* 2008;152:552-8.
381. Lohi S. Keliakian esiintyvyys on kaksinkertaistunut 1980-luvulta. *Keliakia* 2/2008:6.
382. Lohi S, Mäki M, Montonen J, Knekt P, Pukkala E, Reunanen A, Kaukinen K. Tunnistamaton keliakia ei lisää yleistä syöpäriskiä. *Duodecim* 2008;124:2484.
383. Myrsky E, Kaukinen K, Syrjänen M, Korponay-Szabo IR, Mäki M, Lindfors K. Celiac disease-specific autoantibodies targeted against transglutaminase 2 disturb angiogenesis. *Clin Exp Immunol* 2008;152:111-9.
384. Mäki M. Celiac disease. In: Kleinman RE, Goulet O, Mieli-Vergani G, Sanderson IR, Sherman P, Shneider BL, eds. *Walker's Pediatric Gastrointestinal Disease*, 5th Edition. BC Decker Inc, Hamilton 2008;319-27.
385. Nemes E, Lefler E, Szegedi L, Kapitány A, Kovács JB, Balogh M, Szabados K, Tumppek J, Sipka S, Korponay-Szabó IR. Gluten intake interferes with the humoral immune response to recombinant hepatitis B vaccine in patients with celiac disease. *Pediatrics* 2008;121:e1570-6.
386. Oikarinen M, Tauriainen S, Honkanen T, Oikarinen S, Vuori K, Kaukinen K, Rantala I, Mäki M, Hyöty H. Detection of enteroviruses in the intestine of type 1 diabetic patients. *Clin Exp Immunol* 2008;151:71-5.
387. Raivio T. Keliakian toteaminen helpottuu. *Keliakia* 2008;(3):6-7.
388. Raivio T, Korponay-Szabo IR, Paajanen T, Ashorn M, Iltanen S, Collin P, Laurila K, Nemes E, Kovacs JB, Carrard G, Saramäki M, Mäki M, Kaukinen K. Comparison of a novel whole blood transglutaminase-based ELISA with a whole blood rapid antibody test and established conventional serological coeliac disease assays. *J Pediatr Gastroenterol Nutr* 2008;47:562-7.
389. Salmi T. Keliakian seulonnalla diagnoosiin, diagnostiset kriteerit. In: Anttila S et al, eds. XXIX Tampereen Lääkärpäivät Tampereella 13.-15.3.2008, luennot. Lege Artis Oy, Vammala 2008;21-2.
390. Salo M, Collin P, Kyrönpalo S, Rasmussen M, Huhtala H, Kaukinen K. Age, symptoms and upper gastrointestinal malignancy in primary care endoscopy. *Scand J Gastroenterol* 2008;43:122-7.
391. Stenman S, Lindfors K, Korponay-Szabo IR, Lohi O, Saavalainen P, Partanen J, Haimila K, Wieser H, Mäki M, Kaukinen K: Secretion of celiac disease autoantibodies after in vitro gliadin challenge is dependent on small-bowel mucosal transglutaminase 2-specific IgA deposits. *BMC Immunology* 2008;9:6.
392. Troncone R, Ivarsson A, Szajewska H, Mearin ML, the European Multistakeholder Platform on CD (CDEUSSA), incl. Mäki M, Kaukinen K, Lähdeaho ML. Review article: future research on coeliac disease - a position report from the European multistakeholder platform on coeliac disease (CDEUSSA). *Aliment Pharmacol Therap* 2008;27:1030-43.
393. Venäläinen J, Männistö PT, Kaukovirta-Norja A, Loponen J, Salovaara H, Stenman S, Lindfors K, Kaukinen K, Mäki M. Viljojen omista entsyymeistä apua keliakiaan? *Dosis* 2008;24:67-75.
394. Vilppula A, Collin P, Mäki M, Valve R, Luostarinen M, Krekelä I, Patrikainen H, Kaukinen K, Luostarinen L. Undetected coeliac disease in the elderly. A biopsy-proven population-based study. *Digest Liver Dis* 2008;40:809-13.
395. Viiri LE, Viiri KM, Ilveskoski E, Huhtala H, Mäki M, Tienari PJ, Perola M, Lehtimäki T, Karhunen PJ. Interactions of functional apolipoprotein E gene promoter polymorphisms with smoking on aortic atherosclerosis. *Circ Cardiovasc Genet* 2008;1:107-16. (doi:10.1161/Circgenetics.108.791764).
396. Ashorn S, Välineva T, Kaukinen K, Ashorn M, Braun J, Raukola H, Rantala I, Collin P, Mäki M, Luukkaala T, Iltanen S. Serological responses to microbial antigens in celiac disease patients during a gluten-free diet. *J Clin Immunol* 2009;29:190-5.
397. Collin P, Kaukinen K, Mäki M. Duodenal intraepithelial lymphocytosis: Celiac disease or not? Letter to the Editor. *Am J Gastroenterol* 2009;104:1847.
398. Einarsdottir E, Koskinen LLE, Dukes E, Kainu K, Suomela S, Lappalainen M, Zibera F, Korponay-Szabo IR, Kurppa K, Kaukinen K, Adany R, Pocsai Z, Széles G, Färkkilä M, Turunen U, Halme L, Paavola-Sakki P, Not T, Vatta S, Ventura A, Löfberg R, Torkvist P, Bresso F, Halfvarson J, Mäki M, Kontula K, Saarialho-Kere U, Kere J, D'Amato M, Saavalainen P. IL23R in the Swedish, Finnish, Hungarian and Italian populations: association with IBD and psoriasis, and linkage to coeliac disease. *BMC Medical Genetics* 2009;10:8. (doi:10.1186/1471-2350-10-8.)
399. Haimila K, Einarsdottir E, de Kauwe A, Koskinen LLE, Pan-Hammarström Q, Kaartinen T, Kurppa K, Zibera F, Not T, Vatta S, Ventura A, Korponay-Szabo IR, Ádány R, Pocsai Z, Széles G, Dukes E, Kaukinen K, Mäki M, Koskinen S, Partanen J, Hammarström L, Saavalainen P. The shared CTLA4-ICOS risk locus in celiac disease, IgA deficiency and common variable immunodeficiency. *Genes Immun* 2009;10:151-61 (doi:10.1038/gene.2008.89).
400. Heinonen TY, Mäki M. Peters'-plus syndrome is a congenital disorder of glycosylation caused by a defect in the  $\beta$ 1,3-glucosyltransferase that modifies thrombospondin type 1 repeats. *Ann Med* 2009;41:2-10. (DOI: 10.1080/07853890802301975).

401. Juuti-Uusitalo K, Lindfors K, Mäki M, Patrikainen M, Isola J, Kaukinen K. Inhibition of epithelial growth factor receptor signalling does not preserve epithelial barrier function after *in vitro* gliadin insult. *Scand J Gastroenterol* 2009;44:820-5.
402. Kaukinen K, Collin P, Mäki M. Keliakian nykyhoito: ”vältä viljoja” ei riitä ohjeeksi. *S Lääkäril* 2009;64:277-83.
403. Koskinen L, Romanos J, Kaukinen K, Mustalahti K, Korponay-Szabo I, Barisani D, Bardella MT, Zibera F, Vatta S, Széles G, Pocsai Z, Karell K, Haimila K, Ádány R, Not T, Ventura A, Mäki M, Partanen J, Wijmenga C, Saavalainen P. Cost-effective HLA-typing with tagging SNPs predicts celiac disease risk haplotypes in the Finnish, Hungarian and Italian populations. *Immunogenetics* 2009;61:247-56 (DOI 10.1007/s00251-009-0361-3).
404. Koskinen LLE, Einarsdóttir E, Dukes E, Heap GAR, Dubois P, Korponay-Szabo IR, Kaukinen K, Kurppa K, Zibera F, Vatta S, Not T, Ventura A, Sistonen P, Ádány R, Pocsai Z, Széles G, Mäki M, Kere J, Wijmenga C, van Heel DA, Saavalainen P. Association study of the IL18RAP locus in three European populations with coeliac disease. *Human Molec Genet* 2009;18:1148-55.
405. Koskinen O, Villanen M, Korponay-Szabo I, Lindfors K, Mäki M, Kaukinen K. Oats do not induce systemic or mucosal autoantibody response in children with coeliac disease. *J Pediatr Gastroenterol Nutr* 2009;48:559-65.
406. Kurppa K, Collin P, Viljamaa M, Haimila K, Saavalainen P, Partanen J, Laurila K, Huhtala H, Paasikivi K, Mäki M, Kaukinen K. Diagnosing mild enteropathy celiac disease: a randomized, controlled clinical study. *Gastroenterology* 2009;136:816-23. (doi:10.1053/j.gastro.2008.11.040).
407. Kurppa K, Collin P, Viljamaa M, Haimila K, Saavalainen P, Partanen J, Laurila K, Huhtala H, Paasikivi K, Mäki M, Kaukinen K. Keliakia tulee hoitaa riippumatta ohutsuulilöydöksestä. *Duodecim* 2009;125:307.
408. Lindfors K, Kaukinen K, Mäki M. A role for anti-transglutaminase 2 autoantibodies in the pathogenesis of coeliac disease? *Amino Acids* 2009;36:685-91 (DOI 10.1007/s00726-008-0127-5).
409. Lohi S, Mäki M, Montonen J, Knekt P, Pukkala E, Reunanen A, Kaukinen K. Malignancies in cases with screening-identified evidence of coeliac disease: a long-term population-based cohort study. *Gut* 2009;58:643-7 (doi:10.1136/gut.2007.140970).
410. Myrsky E, Syrjänen M, Korponay-Szabo IR, Mäki M, Kaukinen K, Lindfors K. Altered small-bowel mucosal vasculature network in untreated coeliac disease. *Scand J Gastroenterol* 2009;44:162-7.
411. Peltola M, Kaukinen K, Dastidar P, Haimila K, Partanen J, Haapala AM, Mäki M, Keränen T, Peltola J. Hippocampal sclerosis in refractory temporal lobe epilepsy is associated with gluten sensitivity. *J Neurol Neurosurg Psychiatry* 2009;80:626-30 (doi:10.1136/jnnp.2008.148221).
412. Rauhavirta T. 13. kansainvälinen keliakiakokous 6.-8.4.2009 Amsterdamissa. Tulppaneja ja tutkimustuloksia. *Keliakia* 2009;(3):6-8.
413. Salovaara H, Kanerva P, Kaukinen K, Sontag-Strohm T. Oats - an overview from a celiac disease point of view. In: Arendt EK, Dal Bello F. *The Science of Gluten-Free Foods and Beverages. Proceedings of the First International Conference on Gluten-Free Cereal Products and Beverages.* AACC International, Inc. St. Paul, Minnesota 2009;69-87.
414. Sareneva I, Koskinen LLE, Korponay-Szabo IR, Kaukinen K, Kurppa K, Zibera F, Vatta S, Not T, Ventura A, Ádány R, Pocsai Z, Széles G, Mäki M, Saavalainen P, Einarsdóttir E. Linkage and association study of FCγR polymorphisms in celiac disease. *Tissue Antigens* 2009;73:54-8.
415. Solakivi T, Kaukinen K, Kunnas T, Lehtimäki T, Mäki M, Nikkari ST. Serum fatty acid profile in celiac disease patients before and after a gluten-free diet. *Scand J Gastroenterol* 2009;44:826-30.
416. Stenman SM, Venäläinen JI, Lindfors K, Auriola S, Mauriala T, Kaukovirta-Norja A, Jantunen A, Laurila K, Quiao S-W, Sollid LM, Männistö PT, Kaukinen K, Mäki M. Enzymatic detoxification of gluten by germinating wheat proteases: Implications for new treatment of celiac disease. *Ann Med* 2009;41:390-400.
417. Viiri KM, Jänis J, Siggers T, Heinonen TYK, Valjakka J, Bulyk ML, Mäki M, Lohi O. DNA-binding and – bending activities of SAP30L and SAP30 are mediated by a zinc-dependent module and monophosphoinositides. *Molec Cell Biol* 2009;29:342-56.
418. Viiri KM, Heinonen TYK, Mäki M, Lohi O. Phylogenetic analysis of the SAP30 family of transcriptional regulators reveals functional divergence in the domain that binds the nuclear matrix. *BMC Evol Biol* 2009;9:149. (doi:10.1186/1471-2148-9-149).
419. Vilppula A, Kaukinen K, Luostarinen L, Krekelä I, Patrikainen H, Valve R, Mäki M, Collin P. Increasing prevalence and high incidence of celiac disease in elderly people: A population-based study. *BMC Gastroenterology* 2009;9:49 (doi:10.1186/1471-230X-9-49).
420. Vilppula A. Keliakia ikääntyvässä väestössä. *Keliakia* 2009;(1):6-7.
421. Koskinen LLE, Einarsdóttir E, Korponay-Szabo IR, Kurppa K, Kaukinen K, Sistonen P, Pocsai Z, Széles G, Ádány R, Mäki M, Kere J, Saavalainen P. Finemapping of the *CELIAC2* locus on chromosome 5q31-q33 in the Finnish and Hungarian populations. *Tissue Antigens*, in press.
422. Koskinen O, Collin P, Lindfors K, Laurila K, Mäki M, Kaukinen K. Usefulness of small-bowel mucosal transglutaminase-2 specific autoantibody deposits in the diagnosis and follow-up of celiac disease. *J Clin Gastroenterol*, in press.
423. Lohi S, Mäki M, Rissanen H, Knekt P, Reunanen A, Kaukinen K. Prognosis of unrecognized coeliac disease as regards mortality: A population-based cohort study. *Ann Med*, in press (2009,1-8,iFirst).
424. Lohi S, Lohi O, Vierula M, Mäki M, Toppari J. Coeliac disease autoantibodies in seminal plasma from cases with screen-detected coeliac disease. *Scand J Gastroenterol*, in press.

425. Myrsky E, Caja S, Simon-Vecsei Z, Korponay-Szabo IR, Nadalutti C, Collighan R, Mongeot A, Griffin M, Mäki M, Kaukinen K, Lindfors K. Coeliac disease IgA modulates vascular permeability in vitro through the activity of transglutaminase 2 and RhoA. *Cell Mol Life Sci*, in press (DOI 10.1007/s00018-009-0116-1).
426. Salmi TT, Collin P, Reunala T, Mäki M, Kaukinen K. Diagnostic methods beyond conventional histology in coeliac disease diagnosis. *Digest Liver Disease*, in press (doi:10.1016/j.dld.2009.04.004).
427. Virta LJ, Kaukinen K, Collin P. The incidence and prevalence of diagnosed coeliac disease in Finland. Results of effective case finding in adults. *Scand J Gastroenterol*, in press.

## Patents

1. Mäki M, Korponay-Szabo I. USA States Patent Number 7,361,480 - USA, Methods and Means for Detecting Gluten-Induced Diseases, Patent Granted 22.4.2008.
2. Mäki M, Korponay-Szabo I. European Patent No. 1390753. Methods and Means for Detecting Gluten-Induced Diseases, European Patent Office 22.10.2008.
3. Viiri K, Mäki M, Lohi O. Patent Application no 20086014-Finland. 24.10.2008.

## Films

1. Mäki M. Small bowel biopsy in children. Rapid technique and handling of the biopsy specimen. Video 17 min., U-matic, Alfa-Video 1984, Tampere, Finland.

## Abstracts

1. Huupponen T, Mäki M, Hällström O, Reunala T, Vesikari T, Visakorpi JK. Reticulin antibodies as an indicator of coeliac disease in diabetic children. Scandinavian Society for the study of diabetes. 18th Annual Meeting, June 1-3, 1983, Kuopio, Finland. Abstract 30.
2. Hietanen E, Heinonen T, Ståhlberg M-R, Mäki M. Metabolism of xenobiotics in the small intestinal mucosa of children. Extrahepatic drug metabolism and chemical carcinogenesis. Univ. Stockholm and the Royal Swedish Academy of Sciences, IUB Symposium nr 125, Stockholm, Sweden, May 17-20, 1983. Abstract A 38.
3. Mäki M. Evaluation of a serum IgA class reticulin antibody test for the detection of childhood coeliac disease. 6. Symposium der Arbeitsgemeinschaft Gastroenterologie, Ernährung und Stoffwechsel der Gesellschaft für Pädiatrie der DDR. Oberhof 21.-23.5.1984, DDR. Abstract.
4. Mäki M, Hällström O, Vesikari T, Visakorpi JK. Serum IgA reticulin antibody detects childhood coeliac disease. *Pediatr Res* 1984; 18:1054.
5. Mäki M, Alfa-Video Tampere. Small bowel biopsy in children. Rapid technique and handling of the biopsy specimen. Teaching videotape, 17 minutes. *Acta Paediatr Scand Suppl* 1985; Suppl 322:33.
6. Aine L, Mäki M, Aine E, Visakorpi JK. Enamel hypoplasia (EH) of permanent teeth in coeliac disease (CD). *Pediatr Res* 1986; 20:696.
7. Ståhlberg M-R, Hietanen E, Mäki M. Detoxication enzymes in small intestinal mucosa in children. *Pediatr Res* 1986; 20:697.
8. Mäki M, Hällström O, Verronen P, Visakorpi JK. Arthritis in childhood coeliac disease (CD). *Pediatr Res* 1987; 22:101.
9. Mäki M, Ståhlberg M-R, Hällström O, Viander M, Visakorpi JK. Predictability of small bowel mucosal damage by reticulin (ARA) and gliadin antibodies (AGA). *Pediatr Res* 1987; 22:103.
10. Mäki M, Hällström O, Visakorpi JK. Gluten challenge in postpubertal children with coeliac disease (CD): Detection of mucosal relapse with IgA-class reticulin antibody (ARA). *Pediatr Res* 1987; 22:103.
11. Mäki M, Hällström O, Polanco I, Visakorpi JK. Comparison of reticulin (ARA) and endomysium (EMA) antibodies in coeliac disease. *Pediatr Res* 1988; 24:406.
12. Mäki M, Visakorpi JK. Normal small bowel histology does not exclude coeliac disease (CD). *Pediatr Res* 1988; 24:411.
13. Koskimies S, Mäki M, Lipsanen V, Visakorpi JK. HLA haplotypes in Finnish coeliac patients and their family members. Possible role of protecting haplotypes. Samuel Gee 1888 - 1988 Coeliac Symposium, St. Bartholomew's Hospital, 4th - 6th September, 1988. Abstract T10.
14. Mäki M, Koskimies S, Visakorpi JK. Latent coeliac disease. Samuel Gee 1888 - 1988 Coeliac Symposium, St. Bartholomew's Hospital, 4th - 6th September, 1988. Abstract T31.
15. Mäki M. Milloin on syytä epäillä keliakiaa? Tampereen Lääkärpäivät 1988, Luennot IX, s. 74.
16. Mäki M. Occurrence of coeliac disease in Finland. Tampereen Lääkärpäivät -89. Luennot X, p. 143.
17. Koskimies S, Lipsanen V, Mäki M, Visakorpi JK. Restriction fragment length polymorphism of HLA class II genes in coeliac patients, their family members and controls. *Pediatr Res* 1989;26:274.
18. Lähdeaho M-L, Lehtinen M, Mäki M, Parkkonen P, Rissa H-R, Hällström O, Visakorpi JK. Adenovirus antibodies in childhood coeliac disease, antigliadin antibodies in adenovirus diseased children - what is the chain of events? *Pediatr Res* 1989;26:274.
19. Mäki M, Koskimies S, Visakorpi JK. Latent coeliac disease. In: Kumar PJ, Walker-Smith JA, eds. *Coeliac Disease: One Hundred Years*. Univ. Printing Service, Univ. Leeds, Leeds, 1991:245.
20. Holm K, Mäki M, Hällström O, Koskimies S, Visakorpi JK. Healthy individuals with or without enteropathy but positive for serum reticulin antibodies (RA) are genetically similar to coeliac disease (CD) patients. *J Pediatr Gastroenterol Nutr* 1991;13:321.
21. Savilahti E, Reunala T, Mäki M. Increase of gamma/delta receptor bearing lymphocytes in the jejunum of patients with dermatitis herpetiformis. *J Pediatr Gastroenterol Nutr* 1991;13:323.
22. Mäki M, Hällström O, Marttinen A. Purification of human protein molecules - autoantigens - reacting with coeliac disease (CD) patient sera IgA. *J Pediatr Gastroenterol Nutr* 1991;13:321.
23. Lähdeaho M-L, Mäki M, Parkkonen P, Lehtinen M. Antibodies to a structural and a non-structural adenovirus antigen in adenovirus diseased children and children with coeliac disease. 12th Nordic Virus Symposium, June 13-15, 1991, Espoo, Finland. Abstract.
24. Ascher H, Holm K, Kristiansson B, Mäki M. Different clinical and epidemiological features of childhood coeliac disease in two neighbouring countries, Sweden and Finland. *Epidemiology of Coeliac Disease*, Capri 11-12 October 1991. Abstract p. 34.
25. Mäki M, Savilahti E, Holm K, Lipsanen V, Koskimies S. Association between genetic markers and the density of intraepithelial (IEL) gamma/delta and alpha/beta T cells in healthy first-degree relatives of coeliac disease (CD) patients. Eleventh Annual Bristol Myers Squibb/Mead Johnson Nutrition Research Symposium "Immunophysiology of the Gut", Boston, August 26-28, 1991, Abstract No. 5.
26. Ascher H, Kristiansson B, Holm K, Mäki M. Vad betyder glutenkonsumtionsmönstret för skillnaderna mellan celiaki i Sverige och Finland? XLVIII Svenska Läkaresällskapets Riksstämma, Stockholm 27-29 november 1991. Svenska Läkaresällskapets Handlingar Hygiea, 1991;Band 199, Häfte 2:297.

27. Savilahti E, Mäki M, Holm K, Reunala T, Klemola T, Koskimies S. Intraepithelial gamma/delta T-cells receptor bearing cells as a marker of latent coeliac disease. Workshop on Epidemiology of Coeliac Disease. Capri, 11-12 October 1991. Abstract p. 28.
28. Mäki M. Screening tools to be used. Workshop on Epidemiology of Coeliac Disease. Capri, 11-12 October 1991. Abstract p. 26.
29. Mäki M. Effect of the type and amount of cereals in the diet. Workshop on Epidemiology of Coeliac Disease. Capri, 11-12 October 1991. Abstract p. 23.
30. Collin P, Korpela M, Keyriläinen O, Hällström O, Viander M, Mäki M. Rheumatic complaints as a presenting symptom in patients with coeliac disease. *Scand J Rheumatol* 1991;20:222.
31. Ascher H, Holm K, Kristiansson B, Mäki M. Quality and quantity of gluten intake in infancy in Sweden and Finland; one factor of importance for the clinical expression of coeliac disease? *J Pediatr Gastroenterol Nutr* 1992;15:347.
32. Holm K, Mäki M, Savilahti E, Lipsanen V, Koskimies S, Visakorpi JK. Association between DR, DQA and DQB genes and the density of jejunal intraepithelial gamma/delta T cells. *J Pediatr Gastroenterol Nutr* 1992;15:352.
33. Holm K, Savilahti E, Mäki M, Koskimies S, Lipsanen V. HLA class II antigens in the intestine of first degree relatives of coeliac patients. *J Pediatr Gastroenterol Nutr* 1992;15:352.
34. Holm K, Savilahti E, Koskimies S, Lipsanen V, Mäki M. Coeliac disease family study: immunohistochemical changes in the jejunum of first-degree relatives. International Coeliac Symposium, Dublin 22-25 July, 1992. Abstract 36.
35. Marttinen A, Mäki M. Human fibroblasts synthesise and secrete coeliac disease autoantigen molecules. International Coeliac Symposium, Dublin 22-25 July, 1992. Abstract 47.
36. Reunala T, Mäki M. Gluten-free diet treatment in dermatitis herpetiformis: a 5-year study. International Coeliac Symposium, Dublin 22-25 July, 1992. Abstract 54.
37. Mäki M, Huupponen T, Holm K, Hällström O. Onset of diabetes initiates coeliac disease autoantibody production and development of enteropathy. International Coeliac Symposium, Dublin 22-25 July, 1992. Abstract 44.
38. Mäki M. The role of antibody in coeliac disease pathogenesis. International Coeliac Symposium, Dublin 22-25 July, 1992. Abstract p. 24.
39. Reunala T, Aine L, Hällström O, Laurila K, Mäki M. Celiac-type dental enamel defects and circulating IgA-class antibodies in children with dermatitis herpetiformis. *Ped Dermatol* 1992;9:180.
40. Reunala T, Aine L, Mäki M. Coeliac-type dental enamel defects in patients with dermatitis herpetiformis. 18th World Congress of Dermatology, New York City, June 12-18, 1992. Abstract 189 A.
41. Ascher H, Holm K, Kristiansson B, Mäki M. Different trends in epidemiology and clinical expression of childhood coeliac disease in Sweden and Finland. Is gluten intake in infancy a factor of importance? *Scand J Gastroenterol* 1993;28(suppl. 197):53.
42. Collin P, Helin H, Mäki M, Hällström O, Karvonen AL. Latent coeliac disease in patients positive in reticulin and gliadin antibody tests. *Scand J Gastroenterol* 1993;28(suppl. 197):53.
43. Marttinen A, Hakala S, Mäki M. Reticulin and endomysium antibodies recognize fibroblast-derived molecules. *Scand J Gastroenterol* 1993;28(suppl. 197):54.
44. Holm K, Rantala I, Hällström O, Ivanyi J, Mäki M. Gamma/delta T cell density and heat-shock protein expression in jejunal epithelium in children suspected for coeliac disease. *Scand J Gastroenterol* 1993;28(suppl. 197):54.
45. Hakala S, Mäki M, Marttinen A. Coeliac disease (CD) patient sera IgA recognizes human fibroblast-derived molecules. *J Pediatr Gastroenterol Nutr* 1993;17:462.
46. Lähdeaho ML, Vainio E, Mäki M, Lehtinen M. Identification of a coeliac disease specific B-cell epitope in A-gliadin. *J Pediatr Gastroenterol Nutr* 1993;17:462.
47. Holm K, Rantala I, Hällström O, Ivanyi J, Mäki M. Enhanced expression of the human groEL stress-protein homologue (HSP65) in jejunal enterocytes in children with reticulin autoantibodies with or without mucosal atrophy. *J Pediatr Gastroenterol Nutr* 1993;17:471.
48. Mäki M, Marttinen A. Fibroblast-derived human proteins are targets for genetically determined reticulin and endomysium autoantibodies. *Ped Res* 1994;35:266.
49. Erilä T, Collin P, Pirttilä T, Nurmikko T, Somer H. Neurologic complications of coeliac disease. XIVth World Congress of Neurology. October 23, 1989, New Delhi, India. Abstract
50. Mäki M, Marttinen A. Human fibroblast in vitro produced protein complex contains targets for reticulin and endomysium autoantibodies. 7th Annual Meeting of the North American Society for Pediatric Gastroenterology and Nutrition. November 5-6, 1993, Chicago, USA. Abstract 27.
51. Mäki M, Huupponen T, Holm K, Hällström O. Serum reticulin autoantibody production and development of coeliac disease is triggered by onset of IDDM. In: Greene S (ed.) *Diabetes in the young*. XIXth Annual Meeting Aegean Sea, Greece, 1993, p. 68. Medicom Europe BV, The Netherlands.
52. Mäki M. Keliakian epidemiologia ja patogeneesi. XV Tampereen Lääkäripäivät 17.-19.3.1994, Luennot XV, s. 31.
53. Mäki M. Antibodies in coeliac disease. Mucosal Immunity and the gut epithelium. Interactions in health and disease. Capri 22-23 April 1994, p. 25.
54. Holm K, Mäki M, Eland C, Partanen J, Kainulainen H, Koskimies S, Ruuska T, Rantala I. Jejunal mucosal heat-shock protein 65 expression in coeliac disease. Mucosal Immunity and the gut epithelium. Interactions in health and disease. Capri 22-23 April 1994, Abstract 43, p.77.

55. Halttunen T, Marttinen A, Kainulainen H, Mäki M. Three-dimensional growth and fibroblast-supported differentiation of a gut epithelial cell line (T84). *Experimental Biology 94*, Anaheim, California, April 24-28, 1994. *The FASEB Journal*, Abstracts Part II 1994;8:A1007.
56. Jalava T, Marttinen A, Mäki M, Koskimies S. Immune response in vitro to coeliac disease associated autoantigen, reticulin, is MHC restricted. 12th European Immunology Meeting, 14-17 June 1994, Barcelona, Spain. Abstract p. 341.
57. Rantala I, Holm K, Mustonen J, Collin P, Kainulainen H, Mäki M. Immunohistochemical quantitation of jejunal T cells, HLA-DR antigen, and human groEL stress-protein homologue in IgA nephropathy. *IgA Nephropathy International Symposium 28.2*. - 1.3.1994, Adelaide, Australia. Abstract.
58. Kerttula TO, Hällström O, Mäki M. Analysis of peripheral blood T cells in patients with untreated coeliac disease (CoD) - elevation of CD45RP+ T cells. 12th European Immunology Meeting, 14-17 June 1994, Barcelona, Spain. Abstract p. 343.
59. Sulkanen S, Marttinen A, Mäki M. Human fibroblast-derived molecules as antigen in ELISA for celiac disease-specific IgA. *J Pediatr Gastroenterol Nutr* 1994;19:332.
60. Marttinen A, Jalava T, Koskimies S, Mäki M. HLA restricted immune response in vitro human fibroblast-derived celiac disease-associated autoantigen. *J Pediatr Gastroenterol Nutr* 1994;19:363.
61. Halttunen T, Marttinen A, Kainulainen H, Mäki M. Fibroblast-induced differentiation and three-dimensional organization of a gut epithelial cell line (T84). *FEBS Special Meetings, Biological Membranes*, Helsinki, Finland, June 26 - July 1, 1994. Abstract 390.
62. Sulkanen S, Marttinen A, Hällström O, Laurila K, Halttunen T, Mäki M. Human umbilical cord autoantibodies (HUC-ab). An ideal screening test for coeliac disease. *J Pediatr Gastroenterol Nutr* 1995;20:445.
63. Hacsek G, Savilahti E, Arato A, Reunala T, Mäki M, Nemeth A. Effect of age on jejunal intraepithelial lymphocyte subsets in patients with gluten sensitive enteropathy and in healthy controls. *J Pediatr Gastroenterol Nutr* 1995;20:474.
64. Polvi A, Koskimies S, Mäki M, Partanen J. A gene dosage effect of DQA1\*0501 and DQB1\*0201 alleles in human coeliac disease. *Scand J Immunol* 1995;41:636.
65. Halttunen T, Marttinen A, Mäki M. Fibroblast-induced differentiation of the epithelial cell line T84 is due to TGF- $\beta$ . *Clin Immunol Immunopathol* 1995;76:S3.
66. Koskimies S, Jalava T, Marttinen A, Partanen J, Mäki M. Immune response in vitro to coeliac disease associated auto-antigen is DR and DQ restricted. *Clin Immunol Immunopathol* 1995;76:S72.
67. Mäki M, Sulkanen S, Laurila K, Hällström O, Halttunen T, Marttinen A. Human umbilical cord autoantibodies (IgA-HUC-AB) are sensitive and specific for celiac disease. *Clin Immunol Immunopathol* 1995;76:S73.
68. Marttinen A, Sulkanen S, Mäki M. ELISA for celiac disease-specific IgA with human fibroblast-synthesised molecules as the antigen. *Clin Immunol Immunopathol* 1995;76:S73.
69. Savilahti E, Hacsek G, Arato A, Reunala T, Mäki M, Nemeth A. Age and the density of intraepithelial gamma/delta T cell receptor positive cells in patients with coeliac disease. *Clin Immunol Immunopathol* 1995;76:S73.
70. Collin P, Nurmikko T, Himanen S-L, Eirilä T, Mäki M. Brain atrophy and coeliac disease. International meeting on epilepsy and other neurological disorders in coeliac disease. San Marino 10-12 April 1995
71. Collin P, Mäki M, Rasmussen M, Kyrönpalo S, Pehkonen E, Reunala T. The prevalence of adult coeliac disease in Finland. *Scand J Gastroenterol* 1995;30,Suppl 209:99. Presented at the Twenty-seventh Scandinavian congress of gastroenterology, Uppsala, Sweden; 29-31 May 1995.
72. Collin P, Vilksa S, Hällström O, Pikkarainen P. Coeliac disease in patients with infertility or spontaneous abortions. Beyond the iceberg, present and future of coeliac disease screening. Ancona, Italy October 20-21, 1995.
73. Halttunen T, Mäki M. Purified coeliac disease patient serum IgA inhibits T84 epithelial cell differentiation in vitro. *J Pediatr Gastroenterol Nutr* 1996;22:435.
74. Halttunen T, Marttinen A, Rantala I, Kainulainen H, Mäki M. TGF- $\beta$  is responsible for the fibroblast-induced differentiation of the intestinal epithelial cell line T84. *J Pediatr Gastroenterol Nutr* 1996;22:436.
75. Kaukinen K, Collin P, Holm K, Karvonen A-L, Pikkarainen P, Mäki M. Markers of coeliac disease latency in patients with positive reticulin and gliadin antibodies. In: Collin P, Mäki M, eds. *All on Coeliac Disease*. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 4.
76. Iltanen S, Holm K, Eland C, Partanen J, Mäki M. Immunohistochemical changes, serological markers and HLA genes in patients with coeliac disease and in patients suspected for coeliac disease, but having normal jejunal mucosa. In: Collin P, Mäki M, eds. *All on Coeliac Disease*. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 29.
77. Kaukinen K, Collin P, Holm K, Mäki M, Salmi J. Coeliac disease and endocrinologic disorders. In: Collin P, Mäki M, eds. *All on Coeliac Disease*. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 38.
78. Jokinen J, Peters U, Mäki M, Miettinen A, Collin P. Coeliac disease in patients with oral mucosal symptoms. In: Collin P, Mäki M, eds. *All on Coeliac Disease*. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 44.

79. Lähteenoja H, Toivanen A, Irjala K, Viander M, Mäki M, Syrjänen S. Oral mucosal changes in patients with coeliac disease. In: Collin P, Mäki M, eds. All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 44.
80. Patinen P, Savilahti E, Hietanen J, Malmström M, Mäki M, Reunala T. Intraepithelial lymphocytes bearing gamma/delta receptor in the oral and jejunal mucosa of patients with dermatitis herpetiformis. In: Collin P, Mäki M, eds. All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 45.
81. Sulkanen S, Marttinen A, Laurila K, Halttunen T, Mäki M. Endomysium autoantibodies on human umbilical cord in coeliac disease. In: Collin P, Mäki M, eds. All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 54.
82. Sulkanen S, Halttunen T, Leivo EL, Laurila K, Mäki M. Human umbilical cord-derived fibroblasts as antigen in whole cell Elisa for coeliac disease antibodies. In: Collin P, Mäki M, eds. All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 54.
83. Reunala T, Collin P, Puolakka T, Vuolteenaho N, Miettinen A, Pikkarainen P, Mäki M. Oats for patients with dermatitis herpetiformis. In: Collin P, Mäki M, eds. All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 66.
84. Halttunen T, Mäki M. Purified coeliac disease patient serum IgA inhibits fibroblast and TGF- $\beta$  induced epithelial cell (T84) differentiation in vitro. In: Collin P, Mäki M, eds. All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 72.
85. Jalava T, Polvi A, Partanen J, Marttinen A, Mäki M, Koskimies S. Immune response in vitro to human fibroblast-derived coeliac disease autoantigen is HLA DR/DQ restricted. In: Collin P, Mäki M, eds. All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 88.
86. Kerttula TO, Holm K, Polvi A, Partanen J, Mäki M. Circulating CD45RO+ TcR $\gamma\delta$  cells in coeliac disease (CoD) patients and their healthy family members. In: Collin P, Mäki M, eds. All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 88.
87. Luostarinen L, Collin P, Pirttilä T, Dastidar P, Erilä T. Nervous system manifestations in patients with coeliac disease. In: Collin P, Mäki M, eds. All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 41.
88. Keyriläinen O, Vuorinen P, Koivula T. Lactulose-mannitol test, antigliadin- and antireticulin antibodies in the screening for coeliac disease. In: Collin P, Mäki M, eds. All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 50.
89. Polvi A, Garden OA, Elwood CM, Batt RM, Partanen J. DLA-DQ alleles in Irish setter dogs with naturally occurring gluten-sensitive enteropathy. In: Collin P, Mäki M, eds. All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 92.
90. Halttunen T, Mäki M. Fibroblasts induce differentiation of the crypt-like epithelial cell line T84 in 3-dimensional co-culture. *Mol Biol Cell* 1996;7:113a.
91. Iltanen S, Holm K, Eland C, Partanen J, Mäki M. Markers of coeliac disease latency in children having normal jejunal morphology. *J Pediatr Gastroenterol Nutr* 1997;24:453.
92. Sulkanen S, Halttunen T, Leivo E-L, Laurila K, Mäki M. Whole cell ELISA for coeliac disease antibodies using human umbilical cord-derived fibroblasts as antigen. *J Pediatr Gastroenterol Nutr* 1997;24:463.
93. Iltanen S, Collin P, Korpela M, Holm K, Partanen J, Polvi A, Mäki M. Celiac disease and primary Sjögren's syndrome. American Gastroenterological Association, Washington, USA, May 12, 1997. Abstract 960.
94. Mäki M. Nutrition, small bowel mucosa, immunology and coeliac disease. Nordiske barnelegedager, Trondheim, June 11-14, 1997. Abstract I 23.
95. Halttunen T, Lindfors K, Mäki M. Coeliac disease patient serum IgA reduces T84 epithelial cell differentiation rate in vitro. FASEB Summer Research Conference 1997, Copper Mountain, Colorado, USA, August 3-8, 1997. Abstract 1:8.
96. Reunala T, Collin P, Mäki M. Gluten-free diet in dermatitis herpetiformis: With or without oats? 6th Congress of the European Academy of Dermatology and Venereology September 11-15, 1997, Dublin, Ireland. *J EADV* 1997;9;(Suppl.1):7.
97. Halttunen T, Mäki M. Celiac disease autoantibodies and tissue transglutaminase antibodies interfere similarly fibroblast-induced epithelial cell (T84) differentiation. *Gastroenterology* 1998;114:A376.
98. Kaukinen K, Collin P, Holm K, Rantala I, Vuolteenaho N, Reunala T, Mäki M. Wheat starch containing gluten-free products in the treatment of gluten intolerance. *Gastroenterology* 1998;114:A384-5.
99. Sulkanen S, Halttunen T, Korponay-Szabo I, Laurila K, Collin P, Mäki M. Tissue transglutaminase autoantibody enzyme-linked immunosorbent assay in detecting celiac disease. *Gastroenterology* 1998;114:A420.
100. Lindfors K, Halttunen T, Kainulainen H, Mäki M. Transforming growth factor- $\beta$ 1 downregulates rab11 expression in differentiating T84 intestinal epithelial cells. *Gastroenterology* 1998;114:A892.
101. Hallert C, Svensson H, Collin P, the Finnish-Swedish Study Group on Celiac Disease. Gastrointestinal symptoms and well-being of celiac patients treated for 10 years. *Gastroenterology* 1998;114:A376-7.
102. Holm K, Vuolteenaho N, Mäki M. No harm of oats in the diet of children with newly or previously diagnosed coeliac disease (CD). *J Pediatr Gastroenterol Nutr* 1998;26:549.

103. Korponay-Szabo IR, Sulkanen S, Halttunen T, Laurila K, Maurano F, Rossi M, Troncone R, Mäki M. Complete overlapping of tissue staining patterns of anti-transglutaminase antibodies and coeliac disease-specific IgA antibodies. *J Pediatr Gastroenterol Nutr* 1998;26:553.
104. Lindfors K, Halttunen T, Kainulainen H, Mäki M. rab11 expression is suppressed by TGF- $\beta$ 1 during differentiation of deep crypt T84 intestinal epithelial cells. *J Pediatr Gastroenterol Nutr* 1998;26:554.
105. Santtila S, Partanen J, Mäki M, Hurme M. Polymorphisms of the genes of the pro-inflammatory cytokines in coeliac disease: Association of the IL1A2 and TNF2 alleles. *J Pediatr Gastroenterol Nutr* 1998;26:564.
106. Sulkanen S, Laurila K, Kolho K-L, Sarnesto A, Halttunen T, Korponay-Szabo I, Savilahti E, Collin P, Mäki M. Immunological detection of tissue transglutaminase autoantibodies in coeliac disease. *J Pediatr Gastroenterol Nutr* 1998;26:566.
107. Lohiniemi S, Mäki M, Hallert C, Collin P. Quality of life in coeliac disease patients treated with gluten-free diet for 10 years. *Scand J Gastroenterol* 1998;33 Suppl 227:F2
108. Holm K, Vuolteenaho N, Mäki M. No harm of oats in the diet of children with newly or previously diagnosed coeliac disease (CD). In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;116 (A3).
109. Kaukinen K, Collin P, Holm K, Rantala I, Vuolteenaho N, Reunala T, Mäki M. Wheat starch containing gluten-free products are well tolerated in the treatment of coeliac disease. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;117 (A5).
110. Mustalahti K, Sulkanen S, Holopainen P, Laurila K, Collin P, Partanen J, Mäki M. Undetected coeliac disease in multiple case families. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;125 (A22).
111. Lohiniemi S, Mustalahti K, Collin P, Vuolteenaho N, Rostila I, Mäki M. Quality of life in Finnish coeliac disease patients. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;139 (A50)
112. Iltanen S, Holm K, Ashorn M, Ruuska T, Mäki M. The density of jejunal intraepithelial  $\gamma\delta$ + T cells increases during development of coeliac disease and decreases on gluten-free diet. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;142 (A55).
113. Kaukinen K, Salmi J, Lahtela J, Oksa H, Siljamäki-Ojansuu U, Mäki M, Collin P. Gluten-free diet does not change the metabolic control of diabetes in patients suffering from diabetes mellitus and coeliac disease. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;149 (A70).
114. Korponay-Szabo IR, Sulkanen S, Halttunen T, Laurila K, Mäki M. Coeliac disease-specific autoantibodies recognize epitopes involving the transamidase active site of the calcium-activated tissue transglutaminase. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;157 (A86).
115. Korponay-Szabo IR, Laurila K, Szalai Zs, Török E, Sulkanen S, Kovacs J, Mäki M. Comparison of transglutaminase ELISA and endomysial antibody results in children with dermatitis herpetiformis. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;158 (A87).
116. Sulkanen S, Halttunen T, Laurila K, Kolho K-L, Korponay-Szabo IR, Sarnesto A, Savilahti E, Collin P, Mäki M. Tissue transglutaminase as antigen in enzyme-linked immunosorbent assay for coeliac disease autoantibodies. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;159 (A89)
117. Polvi A, Arranz E, Fernandez-Arquero M, Collin P, Mäki M, Sanz A, Calvo C, Maluenda C, Westman P, de la Concha EG, Partanen J. HLA-DQ2-negative celiac disease in Finland and Spain. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;161 (A93).
118. Arvas M, Mustalahti K, Holopainen P, Sistonen P, Polvi A, Collin P, Mäki M, Partanen J. Evaluation of a major non-MHC candidate susceptibility locus (6p23) for celiac disease in the Finnish population. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;161 (A94).
119. Holopainen P, Mustalahti K, Collin P, Mäki M, Partanen J. HLA DQ in families with both full and clinically silent (ARA/EMA and biopsy proven) celiac disease. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;162 (A95).
120. Hakonen M, Kaukinen K, Collin P. Autoimmune endocrine diseases in patients with celiac disease. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.245.
121. Halttunen T, Mäki M. Purified coeliac disease patient serum IgA interferes with fibroblast - extracellular matrix interactions. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.47.
122. Holopainen P, Arvas M, Sistonen P, Mustalahti K, Collin P, Mäki M, Partanen J. CD28/CTLA4 gene region on chromosome 2q33 confers genetic susceptibility to celiac disease. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.53.
123. Holopainen P, Arvas M, Sistonen P, Mustalahti K, Collin P, Mäki M, Partanen J. CD28/CTLA4 gene region on chromosome 2q33 confers genetic susceptibility to celiac disease. *J Pediatr Gastroenterol Nutr* 1999;28:558.

124. Holopainen P, Arvas M, Sistonen P, Mustalahti K, Polvi A, Collin P, Mäki M, Partanen J. Linkage and family-based association study of candidate regions 11qTER, 5qTER, 6p23 and 6p21.3 in celiac disease in Finland. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.10.
125. Holopainen P, Ollikainen V, Klinger N, Mustalahti K, Mäki M, Kere J, Partanen J. Evaluation of celiac disease risk markers in a founder population. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.56.
126. Holopainen P, Ollikainen V, Klinger N, Mustalahti K, Mäki M, Kere J, Partanen J. Evaluation of celiac disease risk markers in a founder population. *J Pediatr Gastroenterol Nutr* 1999;28:557.
127. Iltanen S, Collin P, Korpela M, Holm K, Partanen J, Polvi A, Mäki M. Association between coeliac disease and Sjögren's syndrome. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.105
128. Iltanen S, Holm K, Ashorn M, Ruuska T, Mäki M. Jejunal intraepithelial  $\gamma\delta$ + T cell density increases during coeliac disease development and decreases on gluten-free diet. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.66.
129. [Kainulainen H, Rantala I, Halttunen T, Lindfors K, Collin P, Mäki M. Cyclooxygenase-2 \(COX-2\) is expressed in mucosal lesions of untreated coeliac patients. \*Immunology Letters\* 1999;69:101.](#)
130. Kaukinen K, Mäki M, Turjanmaa K, Reunala T, Karvonen A-L, Partanen J, Collin P. Abdominal complaints after ingestion of cereals. Coeliac disease or food allergy? Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.106.
131. Korponay-Szabo IR, Halttunen T, Laurila K, Szalai Zs., Török E, Sulkanen S, Szabo T, Kovacs J-B, Mäki M. Transglutaminase ELISA with human red blood cells. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.201.
132. Korponay-Szabo IR, Halttunen T, Sulkanen S, Szalai Zs, Laurila K, Török E, Kovacs JB, Mäki M. Coeliac disease-specific and coeliac disease non-specific transglutaminase autoantibodies. 7th UEGW Oral presentations. *Gut* 1999;45 (Suppl V): A27.
133. Korponay-Szabo IR, Sulkanen S, Halttunen T, Maurano F, Rossi M, Laurila K, Troncone R, Mäki M. Tissue transglutaminase is the target in rodent and primates tissues for reticulín/endomysial antibodies. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.73.
134. Lähteenoja H, Toivanen A, Viander M, Riihää I, Vilja P, Mäki M, Syrjänen S. Local challenge on oral mucosa with gliadin and a gliadin derived peptide: is it helpful as second challenge method in CD? Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.286.
135. Lähteenoja H, Viander M, Riihää I, Toivanen A, Syrjänen S, Mäki M. T cell subsets of oral mucosa in patients with coeliac disease: increase of T cell subpopulations despite treatment. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.285.
136. Lindfors K, Halttunen T, Kainulainen H, Mäki M. The expression of rab11 and TIP30/CC3 is modulated during intestinal epithelial cell differentiation. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.122.
137. Lohiniemi S, Mustalahti K, Collin P, Vuolteenaho N, Rostila I, Mäki M. Quality of life in Finnish coeliac disease patients - a national cross sectional survey. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.266.
138. Luostarinen L, Pirttilä T, Collin P. Coeliac disease presenting with neurological disorders. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.146.
139. Luostarinen L, Pirttilä T, Erilä T, Collin P. Prevalence of coeliac disease in patients with epilepsy. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.225.
140. Mäki M. Antibody testing: new diagnostic strategies. Scientific Conference on coeliac disease. Frankfurt/Main, 1999. Abstracts.
141. Mäki M. Autoantibodies and autoantigens in coeliac disease. 7th UEGW Oral presentations. *Gut* 1999;45 (Suppl V): A27.
142. Mäki M. Autoantibodies and autoantigens in coeliac disease. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.31.
143. Mäki M. Autoantibodies, autoantigens and coeliac disease. *It J Anat Embr* 1999;104 (Suppl 1):422.
144. Mustalahti K, Collin P, Sievänen H, Salmi J, Mäki M. Bone mineral density in clinically silent celiac disease. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.256.
145. Mustalahti K, Lohiniemi S, Collin P, Mäki M. Effect of gluten-free diet on quality of life of silent celiac disease (CD) patients. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.269.
146. Mustalahti K, Sulkanen S, Holopainen P, Laurila K, Collin P, Partanen J, Mäki M. Undetected coeliac disease in Finnish multiple case families. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.133.

147. Mustalahti K, Sulkanen S, Holopainen P, Laurila K, Collin P, Partanen J, Mäki M. Strategy of indentifying undetected coeliac disease in Finnish multiple case families. *J Pediatr Gastroenterol Nutr* 1999;28:562.
148. Patinen P, Aine L, Korpela M, Collin P, Reunala T. Oral findings in coeliac disease associated with Sjögren's syndrome. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.248.
149. Reunala T, Hervonen K, Collin P. Update on dermatitis herpetiformis. 8th Congress of the European Academy of Dermatology and Venereology. 3 Oct 1999. *JEAVEQ* 12 (suppl) 2 S38.
150. Reunala T, Rikkinen S, Ponkkonen H, Sihvonen M, Collin P. Long-term dietary complaints in dermatitis herpetiformis. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.270.
151. Hakanen M, Luotola K, Kaukinen K, Collin P. Clinical and subclinical autoimmune thyroid disease in patients with coeliac disease. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S21.
152. Halttunen T, Mäki M. Disrupted fibroblast - extracellular matrix interaction in celiac disease? World Congress of Pediatric Gastroenterology, Hepatology, and Nutrition. Abstracts. Boston, Massachusetts, USA, August 5-9, 2000. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl 2):S59.
153. Halttunen T, Mäki M. Disrupted fibroblast extracellular matrix interaction in celiac disease? Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S20.
154. Holm K, Vuolteenaho N, Mäki M. Two years follow up of children with celiac disease (CD) using oats in their diet. World Congress of Pediatric Gastroenterology, Hepatology, and Nutrition. Abstracts. Boston, Massachusetts, USA, August 5-9, 2000. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl 2):S284.
155. Holm K, Vuolteenaho N, Mäki M. Two years follow up of children with celiac disease (CD) using oats in their diet. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S23.
156. Holopainen P, Ollikainen V, Mustalahti K, Collin P, Mäki M, Partanen J. CTLA4/CD28 gene region in Finnish celiac disease families. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S13.
157. Iltanen S and The Coeliac Disease Study Group of Tampere. Markers of latent coeliac disease. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S23.
158. Iltanen S, Rantala I, Laippala P, Holm K, Partanen J, Mäki M. Enhanced HSP-65 expression as a marker of coeliac disease. World Congress of Pediatric Gastroenterology, Hepatology, and Nutrition. Abstracts. Boston, Massachusetts, USA, August 5-9, 2000. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl 2):S65.
159. Kainulainen H, Rantala I, Collin P, Ruuska T, Päivärinne H, Halttunen T, Lindfors K, Kaukinen K, Mäki M. COX-2 positive T-cells are present in blisters of small bowel mucosa of celiac patients. World Congress of Pediatric Gastroenterology, Hepatology, and Nutrition. Abstracts. Boston, Massachusetts, USA, August 5-9, 2000. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl 2):S9-10.
160. Kainulainen H, Rantala I, Collin P, Ruuska T, Päivärinne H, Halttunen T, Lindfors K, Kaukinen K, Mäki M. Cyclooxygenase-2 and celiac disease. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S8.
161. Karell K, Holopainen P, Mustalahti K, Collin P, Mäki M, Partanen J. Not all HLA DQ2 haplotypes confer equal susceptibility to coeliac disease: a transmission analysis in families. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S18.
162. Kaukinen K, Mäki M, Peräaho M, Collin P. A changing pattern of refractory sprue. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S20.
163. Kaukinen K, Peräaho M, Paasikivi K, Sievänen H, Mäki M, Collin P. Wheat starch-containing gluten-free products in the treatment of newly-detected coeliac disease - prospective and randomised study. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S20.
164. Korponay-Szabo I, Halttunen T, Laurila K, Szalai Z, B.Kovacs J, Collin P, Mäki M. Threshold of human transglutaminase autoantibody positivity in known coeliacs. 8th United European Gastroenterology Week, Brussels, 2000. *Gut* 2000;47, Suppl III, 198.
165. Korponay-Szabo IR, Halttunen T, Laurila K, Sulkanen S, Lörincz M, B-Kovacs J, Mäki M. Transglutaminase ELISA for IgA-deficient subjects. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S9.
166. Korponay-Szabo IR, Halttunen T, Laurila K, Sulkanen S, Lörincz M, B.Kovacs J, Mäki M. Detection of non-IgA class tissue transglutaminase autoantibodies in IgA deficient celiac patients. 6th International Symposium on Transglutaminases and Protein Cross-Linking, Lyon, France, September 16-19, 2000.
167. Korponay-Szabo IR, Halttunen T, Laurila K, Sulkanen S, Szalai Z, Dahlbom I, Kovacs JB, Mäki M. Calreticulin-bound tissue transglutaminase is less antigenic for celiac autoantibodies. World Congress of Pediatric

- Gastroenterology, Hepatology, and Nutrition. Abstracts. Boston, Massachusetts, USA, August 5-9, 2000. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl 2):S10.
168. Korponay-Szabo IR, Halttunen T, Laurila K, Sulkanen S, Szalai Zs, Dahlbom I, B-Kovacs J, Mäki M. Calcireticulin-bound tissue transglutaminase is less antigenic for celiac autoantibodies. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S9.
  169. Lähteenoja H, Viander M, Vilja P, Syrjänen S, Mäki M. Oral mucosa responds by recruitment and non-proliferative activation of T cells after submucosal challenge with a gliadin derived peptide. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S7.
  170. Lindfors K, Halttunen T, Huotari P, Nupponen N, Vihinen M, Visakorpi T, Mäki M, Kainulainen H. Identification of DED, an evolutionary conserved novel putative transcription factor. World Congress of Pediatric Gastroenterology, Hepatology, and Nutrition. Abstracts. Boston, Massachusetts, USA, August 5-9, 2000. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl 2):S280-1.
  171. Lindfors K, Halttunen T, Huotari P, Nupponen N, Vihinen M, Visakorpi T, Mäki M. Identification of DED, an evolutionary conserved novel putative transcription factor. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S22.
  172. Lindfors K, Halttunen T, Kainulainen H, Mäki M. CC3/TIP30 and rab11 - novel markers for in vitro and in vivo intestinal epithelial cell differentiation. World Congress of Pediatric Gastroenterology, Hepatology, and Nutrition. Abstracts. Boston, Massachusetts, USA, August 5-9, 2000. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl 2):S75.
  173. Lindfors K, Halttunen T, Kainulainen H, Mäki M. CC3/TIP30 and rab11: novel markers for in vitro and in vivo intestinal epithelial cell differentiation. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S21.
  174. Liu J, Juo S-H H, Holopainen P, Tong X, Grunn A, Brito ME, Mustalahti K, Mäki M, Gilliam CT, Partanen J. A genome-wide mapping on celiac disease in 60 Finnish families. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S5.
  175. Luostarinen L, Dastidar P, Collin P, Peräaho M, Eirilä T, Mäki M, Pirttilä T. Is coeliac disease associated with epilepsy and intracerebral calcifications? Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S20.
  176. Mustalahti K, Lohiniemi S, Collin P, Mäki M. Gluten-free diet improves the quality of life of silent celiac disease patients. World Congress of Pediatric Gastroenterology, Hepatology, and Nutrition. Abstracts. Boston, Massachusetts, USA, August 5-9, 2000. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl 2):S142.
  177. Mustalahti K, Lohiniemi S, Collin P, Mäki M. Improving quality of life of silent celiac disease (CD) patients during gluten-free diet warrants screening. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S7.
  178. Huotari P, Lindfors K, Halttunen T, Mäki M, Kainulainen H. Is human AATF a multifunctional regulator of epithelial cell differentiation? ESPGHAN 34th Annual Meeting, Geneva, Switzerland 9-12 May 2001. *J Pediatr Gastroenterol Nutr* 2001;32:358.
  179. Korponay-Szabo IR, Halttunen T, Laurila K, Szalai Zs, Kovacs JB, Mäki M. Capillary blood test for coeliac disease. ESPGHAN 34th Annual Meeting, Geneva, Switzerland 9-12 May 2001. *J Pediatr Gastroenterol Nutr* 2001;32:361.
  180. Korponay-Szabo IR, Szondy Zs, Laurila K, Halttunen T, Sarang Zs, Kovacs JB, Fesus L, Mäki M. Coeliac autoantibody testing on tissue transglutaminase knockout mouse tissues. ESPGHAN 34th Annual Meeting, Geneva, Switzerland 9-12 May 2001. *J Pediatr Gastroenterol Nutr* 2001;32:361.
  181. Lähdeaho M-L, Peltonen S, Collin P, Kaukinen K, Ruuska T, Haapala A-M, Mäki M. The development of small bowel lesions in coeliac disease. ESPGHAN 34th Annual Meeting, Geneva, Switzerland 9-12 May 2001. *J Pediatr Gastroenterol Nutr* 2001;32:362.
  182. Kaukinen K, Halme L, Collin P, Färkkilä M, Partanen J, Mäki M, Höckerstedt K. Coeliac disease in patients with severe liver failure. ESPGHAN 34th Annual Meeting, Geneva, Switzerland 9-12 May 2001. *J Pediatr Gastroenterol Nutr* 2001;32:385.
  183. Korponay-Szabo IR, Halttunen T, Laurila K, Szalai Zs., Kovacs JB, Mäki M. Coeliac disease autoantibody detection with patient erythrocyte transglutaminase in capillary blood. In: Mäki M, Tossavainen M, eds. Proceedings of the Workshop on Transglutaminases, Protein Cross-Linking and Coeliac Disease, September 14-15, 2001, Tampere, Finland. Coeliac Disease Study Group, University of Tampere. Tampere, 2001;173.
  184. Mustalahti K, Rannila T, Kuorilehto M, Mäki M. Finnish founder population with coeliac disease. In: Mäki M, Tossavainen M, eds. Proceedings of the Workshop on Transglutaminases, Protein Cross-Linking and Coeliac Disease, September 14-15, 2001, Tampere, Finland. Coeliac Disease Study Group, University of Tampere. Tampere, 2001;177.
  185. Mustalahti K, Rannila T, Kuorilehto M, Mäki M. Finnish founder population with coeliac disease. Abstract book of Annual Meeting of NAPGAN; Helsingör, Denmark. 30.9.-1.10. 2001.

186. Gadhavi PL, Korponay-Szabo I, Halttunen T, Mäki M. Preliminary characterisation of the IgG fraction in IgA-deficient coeliac disease patient sera. International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P76.
187. Haimila K, Holopainen P, Mustalahti K, Mäki M, Partanen J. Candidate genes CD28, CTLA4 and ICOS in celiac disease families. 10th International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002;O1.
188. Halttunen T, Huotari P, Mäki M. Mouse TG-2 antibodies and recombinant coeliac disease (CD) IgA increase fibroblast proliferation but 'natural' CD-IgA has no effect. International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P78.
189. Halttunen T, Huotari P, Mäki M. Mouse TG-2 antibodies and recombinant coeliac disease (CD) IgA increase fibroblast proliferation but "natural" CD-IgA has no effect. 7th International Conference on Transglutaminases and Protein Crosslinking Reactions. *Minerva Biotechnologica* 2002;14:226.
190. Halttunen T, Marzari R, Sblattero D, Mäki M. Biological functions of coeliac disease autoantibodies. In: Conrad K et al, eds. *From Proteomics to Molecular Epidemiology: Relevance of Autoantibodies*. Report on the 6th Dresden Symposium on Autoantibodies held in Dresden on September 4 - 7, 2002. *Autoimmunity, autoantigens, autoantibodies* 2002;3:P502.
191. Heinonen T, Lindfors K, Pasternack L, Korponay-Szabo I, Halttunen T, Kainulainen H, Mäki M. A novel glycosyltransferase isolated from differentiating T84 intestinal epithelial cells. 10th International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P45.
192. Järvinen T, Kaukinen K, Hakanen M, Mäki M, Laurila K, Collin P. Intraepithelial lymphocytes in coeliac disease. International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P125.
193. Juuti-Uusitalo K, Kainulainen H, Halttunen T, Collin P, Kaukinen K, Peräaho M, Vihinen M, Ollila J, Mäki M. Gene expression studies with cDNA microarray in jejunal biopsies and in experimental crypta-villus axis model. 10th International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002;O2.
194. Karell K, Korponay-Szabo I, Mustalahti K, Partanen J. HLA DR;DQ haplotypes do not differ between classical coeliac disease and dermatitis herpetiformis patients within families. 10th International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P15.
195. Korponay-Szabo I, Halttunen T, Szalai Z, Kovacs J, Mäki M. In vivo targeting of jejunal and extrajejunal transglutaminase 2. International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P85.
196. Korponay-Szabo I, Halttunen T, Szalai Zs, Kovacs J, Mäki M. In vivo autoantibody targeting of transglutaminase 2 in coeliac disease. 7th International Conference on Transglutaminases and Protein Crosslinking Reactions. *Minerva Biotechnologica* 2002;14:226.
197. Korponay-Szabo I, Laurila K, Szondy Z, Dahlbom I, Kovacs J, Fésüs L, Mäki M. Is endomysial antibody (EMA) positivity in patients without villous atrophy due to antibodies against transglutaminase (TG2)? International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P84.
198. Korponay-Szabo I, Laurila K, Szondy Zs, Szalai Zs, Dahlbom I, Fesus L, Mäki M. Specificity of the endomysial antibody reaction (EMA) for transglutaminase 2. 7th International Conference on Transglutaminases and Protein Crosslinking Reactions. *Minerva Biotechnologica* 2002;14:226.
199. Korponay-Szabó IR, Halttunen T, Szalai Z, Kovács JB, Mäki M. In vivo targeting of jejunal and extrajejunal tissue transglutaminase by coeliac IgA autoantibodies. ESPGHAN 35th Annual Meeting, Taormina, Italy, June 5-8, 2002. *J Pediatr Gastroenterol Nutr* 2002;34:449.
200. Lindfors K, Halttunen T, Kainulainen H, Mäki M. Wanted: AATF transcriptional regulators. 10th International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P47.
201. Mäki M, Mustalahti K, Kokkonen J, Kulmala P, Haapalahti M, Karttunen T, Ilonen J, Laurila K, Dahlbom I, Hansson T, Höpfl P, Knip M. Coeliac disease among Finnish Schoolchildren. 10th International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P21.
202. Mustalahti K, Kokkonen J, Kulmala P, Haapalahti M, Ilonen J, Karttunen T, Laurila K, Dahlbom I, Hansson T, Knip M, Mäki M. Coeliac disease among Finnish schoolchildren. ESPGHAN 35th Annual Meeting, Taormina, Italy, June 5-8, 2002. *J Pediatr Gastroenterol Nutr* 2002;34:463.
203. Niittynen M, Lindfors K, Mäki M, Kainulainen H. 3G3T, a novel transcript expressed in differentiated T84 cells. International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P49.
204. Peräaho M, Kaukinen K, Vuolteenaho N, Mäki M, Collin P. Oats and quality of life. International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P151.
205. Woolley N, Holopainen P, Ollikainen V, Mustalahti K, Mäki M, Kere J, Partanen J. Genome scan for coeliac disease in a Finnish founder family. 10th International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P30.
206. Juuti-Uusitalo Kati, Markku Mäki, Katri Kaukinen, Markku Peräaho, Juha Ollila, Pekka Collin, Mauno Vihinen, Heikki Kainulainen. cDNA MICROARRAY METHOD IN STUDYING THE COELIAC DISEASE PATHOGENESIS. IMT Research Day, Tuesday 16 December 2003, Tampere.
207. Kapitány Anikó, Korponay-Szabó Ilma, Tóth László, Niina Woolley, Jukka Partanen, Sipka Sándor. HLA-DQ allelek diagnosztikus jelentősége suboptimális módon felállított coeliakia diagnózis esetén. Hungarian Paediatric Gastroenterology Meeting, Szolnok, 2003.10.11.

208. Király Róbert, Vecsei Zsófia, Korponay-Szabó Ilma, Fésüs László. Coeliakia-autoantitestek hatása a szöveti transzglutamináz (TG2) enzimaktivitására. Hungarian Paediatric Gastroenterology Meeting, Szolnok, 2003.10.11.
209. Korponay-Szabó I, B.Kovács J, Opre J, Mäki M. Point-of-care test for coeliac disease. *Gut* 2003, 52; Suppl II. A66. 11th United European Gastroenterology Week, Madrid, 2003.
210. Korponay-Szabó I, Dahlbom I, B.Kovács J, Hansson T. IgG class anti-tissue transglutaminase antibodies and the clinical presentation type of coeliac disease. *Gut* 2003, 52; Suppl. II. A66. 11th United European Gastroenterology Week, Madrid, 2003.
211. Korponay-Szabó I, Dahlbom I, Laurila K, Koskinen S, Karell K, Partanen J, Kovács J B, Mäki M, Hansson T. IgG transzglutamináz autoantitest vizsgálat diagnosztikus szerepe szelektív IgA hiányban. Hungarian Paediatric Gastroenterology Meeting, Szolnok, 2003.
212. Korponay-Szabó I, Opre J, Laurila K, B.Kovács J, Mäki M. Helyszíni (point-of-care) coeliakia szerológiai teszt. Hungarian Paediatric Gastroenterology Meeting, Szolnok, 2003.10.11.
213. Korponay-Szabó I, Opre J, Oláh É. Tehéntejfehérje allergia a kisdedkor után. Hungarian Paediatric Gastroenterology Meeting, Szolnok, 2003.10.11.
214. Korponay-Szabó I, Halttunen, K Laurila, I. Dahlbom, L. Fésüs, M. Mäki Transglutaminase 2 antibodies: identity and in vivo action. Tampere University Hospital Research Day, 2003.01.29.
215. Korponay-Szabó Ilma, Ingrid Dahlbom, Kaija Laurila, Sinikka Koskinen, Kati Karrell, Jukka Partanen, B. Kovács Judit, Markku Mäki, Tony Hansson. IgG transzglutamináz autoantitest vizsgálat diagnosztikus szerepe szelektív IgA hiányban. Hungarian Paediatric Gastroenterology Meeting, Szolnok, 2003.10.11.
216. Korponay-Szabó Ilma: A coeliakia korszerű diagnosztikája (referátum). Coeliac disease symposium, Szolnok, 2003.10.11.
217. Korponay-Szabó IR, Dahlbom I, Laurila K, Koskinen S, Karell K, Partanen J, Kovács JB, Mäki M, Hansson T. Elevation of IgG antibodies against tissue transglutaminase as a diagnostic tool for coeliac disease in selective IgA deficiency. *J Pediatr Gastroenterol Nutr* 2003;34:Suppl 36th Annual Meeting of the ESPGHAN, Prague, 2003.
218. Opre Judit, Korponay-Szabó Ilma, Nemes Éva, Niina Woolley. Homozygota DR3;DQ2 coeliakiás család. Hungarian Paediatric Gastroenterology Meeting, Szolnok, 2003.10.11.
219. Tumpek Judit, Korponay-Szabó Ilma, Sipka Sándor. A coeliakiás ellenanyagok epitóp-specifitásának jelentősége a transzglutamináz autoantitestek diagnosztikus kimutatásában. Hungarian Paediatric Gastroenterology Meeting, Szolnok, 2003.10.11.
220. Viiri KM, Lindfors K, Laiti M, Niittynen M, Heinonen TYK, Mäki M & Kainulainen H. WHAT'SAP? mSin3A associated protein 30-like (SAP30L) - a new member in mSin3A-HDAC-corepressor complex. IMT Research Day, Tuesday 16 December 2003, Tampere.
221. Collin P, Kaukinen K, Thorell L, Mäki M. The safe threshold for gluten contamination in gluten-free products. Abstracts (P172). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
222. Collin P, Kaukinen K, Vogelsang H, Korponay-Szabó I, Sommer R, Schreier E, Volta U, Granito A, Veronesi L, Mascart F, Ocmant A, Ivarsson A, Lagerqvist C, Bürgin-Wolff A, Hadziselimovic F, Furlano R I, Sidler MA, Mulder C.J.J., Goerres MS, Mearin M.L., Ninaber M.K, Gudmand-Høyer E, Fabiani E, Catassi C, Tidlund H, Alaintalo L, Mäki M. Small bowel biopsy, anti-tissue-transglutaminase and anti-endomysial antibodies in the diagnosis of coeliac disease. A European multicentre study. Abstracts (P13). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
223. Hervonen K, M. Viljamaa, P. Collin and T. Reunala. Lymphoma in patients with dermatitis herpetiformis and their relatives. Odense 5/04, Nordic Congress of Dermatology and Venereology. Odense 2004.
224. Jarvinen TT, Collin P, Rasmussen M, Kyronpalo S, Mäki M, Partanen J, Reunala T, Kaukinen K. Villous tip intraepithelial lymphocyte count in detecting coeliac disease without villous atrophy. Abstracts (P6). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
225. Kaukinen K, Peräaho M, Collin P, Partanen J, Halttunen T, Mäki M, Korponay-Szabó I. Detection of coeliac disease without villous atrophy. Prospective and randomized study. Abstracts (P4). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
226. Korponay-Szabó IR, Halttunen T, Kovács J B, Mäki M. IgM autoantibody deposition on jejunal transglutaminase in IgA deficient coeliac patients. Abstracts (P94). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
227. Korponay-Szabó IR, Laurila K, Opre J, Kovács J B, Mäki M. Diet compliance evaluation of coeliac patients in point of care fashion. Abstracts (P15). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
228. Lähdeaho M-L, Kaukinen K, Collin P, Hartikainen M, Peltonen S, Partanen J, Haapala A-M, Mäki M. The development of small bowel lesion in coeliac disease: A long term follow-up study. Abstracts (P5). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
229. Lohi S, Mustalahti K, Laurila K, Lohi O, Rissanen H, Reunanen A, Mäki M. Prevalence and mortality of undetected coeliac disease. Abstracts (P65). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
230. Mustalahti K, Reunanen A, Heuer M, Metzger M-H, Fabiani E, Catassi C, Murray L, McMillan S, Caradonna M, Bravi E, Mäki M, and the members of the Coeliac EU Cluster, Epidemiology. Tampere, Finland, Freiburg, Germany, Ancona and Trieste, Italy and Belfast, Northern Ireland. Prevalence of coeliac disease in four European countries. Abstracts (P60). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.

231. Peräaho M, Collin P, Kaukinen K, Kekkonen L, Miettinen S, Mäki M. Oats forms an essential part of gluten-free diet in coeliac disease and dermatitis herpetiformis. Abstracts (P173). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
232. Raivio T, Kaukinen K, Korponay-Szabó I, Laurila K, Collin P, Mäki M. Rapid point of care test for coeliac disease. Abstracts (P3). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
233. Siekkinen H, Heinonen T, Mäki M, Kainulainen H. A novel glycosyltransferase, B3GTL, isolated from differentiated T84 intestinal epithelial cells. 7th annual GLIBS meeting 1-3.9.2004, Fiskebäckskil.
234. Siekkinen H, Heinonen T, Mäki M, Kainulainen H. A novel glycosyltransferase, B3GTL, isolated from differentiated T84 intestinal epithelial cells. 5th International EMBL PhD Student Symposium 2.-4.12.2004, Heidelberg.
235. Viiri K, Laiti M, Lindfors K, Lohi O, Peterson P, Mäki M, Kainulainen H. mSin3A-associated protein 30-like (SAP30L) – a new member in mSin3A-HDAC corepressor complex. 5th International EMBL PhD Students Symposium 2.-4.12.2004, Heidelberg.
236. Viljamaa M, Kaukinen K, Huhtala H, Collin P. Duration of gluten exposure does not affect the prevalence of autoimmune disorders. Abstracts (P107). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
237. Viljamaa M, Kaukinen K, Huhtala H, Kyrönpalo S, Rasmussen M, Collin P. Duration of gluten exposure does not affect the prevalence of autoimmune disorders, *Gastropäivät Tampere*, 11.-13.3.2004.
238. Juuti-Uusitalo K, Kaukinen K, Mäki M, Kainulainen H. Gene expression profiling in TGF $\beta$ -mediated T84 epithelial cell differentiation in the three-dimensional crypt-villus axis model. *Murikka Summer School*, August 21st-28th, 2005, Tampere.
239. Juuti-Uusitalo K, Kaukinen K, Mäki M, Kainulainen H. Gene expression profiling in TGF $\beta$ -mediated T84 epithelial cell differentiation in the three-dimensional crypt-villus axis model. *XX Paulo Symposium, Targeting and activation of TGF $\beta$  family growth and differentiation factors*, 2nd -3rd September 2005, Helsinki.
240. Kaukinen K, Holm K, Villanen M, Korponay-Szabo I, Mäki M. Small bowel mucosal transglutaminase autoantibodies during oats and gluten challenge in coeliac disease. Abstracts. 38th Annual Meeting of the European Society for Pediatric Gastroenterology, Hepatology and Nutrition. Porto, Portugal, June 1-4, 2005. *J Pediatr Gastroenterol Nutr* 2005;40:618 (OP2-01).
241. Kaukinen K, Viljamaa M, Huhtala H, Mäki M, Collin P. Is celiac disease screening in risk groups justified? Abstracts. 38th Annual Meeting of the European Society for Pediatric Gastroenterology, Hepatology and Nutrition. Porto, Portugal, June 1-4, 2005. *J Pediatr Gastroenterol Nutr* 2005;40:637 (PG1-14).
242. Korponay-Szabo IR, Hadjivassiliou M, Williamson CA, Laurila K, Haimila K, Woodroffe N, Sanders DS, Mäki M. Gluten ataxia: intestinal and brain targeting by tissue transglutaminase antibodies. Abstracts. 38th Annual Meeting of the European Society for Pediatric Gastroenterology, Hepatology and Nutrition. Porto, Portugal, June 1-4, 2005. *J Pediatr Gastroenterol Nutr* 2005;40:616 (OP1-02).
243. Korponay-Szabo IR, Raivio TM, Laurila K, Kovacs JB, Nemes E, Kaukinen K, Mäki M. Rapid detection of coeliac autoantibodies in the office. Abstracts. 38th Annual Meeting of the European Society for Pediatric Gastroenterology, Hepatology and Nutrition. Porto, Portugal, June 1-4, 2005. *J Pediatr Gastroenterol Nutr* 2005;40:619 (OP2-03).
244. Myrsky E, Lindfors K, Kainulainen H, Mäki M. A cell culture model for small intestinal stem cell function and differentiation. Abstracts. 38th Annual Meeting of the European Society for Pediatric Gastroenterology, Hepatology and Nutrition. Porto, Portugal, June 1-4, 2005. *J Pediatr Gastroenterol Nutr* 2005;40:661 (PG4-19).
245. Myrsky E, Lindfors K, Kainulainen H, Mäki M. Study on small intestinal epithelial stem cells in normal and coeliac disease affected small intestinal mucosa. *Kudosteknologiasymposium* 12.4.2005, Tampere.
246. Raivio T, Kaukinen K, Korponay-Szabó I, Laurila K, Mäki M. Innovative rapid point of care test in coeliac disease antibody detection. *Valtakunnalliset lääkäripäivät, Helsinki* 9.-13.1.2005.
247. Raivio T, Kaukinen K, Korponay-Szabó I, Laurila K, Mäki M. Innovative rapid point of care test in coeliac disease antibody detection. *Gastropäivät* 17.-18.2005, Jyväskylä.
248. Raivio T, Kaukinen K, Korponay-Szabó I, Laurila K, Mäki M. Innovative rapid point of care test in coeliac disease antibody detection. *Tampereen lääkäripäivät* 17.-19.3.2005, Tampere.
249. Raivio T, Kaukinen K, Laurila K, Collin P, Kovacs J, Mäki M, Korponay-Szabo IR. Self transglutaminase-based rapid coeliac antibody detection in the office. *Biannual meeting of NAPGHAN, Nordic Association for Paediatric Gastroenterology, Hepatology and Nutrition*. 24-25 November 2005, Helsinki.
250. Raivio T. Rapid point of care test in coeliac disease antibody detection. *Suomen lastenlääkäriyhdistyksen 75-vuotisjuhlakokous*, 10.-11.2.2005, Helsinki.
251. Siekkinen H, Heinonen T, Mäki M, Kainulainen H. A glycosyltransferase, B3GTL, isolated from differentiated T84 intestinal epithelial cells. *Murikka Summer School*, August 21-28, 2005, Tampere.
252. Siekkinen H, Heinonen T, Mäki M, Kainulainen H. A glycosyltransferase, B3GTL, isolated from differentiated T84 intestinal epithelial cells. *Finbionet National PhD Student Symposium*, 6-7.10.2005, Vammala.
253. Siekkinen H, Heinonen T, Mäki M, Kainulainen H. A glycosyltransferase, B3GTL, isolated from differentiated T84 intestinal epithelial cells. *IMT Research Day*, 14.12.2005, Tampere.
254. Siekkinen H, Heinonen T, Mäki M, Kainulainen H. A novel glycosyltransferase, B3GTL, isolated from differentiated T84 intestinal epithelial cells. *1st International EMBL Monterotondo International PhD Students Minisymposium, Animal Models, Tips and Tricks from Nature*. 12.-14.5.2005, Rome, Italy.

255. Stenman S, Lindfors K, Lohi O, Mäki M, Kaukinen K. Production of coeliac disease specific antibodies after in vitro gliadin challenge is dependent on small-bowel mucosal transglutaminase 2-specific IgA deposits. *IMT Research Day* 14.12.2005.
256. Stenman S, Lindfors K, Mäki M, Kaukinen K. The organ culture method in investigating the effects of wheat gliadin on coeliac disease ex vivo. *Murikka Summer School*, August 21-28 2005, Tampere.
257. Stenman S, Lindfors K, Mäki M, Kaukinen K. The organ culture method in investigating the effects of wheat gliadin on coeliac disease ex vivo. *Finbionet National PhD Student Symposium*, 6-7.10.2005, Vammala.
258. Syrjänen M, Myrsky E, Kaukinen K, Mäki M, Lindfors K. Angiogenic differentiation of endothelial and smooth muscle cells in coeliac disease. *IMT Research Day* 14.12.2005.
259. Viiri KM, Kukkonen M, Korkeamäki H, Nieminen LK, Lindfors K, Peterson P, Mäki M, Kainulainen H, Lohi O. SAP30L interacts with members of the Sin3A corepressor complex and targets Sin3A to the nucleolus. *FinBioNet PhD-Symposium 2005*, October 6-7, 2005, Vammala.
260. Viiri KM, Kukkonen M, Korkeamäki H, Nieminen LK, Lindfors K, Peterson P, Mäki M, Kainulainen H, Lohi O. SAP30L interacts with members of the Sin3A corepressor complex and targets Sin3A to the nucleolus. *IMT Research Day*, 14.12.2005, Tampere.
261. Viiri KM, Laiti M, Kokeamäki H, Nieminen LK, Lindfors K, Peterson P, Mäki M, Kainulainen H, Lohi O. SIN3A-associated protein 30-like (SAP30L) is degraded in the nucleolus and enhances the repression capability of the SIN3A-HDAC corepressor complex. *1st EMBL Monterotondo International PhD Students Minisymposium, Animal Models, Tips and Tricks from Nature*. 12. - 14.5.2005, Rome, Italy.
262. Collin P, Rondonotti E, Jacobs M, Edery J, Kaukinen K, Cellier C, Villa F, Signorelli C, Green P, Mulder C. Video capsule endoscopy in celiac disease. *Abstracts. XII International Celiac Disease Symposium*. Nov. 9-11, 2006, New York, NY.
263. Iltanen S, Raukola H, Rantala I, Kaukinen K, Collin P, Mäki M, Luukkaala T, Hurme M, Ashorn M. Elevated serum Anti-Saccharomyces Cerevisiae antibodies in celiac disease patients. *Abstracts. XII International Celiac Disease Symposium*. Nov. 9-11, 2006, New York, NY.
264. Järvinen O, Collin P, Salmi T, Iltanen S, Mäki M, Kaukinen K. Small-bowel mucosal transglutaminase-2-specific IgA-deposits as a marker of gluten sensitivity: changes with time and relationship to enteropathy. *Abstracts. XII International Celiac Disease Symposium*. Nov. 9-11, 2006, New York, NY.
265. Juuti-Uusitalo K, Mäki M, Kainulainen H, Isola J, Kaukinen K. A common set of genes regulated in epithelial cell differentiation and celiac disease. *IMT Research Day*, 19 December 2006, Tampere.
266. Juuti-Uusitalo K, Mäki M, Kainulainen H, Isola J, Kaukinen K. A common set of genes regulated in epithelial cell differentiation and celiac disease. *Abstracts. XII International Celiac Disease Symposium*. Nov. 9-11, 2006, New York, NY.
267. Juuti-Uusitalo K, Mäki M, Kainulainen H, Isola J, Kaukinen K. Common set of genes regulated in coeliac disease and in three-dimensional crypt-villus axis model. *Program and Abstract Book. 1st Nordic Meeting on Genetics and Pathogenesis of Immunological Diseases*. 4-5 September 2006, Helsinki.
268. Kaukinen K, Collin P, Laurila K, Mäki M. Resurrection of gliadin antibodies in the diagnosis of celiac disease. *Abstracts. XII International Celiac Disease Symposium*. Nov. 9-11, 2006, New York, NY.
269. Kaukinen K, Peräaho M, Lindfors K, Pikkarainen P, Karvonen A-L, Laasanen T, Sievänen H, Mäki M, Collin P. Persistent villous atrophy in symptom-free celiac disease. Does it matter? *Abstracts. XII International Celiac Disease Symposium*. Nov. 9-11, 2006, New York, NY.
270. Kaukinen K. Should we treat asymptomatic patient having coeliac disease? *Workshop CDEUSSA, Integrated knowledge on coeliac disease: identification of prolems and opportunities*. February 17-18, 2006, Naples.
271. Kekkonen L, Virta L, Kaukinen K, Reunala T. When celiac is not celiac. Causes of false diagnoses. *Abstracts. XII International Celiac Disease Symposium*. Nov. 9-11, 2006, New York, NY.
272. Kondrashova A, Mustalahti K, Kaukinen K, Viskari H, Volodicheva V, Haapala A-M, Ilonen J, Knip M, Mäki M, Hyöty H, the EPIVIR Study Group. Conspicuous difference between Russian Karelia and Finland in the prevalence of celiac disease among children. *Program and Abstract Book. 1st Nordic Meeting on Genetics and Pathogenesis of Immunological Diseases*. 4-5 September 2006, Helsinki.
273. Kondrashova A, Mustalahti K, Kaukinen K, Volodicheva v, Haapala AM, Knip M, Höty H. Difference in CD between Russian Karelia and Finland. *XII International Celiac Disease Symposium*. Nov. 9-11, 2006, New York, NY.
274. Korponay-Szabo IR, Szabados K, Pusztai J, Ludmany E, Uhrin K, Kaukinen K, Imre , Mäki M. Population screening by decentralized rapid antibody testing. *Abstracts. XII International Celiac Disease Symposium*. Nov. 9-11, 2006, New York, NY.
275. Koskinen L, Korponay-Szabo I, Kere J, Mäki M, Partanen J, Holopainen P. Linkage study of CD candidate loci in Hungarian families. *Abstracts. XII International Celiac Disease Symposium*. Nov. 9-11, 2006, New York, NY.
276. Koskinen L, Korponay-Szabo I, Mustalahti K, Lindfors K, Kaukinen K, Kurppa K, Juuti-Uusitalo K, Kere J, Wijmenga C, Mäki M, Partanen J, Holopainen P. Linkage and association study of MYO9B variants in Finnish and Hungarian populations supports the role of CELIAC4 locus (19p13) in celiac disease. *HGM2006 Workshop Abstracts. Human Genome Meeting, Helsinki* 31.5.-3.6.2006.
277. Koskinen L, Viiri K, Korponay-Szabo I, Mustalahti K, Kurppa K, Kaukinen K, Wijmenga C, Kere J, Mäki M, Partanen J, Holopainen P. Linkage and association study of MYO9B variants in Finnish and Hungarian

- populations supports the role of celiac4 locus (19P13) in celiac disease. Program and Abstract Book. 1st Nordic Meeting on Genetics and Pathogenesis of Immunological Diseases. 4-5 September 2006, Helsinki.
278. Koskinen L, Viiri K, Korponay-Szabo I, Mustalahti K, Kurppa K, Kaukinen K, Wijmenga C, Kere J, Mäki M, Partanen J, Holopainen P. Study of MYO9B variants in Finnish and Hungarian populations. Abstracts. XII International Celiac Disease Symposium. Nov. 9-11, 2006, New York, NY.
  279. Kurppa K, Mäki M, Collin P, Kaukinen K. Changing the phenotype of celiac disease after a long-term gluten challenge. Abstracts. XII International Celiac Disease Symposium. Nov. 9-11, 2006, New York, NY.
  280. Lindfors K, Myrsky E, Kainulainen H, Mäki M. A cell culture system modelling the entire small intestinal crypt villus axis. Digestive Disease Week 2006, May 20-25, 2006, Los Angeles.
  281. Lindfors K, Myrsky E, Syrjänen M, Korponay-Szabo I, Mäki M, Kaukinen K. Celiac disease IGA class autoantibodies disturb normal capillary formation in angiogenesis. Digestive Disease Week 2006, May 20-25, 2006, Los Angeles.
  282. Lohi S, Mustalahti K, Kaukinen K, Laurila K, Collin P, Rissanen H, Lohi O, Bravi E, Reunanen A, Mäki M. The prevalence of celiac disease is increasing in process of time. Digestive Disease Week 2006, May 20-25, 2006, Los Angeles.
  283. Lohi S, Mustalahti K, Kaukinen K, Laurila K, Collin P, Rissanen H, Lohi O, Bravi E, Gasparin M, Reunanen A, Mäki M. Increasing prevalence of celiac disease in process of time. 8. Gastropäivät, 16.-17.2.2006, Helsinki.
  284. Lohi S, Mustalahti K, Kaukinen K, Laurila K, collin P, Rissanen H, Lohi O, Bravi E, Gasparin M, Reunanen A, Mäki M. Increasing prevalence of celiac disease. Abstracts. XII International Celiac Disease Symposium. Nov. 9-11, 2006, New York, NY.
  285. Mäki M, Hernando A, Mena MC, Lombardia M, Lehtonen P, Pöyri S, Soininen-Tengvall P, Pajunen E, Mendez E. Barley malt beer rendered gluten free. Abstracts. XII International Celiac Disease Symposium. Nov. 9-11, 2006, New York, NY.
  286. Mäki M, Kaukinen K, Collin P, Thorell L, Hernando A, Mujico JR, Lombardia M, Mendez E. Long-term consumption of commercial oat products in coeliac disease and product contamination measured by R5-ELISA. Abstracts. XII International Celiac Disease Symposium. Nov. 9-11, 2006, New York, NY.
  287. Mustalahti K, Rask S, Mattila A, Mäki M, Joukamaa M. Depression and alexithymia in screen detected celiac disease. Abstracts. XII International Celiac Disease Symposium. Nov. 9-11, 2006, New York, NY.
  288. Myrsky E, Kaukinen K, Mäki M, Lindfors K. A cell culture system modelling the entire small intestinal crypt villus axis. 8. Gastropäivät, 16.-17.2.2006, Helsinki.
  289. Myrsky E, Kaukinen K, Syrjänen M, Korponay-Szabo IR, Mäki M, Lindfors K. Celiac disease autoantibodies disturb normal capillary formation in angiogenesis. 1st Nordic Meeting on Genetics and Pathogenesis of Immunological Diseases. 4-5 September 2006, Helsinki.
  290. Myrsky E, Kaukinen K, Syrjänen M, Korponay-Szabo IR, Mäki M, Lindfors K. Coeliac disease IgA class autoantibodies disturb normal angiogenesis. IMT Research Day, 19 December 2006, Tampere.
  291. Myrsky E, Kaukinen K, Syrjänen M, Krponay-Szabo IR, Mäki M, Lindfors K. Coeliac disease specific IgA class autoantibodies targeting transglutaminase 2 disturb normal angiogenesis. Abstracts. XII International Celiac Disease Symposium. Nov. 9-11, 2006, New York, NY.
  292. Myrsky E, Lindfors K, Kainulainen H, Kaukinen K, Mäki M. A cell culture system modelling the entire small intestinal crypt-villus axis. 20th Anniversary Symposium of the Finnish Cancer Institute: Stem cells and cancer. Biomedicum Helsinki, 1 Aug. 31-Sept. 1, 2006.
  293. Myrsky E, Syrjänen M, Kaukinen K, Korponay-Szabo I, Mäki M, Lindfors K. Celiac disease IgA class autoantibodies inhibit the recruitment of vascular smooth muscle cells to capillary walls. The 39th Annual Meeting of ESPGHAN, June 7-10, 2006, Dresden.
  294. Myrsky Essi, Kaukinen Katri, Syrjänen Mari, Korponay-Szabo Ilma R, Mäki Markku, Lindfors Katri. CELIAC DISEASE SPECIFIC IgA CLASS AUTOANTIBODIES DISTURB NORMAL ANGIOGENESIS. Lastenlääkäripäivät, Tampere 25.-27.11.2006.
  295. Raivio T, Korponay-Szabo I, Paajanen T, Iltanen S, Ashorn M, Collin P, Laurila K, Nemes E, Kovacs JB, Carrard G, Mäki M, Kaukinen K. A novel whole blood self tissue transglutaminase-based enzyme linked immunosorbent assay for celiac disease antibody detection. Abstracts. XII International Celiac Disease Symposium. Nov. 9-11, 2006, New York, NY.
  296. Salmi TT, Collin P, Järvinen O, Haimila K, Partanen J, Laurila K, Korponay-Szabo IR, Huhtala H, Reunala T, Mäki M, Kaukinen K. IgA autoantibodies against transglutaminase 2 in the small intestinal mucosa predict forthcoming coeliac disease. 8. Gastropäivät, 16.-17.2.2006, Helsinki.
  297. Siekkinen H. Glycosylation and disease: characterisation of a glycosyltransferase, B3GALTL. IMT Research Day, 19 December 2006, Tampere.
  298. Stenman S, Lindfors K, Korponay-Szabo I, Mäki M, Kaukinen K. Production of coeliac disease autoantibodies after in vitro gluten challenge is dependent on small-bowel mucosal transglutaminase 2-specific IgA deposits. The 39th Annual Meeting of ESPGHAN, June 7-10, 2006, Dresden.
  299. Stenman S, Lindfors K, Korponay-Szabo IR, Wieser H, Mäki M, Kaukinen K. Production of celiac disease autoantibodies after in vitro gliadin challenge is dependent on small-bowel mucosal transglutaminase 2-specific IgA deposits. IMT Research Day, 19 December 2006, Tampere.

300. Stenman S, Lindfors K, Korponay-Szabo IR, Wieser H, Mäki M, Kaukinen K. Production of celiac disease autoantibodies after in vitro gliadin challenge is dependent on small-bowel mucosal transglutaminase 2-specific IgA deposits. Abstracts. XII International Celiac Disease Symposium. Nov. 9-11, 2006, New York, NY.
301. Stenman SM, Lindfors K, Korponay-Szabo IR, Mäki M, Kaukinen K. Production of coeliac disease autoantibodies after in vitro gluten challenge is dependent on small-bowel mucosal transglutaminase 2-specific IgA deposits. 8. Gastropäivät, 16.-17.2.2006, Helsinki.
302. Sundman L, Saarialho-Kere U, Kaukinen K, Westerholm-Ormio M, Pulkkinen V, Kere J, Holopainen P. Expression of the novel asthma gene GPR154 in diseases of the skin and small intestine. Program and Abstract Book. 1st Nordic Meeting on Genetics and Pathogenesis of Immunological Diseases. 4-5 September 2006, Helsinki.
303. Viiri KM, Nieminen LK, Korkeamäki H, Kouvonen P, Lahesmaa R, Kainulainen H, Mäki M, Lohi O. SIN3A corepressor act on 60S ribosome biogenesis and is physically linked to the 60S pre-ribosomes by SAP30 and SAP30L proteins and this recruitment is necessary for the cell cycle progression. IMT Research Day, 19 December 2006, Tampere.
304. Vilppula A, Collin P, Mäki M, Valve R, Luostarinen M, Krekelä I, Kaukinen K, Luostarinen L. Keliakian esiintyvyys ja taudinkuva ikääntyvässä väestössä. 8. Gastropäivät, 16.-17.2.2006, Helsinki.
305. Vilppula A, Collin P, Mäki M, Valve R, Luostarinen M, Krekelä I, Patrikainen H, Kaukinen K, Luostarinen L. High prevalence of coeliac disease in elderly. Abstracts. XII International Celiac Disease Symposium. Nov. 9-11, 2006, New York, NY.
306. Einarsdottir E, Koskinen L, Korponay-Szabo I, Mustalahti K, Kurppa K, Kere J, Mäki M, Partanen J, Holopainen P. A whole-genome scan reveals significant linkage of celiac disease to 3p14-p13 and 2q14-q23 in extended pedigrees from Hungary and Finland. 40th Annual Meeting of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition, 9-12 May 2007, Barcelona, Spain.
307. Juuti-Uusitalo K, Mäki M, Kainulainen H, Isola J, Kaukinen K. Genes regulated in an in vitro epithelial cell differentiation model operate also in gluten-induced small intestinal lesion in coeliac disease. Suomen Gastroenterologiyhdistys ry, 9. Gastropäivät 15.-16.2.2007, Helsinki.
308. Juuti-Uusitalo K, Mäki M, Kainulainen H, Isola J, Kaukinen K. Genes regulated in an in vitro epithelial cell differentiation model operate also in gluten-induced small intestinal lesion in coeliac disease. 40th Annual Meeting of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition, 9-12 May 2007, Barcelona, Spain.
309. Juuti-Uusitalo K, Mäki M, Kainulainen H, Isola J, Kaukinen K. Genes regulated in an in vitro epithelial cell differentiation model operate also in gluten-induced small intestinal lesion in coeliac disease. XXXVIII Nordic Meeting of Gastroenterology, 6-8 June 2007, Reykjavik, Iceland.
310. Juuti-Uusitalo K, Mäki M, Lindfors K, Venäläinen J, Isola J, Kaukinen K. Gliadin affects on epithelial cell junctions via ErbB. IMT Research Day, University of Tampere, December 18, 2007, Tampere.
311. Kaukinen K. Diagnostic criteria for celiac disease – revision of criteria called for. “International Coeliac disease meeting” Maribor, Slovenia, September 13-16th, 2007.
312. Korponay-Szabo IR, Kovacs J, Kiraly R, Nemes E, Mäki M. High efficiency of gluten-dependent transglutaminase-specific intestinal IgA deposits as candidate diagnostic criteria in coeliac disease. 40th Annual Meeting of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition, 9-12 May 2007, Barcelona, Spain.
313. Korponay-Szabo IR, Kovacs J, Nemes E, Dahlbom I, Mäki M. Coeliac antibody testing with deamidated gliadin peptides in difficult patient samples. 40th Annual Meeting of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition, 9-12 May 2007, Barcelona, Spain.
314. Korponay-Szabo IR, Kovacs J, Kiraly R, Nemes E, Mäki M. High efficiency of gluten-dependent transglutaminase-specific intestinal IgA deposits as candidate diagnostic criteria in coeliac disease. “15th United European Gastroenterology week” held in Paris, France October 27-31th, 2007. Gut 2007;56:(Suppl III) A109.
315. Korponay-Szabo IR, Kovacs J, Nemes E, Dahlbom I, Mäki M. Coeliac disease testing with deamidated gliadin peptides in difficult patient samples. “International Coeliac disease meeting” Maribor, Slovenia, September 13-16th, 2007.
316. Korponay-Szabo IR, Kovacs J, Nemes E, Dahlbom I, Mäki M. Coeliac disease testing with deamidated gliadin peptides in difficult patient samples. “15th United European Gastroenterology week” held in Paris, France October 27-31th, 2007.
317. Korponay-Szabo IR, Vecsei Z, Kiraly R, Dahlbom I, Mäki M. Homology of deamidated gliadin peptides and tissue transglutaminase. 40th Annual Meeting of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition, 9-12 May 2007, Barcelona, Spain.
318. Korponay-Szabo IR, Vecsei Z, Kiraly R, Nemes E, Dahlbom I, Fesus L, Mäki M. Homology of deamidated gliadin peptides and tissue transglutaminase. “International Coeliac disease meeting” Maribor, Slovenia, September 13-16th, 2007.
319. Korponay-Szabo IR, Vecsei Z, Kiraly R, Nemes E, Dahlbom I, Fesus L, Mäki M. Homology of deamidated gliadin peptides and tissue transglutaminase. “15th United European Gastroenterology week” held in Paris, France October 27-31th, 2007.
320. Korponay-Szabo IR. Antibody testing in coeliac disease. “International Coeliac disease meeting” Maribor, Slovenia, September 13-16th, 2007.

321. Koskinen L, Korponay-Szabo I, Viiri K, Juuti-uusitalo K, Kaukinen K, Lindfors K, Mustalahti K, Kurppa K, Adany R, Pocsai Z, Szeles G, Wijmenga C, Mäki M, Partanen J, Kere J, Holopainen P. MYO9B gene at 19P13 (CELIACC4) shows genetic linkage and association with dermatitis herpetiformis, the skin manifestation of celiac disease. Suomen Gastroenterologiyhdistys ry, 9. Gastropäivät 15.-16.2.2007, Helsinki.
322. Kurppa K, Collin P, Viljamaa M, Haimila K, Partanen J, Saavalainen P, Laurila K, Mäki M, Kaukinen K. Treatment of endomysial antibody-positive patients without villous atrophy: are current diagnostic criteria for celiac disease valid? "International Coeliac disease meeting" Maribor, Slovenia, September 13-16th, 2007.
323. Kurppa K, Mäki M, Collin P, Reunala T, Kaukinen K. A phenotype of celiac disease may change from classical disorder to dermatitis herpetiformis after a long asymptomatic period with gluten challenge. 40th Annual Meeting of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition, 9-12 May 2007, Barcelona, Spain.
324. Lindfors K, Blomqvist T, Kaukinen K, Mäki M. Live bacteria of a probiotic bacterial strain inhibit gliadin-induced damage. 40th Annual Meeting of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition, 9-12 May 2007, Barcelona, Spain.
325. Lohi S, Mäki M, Laurila K, Montonen J, Bravi E, Gasparin M, Knekt P, Reunanen A, Kaukinen K. Malignancy in unrecognized celiac disease: population based cohort study. Suomen Gastroenterologiyhdistys ry, 9. Gastropäivät 15.-16.2.2007, Helsinki.
326. Lohi S, Mäki M, Laurila K, Montonen J, Bravi E, Gasparin M, Knekt P, Reunanen A, Kaukinen K. Malignancy in unrecognized celiac disease: population based cohort study. 40th Annual Meeting of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition, 9-12 May 2007, Barcelona, Spain.
327. Myrsky E, Kaukinen K, Syrjänen M, Korponay-Szabo IR, Mäki M, Lindfors K. Celiac disease-specific IgA class autoantibodies disturb angiogenesis. Suomen Gastroenterologiyhdistys ry, 9. Gastropäivät 15.-16.2.2007, Helsinki.
328. Myrsky E, Kaukinen K, Syrjänen M, Korponay-Szabo IR, Mäki M, Lindfors K. Coeliac disease-specific iGA class autoantibodies disturb angiogenesis. 9th International Conference of Transglutaminases and Protein Cross-linking, Marrakech September 1-5, 2007.
329. Myrsky E, Kaukinen K, Syrjänen M, Korponay-Szabo IR, Mäki M, Lindfors K. Coeliac disease-specific IgA class autoantibodies disturb angiogenesis. 8th Dresden symposium on autoantibodies, September 12-15th, 2007, Dresden, Germany.
330. Rauhavirta T, Mäki M, Kaukinen K, Lindfors K. Coeliac disease IgA - a wolf in sheep's clothing? IMT Research Day December 18, 2007, Tampere.
331. Stenman S, Lindfors K, Korponay-Szabo IR, Lohi O, Wieser H, Mäki M, Kaukinen K. Production of celiac disease autoantibodies after in vitro gliadin challenge is dependent on small-bowel mucosal transglutaminase 2-specific IgA deposits. 8th Dresden symposium on autoantibodies, September 12-15th, 2007, Dresden, Germany
332. Stenman S, Männistö P, Kaukovirta-Norja A, Lindfors K, Kaukinen K, Mäki M, Venäläinen J. Idätetyistä viljoista eristetyt entsyymit pilkkovat vehnän gliadiinin keliaakikoille vaarattomampaan muotoon in vitro. Suomen Gastroenterologiyhdistys ry, 9. Gastropäivät 15.-16.2.2007, Helsinki.
333. Stenman S, Venäläinen J, Männistö P, Lindfors K, Kaukinen K, Mäki M. Proteases from germinating cereals decrease the toxic effects of PT-gliadin in coeliac disease in vitro. 40th Annual Meeting of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition, 9-12 May 2007, Barcelona, Spain.
334. Villa F, Rondonotti E, Soada C, Cave D, Pennazio M, Riccioni ME, De Vitis I, Schneider D, Sprujevnik T, Langelier J, Arrigoni A, Laasanen T, Kaukinen K, Collin P, Costamagna G, de Franchis R. Capsule endoscopy in the diagnosis of gluten sensitive enteropathy: interim analysis of a multicenter international trial. "15th United European Gastroenterology week" held in Paris, France October 27-31th, 2007
335. Zibera F, Braicu O, Kaukinen K, Mäki M, Collin P, Holopainen P. Serum sHLA-G in celiac disease and co-associated autoimmunity. 40th Annual Meeting of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition, 9-12 May 2007, Barcelona, Spain.
336. Ashorn SM, Iltanen S, Honkanen T, Raukola H, Välineva T, Kaukinen K, Rantala I, Wei B, Braun J, Mäki M, Ashorn M. Serological responses to microbial and autoantigens in children with gut inflammation. 16<sup>th</sup> United European Gastroenterology Week, Vienna, Austria, 18-22 October 2008. Gut 2008;57(Suppl II):A125.
337. Caja S, Martucciello S, Lauvric M, Sblattero D, Marzari R, Mäki M, Kaukinen K, Lindfors K. Coeliac disease-specific autoantibodies targeted against transglutaminase 2 disturb vessel morphogenesis in vitro and in vivo through RhoB. IMT Research day 2008, December 17, Tampere.
338. Dahlbom I, Korponay-Szabo IR, Kovacs JB, Szalai Z, Mäki M. Prediction of clinical and mucosal severity of celiac disease by quantification of IgA/IgG serum antibodies to tissue transglutaminase. 16<sup>th</sup> United European Gastroenterology Week, Vienna, Austria, 18-22 October 2008. Gut 2008;57(Suppl II):A5.
339. Einarsdottir E, Dukes E, de Kauwe A, Koskinen L, Zibera F, Korponay-Szabo IR, Kurppa K, Mustalahti K, Not T, Vatta S, Ventura A, Ádány R, Pocsai Z, Széles G, Mäki M, Kere J, Anderson R, Kaukinen K, Saavalainen P. The IBD risk gene IL23R is linked to and associated with celiac disease and points to shared Th17 pathways in intestinal inflammatory diseases. World Immune Regulation Meeting II, March 17-20, 2008. Poster Abstracts WIRM II, Davos 2008:108.
340. Einarsdottir E, Koskinen LLE, Dukes E, Kainu K, Zibera F, Korponay-Szabo IR, Kurppa K, Kaukinen K, Ádány R, Not T, Ventura A, Löfberg R, Torkvist L, Bresso F, Halfvarson J, Mäki M, Saarialho-Kere U, Kere J, D'Amato M, Saavalainen P. IL23R in the Swedish, Finnish, Hungarian and Italian populations: Association with inflammatory

- bowel disease and psoriasis, and linkage to celiac disease. 58th Annual Meeting of the American Society of Human Genetics; November 11-15, 2008, Philadelphia, USA.
341. Hamilton G, Mäki M, Lähdeaho M, Bekker P, Schall T. A randomized, double-blind, placebo-controlled, phase II study testing CCX282-B (Traficet-EN) in the treatment of celiac disease. Digestive Disease Week and the 109<sup>th</sup> Annual Meeting of the AGA Institute May 17-22, 2008, San Diego, CA, USA.
  342. Kataja L, Kaukinen K, Mäki M, Lindfors K. Oats and celiac disease – study of oats-derived compounds in vitro. IMT Research day 2008, December 17, Tampere.
  343. Kaukinen K. Alkava keliakia vatsavaivojen syynä. Jatko- ja täydennyskoulutuspäivät, Krooniset vatsakivut, abstraktit. Suomen Gastroenterologiyhdistys ry, 11.-12.9.2008, Oulu, 2008:32-33.
  344. Kivistö R, Rauhavirta T, Mongeot A, Collighan R, Griffin M, Mäki M, Lindfors K. Transglutaminase 2 inhibitors – protecting by preventing? IMT Research day 2008, December 17, Tampere.
  345. Korponay-Szabo IR, Nemes E, Vecsei Z, Kiraly R, Mäki M. Rapid bedside detection of humoral IgA deficiency and celiac disease antibodies from capillary blood. 3<sup>rd</sup> World Congress of Pediatric Gastroenterology, Hepatology and Nutrition. 16-20 August 2008, Iguassu Falls, Brazil.
  346. Korponay-Szabo IR, Nemes E, Vecsei Z, Mäki M. Rapid bedside detection of humoral IgA deficiency and celiac disease antibodies from capillary blood. 16<sup>th</sup> United European Gastroenterology Week, Vienna, Austria, 18-22 October 2008. Gut 2008;57(Suppl II):A4.
  347. Korponay-Szabó I, Myrsky E, Caja S, Nemes É, Lindfors K, Mäki M. Az intracelluláris transzglutamináz vizsgálata coeliakiában (Investigation of intracellular transglutaminase in coeliac disease). Annual Paediatric Gastroenterology Meeting, 10-11 October, 2008, Budapest, Hungary.
  348. Koskinen LLE, Einarsdóttir E, Dukes E, Heap GAR, Dubois P, Korponay-Szabo IR, Kaukinen K, Kurppa K, Zibera F, Vatta S, Not T, Ventura A, Ádány R, Pocsai Z, Széles G, Mäki M, Kere J, Wijmenga C, van Heel DA, Saavalainen P. Association study of IL18RAP in three European populations with celiac disease. 58th Annual Meeting of the American Society of Human Genetics; November 11-15, 2008, Philadelphia, USA.
  349. Kurppa K, Collin P, Huhtala H, Haimila K, Saavalainen P, Partanen J, Laurila K, Mäki M, Kaukinen K. Keliakian diagnostiset kriteerit tulee uudistaa. Yleislääketieteen päivät 2008 Tampereella 23.-24.10.2008.
  350. Kurppa K, Collin P, Sievänen H, Huhtala H, Mäki M, Kaukinen K. Reduced quality of life and bone mineral density in mild enteropathy celiac disease warrants new diagnostic criteria. 16<sup>th</sup> United European Gastroenterology Week, Vienna, Austria, 18-22 October 2008. Gut 2008;57(Suppl II):A4.
  351. Kurppa K, Collin P, Viljamaa M, Haimila K, Saavalainen P, Partanen J, Laurila K, Huhtala H, Paasikivi K, Mäki M, Kaukinen K. Gluteeniherkkyys alkavassa keliakiassa: satunnaistettu, kontrolloitu kliininen tutkimus. XXIX Tampereen Lääkäripäivät Tampereella 13.-15.3.2008, luennot. Tampereen lääkäripäivät 2008 järjestelytoimikunta. Lege Artis Oy, Vammala 2008:263.
  352. Kurppa K, Collin P, Viljamaa M, Holopainen P, Haimila K, Laurila K, Paasikivi K, Mäki M, Kaukinen K. Gluten sensitivity of endomysial antibody-positive adults without villous atrophy. Should the current diagnostic criteria of celiac disease be revisited? A prospective, randomized and controlled clinical trial. Digestive Disease Week and the 109<sup>th</sup> Annual Meeting of the AGA Institute May 17-22, 2008, San Diego, CA, USA. Gastroenterology 2008;134 (Suppl 1):A-80.
  353. Kurppa K. Gluteiiniherkkyys alkavassa keliakiassa: satunnaistettu, kontrolloitu kliininen tutkimus. Lastenlääkäripäivät 6.-7.11.2008 Helsinki.
  354. Martucciello C, Caja S, Mäki M, Kaukinen K, Lindfors K. The autoantibodies anti-TG2 can modulate gene expression related with angiogenesis in endothelial cells. IMT Research day 2008, December 17, Tampere.
  355. Myrsky E, Caja S, Korponay-Szabo I, Collighan R, Mongeot A, Griffin M, Mäki M, Kaukinen K, Lindfors K. Coeliac disease autoantibodies modulate vascular permeability in vitro. IMT Research day 2008, December 17, Tampere.
  356. Myrsky E, Rauhavirta T, Kaukinen K, Korponay-Szabo IR, Mäki M, Lindfors K. Celiac disease autoantibodies targeted against transglutaminase 2 increase endothelial permeability. Helsinki Biomedical Graduate School: Cell Structure and Organelles. May 8-9, 2008, Helsinki.
  357. Myrsky E, Rauhavirta T, Kaukinen K, Korponay-Szabo IR, Mäki M, Lindfors K. Celiac disease autoantibodies targeted against transglutaminase 2 increase endothelial permeability. 5th International EDHF Symposium, June 24-27, 2008. Tampere, Finland.
  358. Myrsky E, Rauhavirta T, Korponay-Szabo I, Kaukinen K, Mäki M, Lindfors K. Celiac disease-specific autoantibodies targeted against transglutaminase 2 increase endothelial permeability in vitro. Digestive Disease Week and the 109<sup>th</sup> Annual Meeting of the AGA Institute May 17-22, 2008, San Diego, CA, USA. Gastroenterology 2008;134 (Suppl 1):A-519.
  359. Nadalutti C, Viiri K, Kaukinen K, Mäki M, Lindfors K. Downregulation of tissue transglutaminase and its implications for coeliac disease. IMT Research day 2008, December 17, Tampere.
  360. Nadalutti C, Viiri K, Kaukinen K, Mäki M, Lindfors K. G1 cell cycle arrest and anoikis induction by siRNA tissue transglutaminase in HUVECs. Murikka Summer School in Molecular Biology 2008, Cell signalling and its transcriptional readout. Tampere, August 12-19, 2008.
  361. Rauhavirta T, Kaukinen K, Mäki M, Lindfors K. Coeliac disease specific IgA enhances the transepithelial passage of gliadin peptides. 16<sup>th</sup> United European Gastroenterology Week, Vienna, Austria, 18-22 October 2008. Gut 2008;57(Suppl II):A4.

362. Rauhavirta T, Kaukinen K, Korponay-Szabó IR, Mongeot A, Collighan R, Griffin M, Mäki M, Lindfors K. The transepithelial passage of gliadin peptides is enhanced in the presence of coeliac disease-specific IgA. *IMT Research day 2008*, December 17, Tampere.
363. Rauhavirta T, Mäki M, Kaukinen K, Lindfors K. Coeliac disease specific IgA enhances the transepithelial passage of gliadin peptides. *Murikka Minisymposium Cell signalling and its transcriptional readout. Murikka Summer School in Molecular Biology 2008, Cell signalling and its transcriptional readout. Tampere, August 12-19, 2008*
364. Rauhavirta T, Mäki M, Kaukinen K, Lindfors K. Coeliac disease specific IgA enhances the transepithelial passage of gliadin peptides. *Helsinki Biomedical Graduate School, Student Council Symposium: Cell Structure and Organelles, May 8-9, 2008, Helsinki.*
365. Stenman S, Hautala A, Venäläinen J, Männistö P, Kaukovirta-Norja A, Lindfors K, Mäki M, Kaukinen K. Reducing the toxic effects of rye secalin by germinating cereal enzymes. *16<sup>th</sup> United European Gastroenterology Week, Vienna, Austria, 18-22 October 2008. Gut 2008;57(Suppl II):A5.*
366. Stenman S, Lindfors K, Venäläinen J, Männistö PT, Kaukinen K, Mäki M. Proteases from germinating wheat decrease the toxic effects of celiac disease-inducing wheat gliadin. *Digestive Disease Week and the 109<sup>th</sup> Annual Meeting of the AGA Institute May 17-22, 2008, San Diego, CA, USA. Gastroenterology 2008;134 (Suppl 1):A80.*
367. Stenman S, Hautala A, Venäläinen J, Männistö P, Kaukovirta-Norja A, Lindfors K, Mäki M, Kaukinen K. Reducing the toxic effects of rye secalin by germinating cereal enzymes. *IMT Research day 2008, December 17, Tampere.*
368. Viiri K. *8<sup>th</sup> EMBL Transcription Meeting, Heidelberg, Germany 23-27 Aug 2008.*
369. Viiri K. Nuclear monophosphoinositides regulate chromatin association and transcriptional repression of proteins of the SAP30 family. *IMT Research day 2008, December 17, Tampere.*
370. Ukkola A, Mäki M, Kurppa K, Collin P, Kekkonen L, Kaukinen K. The significance of phenotype in health related quality of life in celiac disease: a prospective follow-up study. *Science Day of the Kauppi Campus. Tampere, Jan 23, 2009.*
371. Kataja L, Kaukinen K, Mäki M, Lindfors K. Oats and celiac disease – study of oats-derived compounds in vitro. *Science Day of the Kauppi Campus. Tampere, Jan 23, 2009.*
372. Koskinen O, Collin P, Lindfors K, Laurila K, Mäki M, Kaukinen K. Usefulness of small-bowel mucosal transglutaminase-2 specific IgA autoantibody deposits in celiac disease diagnosis and follow-up. *Science Day of the Kauppi Campus. Tampere, Jan 23, 2009.*
373. Kurppa K, Collin P, Sievänen H, Saavalainen P, Huhtala H, Mäki M, Kaukinen K. Increased morbidity in mild enteropathy coeliac disease. A prospective, randomized clinical trial. *Science Day of the Kauppi Campus. Tampere, Jan 23, 2009.*
374. Kurppa K, Collin P, Viljamaa M, Haimila K, Saavalainen P, Partanen J, Laurila K, Huhtala H, Paasikivi K, Mäki M, Kaukinen K. Diagnosing mild enteropathy coeliac disease: A randomized, controlled clinical study. *Suomen lastenlääkäriyhdistyksen vuosikokous 5.-6.2.2009, Helsinki.*
375. Koskinen O, Collin P, Laurila K, Lindfors K, Mäki M, Kaukinen K. Usefulness of small-bowel mucosal transglutaminase-2 specific autoantibody deposits in coeliac disease diagnosis and follow-up. *XI Gastropäivät 12.-13.2.2009, Helsinki.*
376. Kurppa K, Collin P, Sievänen H, Saavalainen P, Huhtala H, Mäki M, Kaukinen K. Increased morbidity in mild enteropathy coeliac disease. A prospective, randomized clinical trial. *XI Gastropäivät 12.-13.2.2009, Helsinki.*
377. Ukkola A, Mäki M, Kurppa K, Collin P, Kekkonen L, Kaukinen K. The significance of phenotype in health-related quality of life in celiac disease. *XI Gastropäivät 12.-13.2.2009, Helsinki.*
378. Koskinen LLE, Einarsdóttir E, Dukes E, Heap GAR, Dubois P, Korponay-Szabo IR, Kaukinen K, Kurppa K, Zibera F, Vatta S, Not T, Ventura A, Sistonen P, Ádány R, Pocsai Z, Széles G, Mäki M, Kere J, Wijmenga C, van Heel DA, Saavalainen P. IL18RAP – a novel risk locus for coeliac disease. *XI Gastropäivät 12.-13.2.2009, Helsinki.*
379. Lohi S, Mäki M, Rissanen H, Knekt P, Reunanen A, Kaukinen K. Prognosis of unrecognized coeliac disease as regards mortality: a population based cohort study. *XI Gastropäivät 12.-13.2.2009, Helsinki.*
380. Myrsky E, Caja S, Simon-Vecsei Z, Korponay-Szabo IR, Collighan R, Mongeot A, Griffin M, Mäki M, Kaukinen K, Lindfors K. Coeliac disease IgA modulates vascular permeability *in vitro* through the activity of transglutaminase 2 and RhoA. *XI Gastropäivät 12.-13.2.2009, Helsinki.*
381. Rauhavirta T. Coeliac disease IgA enhances the transepithelial passage of gliadin peptides. *XI Gastropäivät 12.-13.2.2009, Helsinki.*
382. Ukkola A, Mäki M, Kurppa K, Collin P, Kekkonen L, Kaukinen K. Relevance of phenotypic characteristics of celiac disease in health-related quality of life: a prospective 1-year follow-up study. *13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:45-6(O1.2).*
383. Kurppa K, Collin P, Sievänen H, Saavalainen P, Huhtala H, Mäki M, Kaukinen K. Increased morbidity in mild enteropathy coeliac disease. A prospective, randomized study with special focus on symptoms, quality of life and bone mineral density. *13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:54-5(O3.5).*
384. Koskinen LLE, Einarsdóttir E, Dukes E, Heap GAR, Dubois P, Korponay-Szabo IR, Kaukinen K, Kurppa K, Zibera F, Vatta S, Not T, Ventura A, Sistonen P, Adany R, Pocsai Z, Szeles G, Mäki M, Kere J, Wijmenga C, Van Heel DA, Saavalainen P. Association study of the IL18RAP locus in three European populations with coeliac disease. *13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:59(O5.1).*
385. Hadjivassiliou M, Aeschlimann P, Sanders S, Korponay-Szabo I, Mäki M, Kaukinen K, Woodroffe N, Aeschlimann D. Antibodies against TG6 as the only serological marker of gluten ataxia. *13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:75(O9.1).*

386. Lohi S, Mäki M, Rissanen H, Knekt P, Reunanen A, Kaukinen K. Prognosis of unrecognized celiac disease as regards mortality. 13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:96(P-035).
387. Ruuskanen AT, Kaukinen K, Collin P, Huhtala H, Valve R, Mäki M, Luostarinen L. Clinical relevance of positive serum antigliadin antibodies in elderly population. 13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:98(P-039).
388. Vilppula R-L, Kaukinen K, Luostarinen L, Krekelä I, Patrikainen H, Mäki M, Collin P. The incidence and prevalence of coeliac disease in elderly people. 13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:100(P-045).
389. Collin P, Kaukinen K, Virta L. The clinical prevalence and incidence of adult coeliac disease in Finland. 13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:100-1(P-046).
390. Kurppa K, Mäki M, Saavalainen P, Partanen J, Collin P, Kaukinen K. Deamidated antigliadin antibodies are accurate markers for mild enteropathy coeliac disease. 13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:105(P-057).
391. Koskinen O, Collin P, Laurila K, Lindfors K, Mäki M, Kaukinen K. Usefulness of small-bowel mucosal transglutaminase-2 specific IgA autoantibody deposits in coeliac disease diagnosis and follow-up. 13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:106-7(P-060).
392. Bhatnagar S, Duttagupta S, Wadhwa N, Sodhi J, Bhatia V, Saini S, Sengupta S, Tandon N, Ranbir A, Jain V, Makharia G, Mäki M, Phillips AD, Not T. Point of contact testing for celiac disease in India. 13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:117(P-087).
393. Einarsdottir E, Koskinen LE, Dukes E, Kainu K, Suomela S, Lappalainen M, Zibera F, Korponay-Szabo IR, Kurppa K, Kaukinen K, Adany R, Pocsai Z, Szeles G, Färkkilä M, Turunen U, Halme L, Paavola-Sakki P, Not T, Vatta S, Ventura A, Löfberg R, Torkvist L, Bresso F, Halfvarson J, Mäki M, Kontula K, Saarialho-Kere U, Kere J, D'Amato M, Saavalainen PM. IL23R in the Swedish, Finnish, Hungarian and Italian populations: association with IBD and psoriasis, and linkage to coeliac disease. 13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:132-3(P-125).
394. Iltanen S, Ashorn S, Kurppa K, Ashorn M, Collin P, Mäki M, Wei B, Braun J, Kaukinen K. Serological responses to microbial antigens in patients with mild enteropathy coeliac disease. 13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:141(P-146).
395. Caja Galan S, Martucciello S, Lavric M, Sblattero D, Marzari R, Mäki M, Kaukinen K, Lindfors K. Coeliac disease-specific autoantibodies targeted against transglutaminase 2 disturb vessel morphogenesis in vitro and in vivo through RhoB. 13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:143(P-150).
396. Nadalutti C, Viiri K, Kaukinen K, Mäki M, Lindfors K. Downregulation of transglutaminase 2 and its implications for coeliac disease. 13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:143(P-151).
397. Martucciello S, Caja S, Mäki M, Kaukinen K, Lindfors K. The autoantibodies anti-TG2 can modulate gene expression related with angiogenesis in endothelial cells. 13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:143-4(P-152).
398. Rauhavirta T, Kaukinen K, Korponay-Szano IR, Mongeot A, Collighan R, Griffin M, Mäki M. The transepithelial passage of gliadin peptides is enhanced in the presence of coeliac disease-specific IgA. 13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:145(P-155).
399. Myrsky E, Caja S, Korponay-Szabo IR, Collighan R, Mongeot A, Griffin M, Mäki M, Kaukinen K, Lindfors K. Coeliac disease-specific autoantibodies modulate vascular permeability in vitro. 13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:146(P-158).
400. Korponay-Szabo I, Kiraly R, Nemes E, Toth B, Vecsei Z, Juuti-Uusitalo K, Csipo A, Fesus L, Mäki M. Materno-fetal transmission of celiac autoimmunity. 13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:149-50(P-168).
401. Hyytiä H, Jaatinen P, Kaukinen K, Huhtala H, Salmi J, Collin P, Metso S. Influence of gluten-free diet on autoimmune thyroiditis in celiac patients: prospective, controlled study. 13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:162(P-200).
402. Pusa E, Kaukinen K, Mäki M, Wacklin P, Partanen J, Mättö J. Composition of the mucosa-associated microbiota in the small intestine of coeliac disease patients and controls. 13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:171(P-224).
403. Korponay-Szabo I, Tumpek J, Nemes E, Papp M, Kovacs JB, Szabados K, Mäki M. Gluten-dependency of thyroid autoimmunity in celiac disease. 13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:185(P-259).
404. Stenman M, Lindfors K, Hautala A, Venäläinen J, Männistö P, Kaukovirta-Norja A, Mäki M, Kaukinen K. Reducing the toxic effects of rye secalin by germinating cereal enzymes in vitro-implications for future therapeutics in coeliac disease. 13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:193(P-278).

405. De Kauwe L, Anderson R, Kaukinen K, Tye-Din J, Kurppa K, Stewart J, Koskinen L, Mäki M, Saavalainen P. Short-term gluten challenge in Finnish patients with DQ2+ coeliac disease. 13<sup>th</sup> International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book:197(P-288).
406. Kurppa K, Mäki M, Saavalainen P, Partanen J, Collin P, Kaukinen K. Gastrointestinal symptoms, quality of life and bone mineral density in mild enteropathy coeliac disease. A prospective, controlled clinical trial. 42<sup>nd</sup> Annual Meeting of the European Society for Paediatric Gastroenterology Hepatology and Nutrition. June 3-6, 2009, Budapest. Book of Abstracts: 14 (G1-02).
407. Korponay-Szabo I, Toth B, Nemes E, Kiraly R, Vecsei Z, Juuti-Uusitalo K, Csipo A, Gyimesi J, Fesus L, Mäki M. Passive transfer model of coeliac autoimmunity in newborns from mothers with coeliac disease and transglutaminase antibodies. 42<sup>nd</sup> Annual Meeting of the European Society for Paediatric Gastroenterology Hepatology and Nutrition. June 3-6, 2009, Budapest. Book of Abstracts:20(PG1-04).
408. Kurppa K, Mäki M, Saavalainen P, Partanen J, Collin P, Kaukinen K. Antibodies against deamidated gliadin peptides in the diagnosis and follow-up of mild enteropathy coeliac disease. 42<sup>nd</sup> Annual Meeting of the European Society for Paediatric Gastroenterology Hepatology and Nutrition. June 3-6, 2009, Budapest. Book of Abstracts:24(PG1-18).
409. Korponay-Szabo I, Tumpek J, Nemes E, Kovacs JB, Szabados K, Papp M, Dinya T, Mäki M. Gluten-dependency of thyroid autoimmunity in coeliac disease. . 42<sup>nd</sup> Annual Meeting of the European Society for Paediatric Gastroenterology Hepatology and Nutrition. June 3-6, 2009, Budapest. Book of Abstracts:41(PG4-14).