

**LIST OF PUBLICATIONS  
MARKKU MÄKI**

**University of Tampere, Medical School  
Tampere University Hospital  
Tampere Finland**

	Page
<b>Thesis</b> .....	<b>2</b>
<b>Infectious diarrhoea</b> .....	<b>2</b>
<b>Pathogenesis and clinics of Yersinia infection</b> .....	<b>2</b>
<b>Coeliac disease</b> .....	<b>3</b>
<b>Clinics</b> .....	<b>3</b>
<b>Epidemiology</b> .....	<b>9</b>
<b>Genetics, immunology, pathogenesis and epithelial cell biology</b> .....	<b>10</b>
<b>Inflammatory bowel disease</b> .....	<b>14</b>
<b>Teeth and gastrointestinal diseases (includes coeliac disease papers)</b> .....	<b>15</b>
<b>Helicobacter pylori, dyspepsia and gastroesophageal reflux</b> .....	<b>15</b>
<b>Neonates in intensive care</b> .....	<b>16</b>
<b>Miscellaneous</b> .....	<b>16</b>
<b>Films</b> .....	<b>16</b>
<b>Patents</b> .....	<b>16</b>
<b>Abstracts</b> .....	<b>17</b>

## Thesis

1. Mäki M. Acute diarrhoea in children. An aetiological and clinical study of hospitalized cases. Academic dissertation, University of Tampere, Tampere, Finland. 1982.

## Infectious diarrhoea

2. Mäki M, Grönroos P, Vesikari T. Ajankohtaista lasten ripulitautien etiologiasta. (The role of bacteria (*E. coli*, *Shigella*) and viruses in the etiology of acute gastroenteritis in children, in Finnish). *Duodecim* 1978; 94:790-801.
3. Mäki M, Mäki R, Vesikari T. Faecal leucocytes in *Campylobacter*-associated diarrhoea in infants. *Acta Paediatr Scand* 1979; 68:271-2.
4. Vesikari T, Mäki M. Lasten ripulitautien mikrobiologia. (Microbial aetiology of childhood diarrhoea, in Finnish) Kirjassa: *Pediatrinen gastroenterologia. Lastentautien tutkimussäätiön julkaisuja XIII*, 1979, s. 5-11.
5. Vesikari T, Mäki M, Grönroos P. Ripulitautien etiologian selvittely (Etiology and diagnosis of gastroenteritis, in Finnish). *Suom lääk-l* 1979; 34:2581-6.
6. Mäki M, Vesikari T, Grönroos P. Enterotoxigenic and invasive *Escherichia coli* as causes of childhood diarrhoea in Finland. *Acta Paediatr Scand* 1980; 69:219-24.
7. Vesikari T, Mäki M, Kuusela A-M, Sarkkinen HK, Grönroos P. Ripulitauti sairaalainfektiona lastenosastoilla. (Diarrhoea as a nosocomial infection in children, in Finnish) *Suom lääk-l* 1980;35:1887-90.
8. Vesikari T, Mäki M, Sarkkinen HK, Arstila P, Halonen PE. Rotavirus, adenovirus, and non-viral enteropathogens in diarrhoea. *Arch Dis Child* 1981; 56:264-70.
9. Mäki M. A prospective clinical study of rotavirus diarrhoea in young children. *Acta Paediatr Scand* 1981; 70:107-13.
10. Vesikari T, Sarkkinen HK, Mäki M. Quantitative aspects of rotavirus excretion in childhood diarrhoea. *Acta Paediatr Scand* 1981; 70:717-21.
11. Mäki M, Harmoinen A, Vesikari T, Visakorpi JK. Faecal excretion of alpha-1-antitrypsin in acute diarrhoea. *Arch Dis Child* 1982; 57:154-6.
12. Vesikari T, Mäki M, Isolauri E. Lasten akuutin ripulitautien hoito. (Treatment of acute diarrhoea in children, in Finnish). *Suom lääk-l* 1983; 38:9-14.
13. Vesikari T, Mäki M, Isolauri E. Epidemiologic background for the need of rotavirus vaccine in Finland. Preliminary experience of RIT 4237 strain of live attenuated rotavirus vaccine in adults. *International Symposium on Enteric Infections in Man and Animals: Standardization of Immunological Procedures*, Dublin, Ireland, 1982, *Develop Biol Standard Vol 53*, pp. 229-36, S. Karger, Basel, 1983.
14. Mäki M, Vesikari T, Isolauri E. Muuttuva lasten ripulitauti. (Changing pattern of childhood diarrhoea, in Finnish). *Duodecim* 1983; 99:754-61.
15. Vesikari T, Isolauri E, Mäki M. Lasten akuutin ripulin avohoito. (Treatment of acute diarrhoea in children, in Finnish). *Lääketehdas Leiraksen julkaisuja XXXVIII*, 1983, pp. 64-70.
16. Vesikari T, Isolauri E, Mäki M, Grönroos P. *Clostridium difficile* in young children: Association with antibiotic usage. *Acta Paediatr Scand* 1984; 73:86-91.
17. Vesikari T, Mäki M, Baer M, Grönroos P. Pseudomembranous colitis with recurring diarrhoea and prolonged persistence of *Clostridium difficile* in a 10-year-old girl. *Acta Paediatr Scand* 1984; 73:135-7.
18. Vesikari T, Isolauri E, Mäki M. Efficacy of cholestyramine in acute infantile diarrhoea: Placebo-controlled double-blind trial in hospitalized children and in outpatients. *J Diar Dis Res* 1984; 2:151-8.
19. Mäki M. Ripulitautien mikrobidiagnostiikka. (Microbial diagnosis of diarrhoea, in Finnish). *Vox* 1984; 6:53-8.
20. Vesikari T, Isolauri E, Mäki M. Lasten akuutin ripulitautien uudet hoitolinjat. (New aspects in the treatment of acute diarrhoea in children, in Finnish). *Suom lääk-l* 1985; 40:429-36.
21. Isolauri E, Jalonen T, Mäki M. Acute gastroenteritis: Changing pattern of clinical features and management. *Acta Paediatr Scand* 1989;78:685-91.
22. Mäki M. Bakterieripulit. (Bacterial diarrhoea in children, in Finnish). In: *Lasten infektiosairaudet (Infectious diseases in children)*, O Ruuskanen, H Peltola, T Vesikari, eds. *Lastentautien tutkimussäätiö*, 1993, pp 141-45.
23. Mäki M, Baer M. Bakterieripulit. (Bacterial diarrhoea in children, in Finnish). In: *Ruuskanen O, Peltola H, Vesikari T, eds. Lasten infektiosairaudet (Infectious diseases in children)*. Pfizer Oy, Turku 2000; pp. 174-80.

## Pathogenesis and clinics of *Yersinia* infection

24. Mäki M, Grönroos P, Vesikari T. In vitro invasiveness of *Yersinia enterocolitica* isolated from children with diarrhoea. *J Infect Dis* 1978; 138:677-80.
25. Mäki M, Vesikari T, Rantala I, Grönroos P. Yersiniosis in children. *Arch Dis Child* 1980; 55:861-5.
26. Vesikari T, Granfors K, Mäki M, Grönroos P. Evaluation of ELISA in the diagnosis of *Yersinia enterocolitica* diarrhoea in children. *Acta Path Microb Scand (B)* 1980; 88:139-42.
27. Vesikari T, Nurmi T, Mäki M, Skurnik M, Sundqvist C, Granfors K, Grönroos P. Plasmids in *Yersinia enterocolitica* serotypes O:3 and O:9. Correlation with epithelial cell adherence in vitro. *Infect Immun* 1981; 33:870-6.

28. Vesikari T, Bromirska J, Mäki M. Enhancement of invasiveness of *Yersinia enterocolitica* and *Escherichia coli* in HEp-2 cells by centrifugation. *Infect Immun* 1982; 36:834-6.
29. Vesikari T, Sundqvist C, Mäki M. Adherence and toxicity of *Yersinia enterocolitica* O:3 and O:9 containing virulence-associated plasmids for various cultured cells. *Acta Path Microb Scand Sect B* 1983; 91:121-7.
30. Mäki M, Vesikari T, Rantala I, Sundqvist C, Grönroos P. Pathogenicity of 42-44 Mdal plasmid positive and negative *Yersinia pseudotuberculosis* I and *Yersinia enterocolitica* O:8 and O:9 studied in the guinea pig eye model (Serény test). *Acta Path Microb Scand Sect B* 1983; 91:241-4.
31. Vesikari T, Isolauri E, Mäki M. Clinical and laboratory features of *Yersinia*, *Campylobacter* and *Salmonella* infections in children. *Klin Pädiatr* 1985; 197:25-9.
32. Saari KM, Mäki M, Päivönsalo T, Leino R, Toivanen A. Acute anterior uveitis and conjunctivitis following yersinia infection in children. *Int Ophthalmol* 1986; 9:237-41.

## Coeliac disease

### Clinics

33. Mäki M. Guide wire manipulation of Crosby jejunal biopsy capsule. *Br Med J* 1984; 288:862-3.
34. Mäki M, Hällström O, Huupponen T, Vesikari T, Visakorpi JK. Increased prevalence of coeliac disease in diabetes. *Arch Dis Child* 1984; 59:739-42.
35. Mäki M, Hällström O, Vesikari T, Visakorpi JK. Evaluation of a serum IgA class reticulin antibody test for the detection of childhood celiac disease. *J Pediatr* 1984; 105:901-5.
36. Vainio E, Kosnai I, Hällström O, Karpati S, Mäki M, Reunala T. Antigliadin and antireticulin antibodies in children with dermatitis herpetiformis. *J Pediatr Gastroent Nutr* 1986; 5:735-9.
37. Teppo A-M, Mäki M, Hällström O, Maury CPJ. Antibodies to 90 kilodalton glycoprotein in childhood and adolescent celiac disease: Relationship to reticulin antibodies. *J Pediatr Gastroent Nutr* 1987; 6:908-14.
38. Auricchio S, Mazzacca G, Tosi R, Visakorpi J, Mäki M, Polanco I. Coeliac disease as a familial condition: Identification of asymptomatic coeliac patients within family groups. *Gastroenterol Int* 1988; 1:25-31.
39. Mäki M, Hällström O, Verronen P et al. Reticulin antibody, arthritis, and coeliac disease in children. *Lancet* 1988; I:479-80.
40. Mäki M, Kallonen K, Lähdeaho M-L, Visakorpi JK. Changing pattern of childhood coeliac disease in Finland. *Acta Paediatr Scand* 1988; 77:408-12.
41. Ståhlberg M-R, Hietanen E, Mäki M. Mucosal biotransformation rates in the small intestine of children. *Gut* 1988; 29:1058-63.
42. Collin P, Salmi J, Hällström O, Oksa H, Oksala H, Mäki M, Reunala T. High frequency of coeliac disease in adult patients with type I diabetes. *Scand J Gastroenterol* 1989; 24:81-4.
43. Mäki M, Lähdeaho M-L, Hällström O, Viander M, Visakorpi JK. Postpubertal gluten challenge in coeliac disease. *Arch Dis Child* 1989;64:1604-07.
44. Collin P, Hällström O, Mäki M, Viander M, Keyriläinen O. Atypical coeliac disease found with serologic screening. *Scand J Gastroenterol* 1990;25:245-50.
45. Mäki M. Keliakiatauti (Coeliac disease, in Finnish). *Suomen Hammläkl* 1990;37:544-47.
46. Mäki M, Holm K, Koskimies S, Hällström O, Visakorpi JK. Normal small bowel biopsy followed by coeliac disease. *Arch Dis Child* 1990;65:1137-41.
47. Koger O, Uibo R, Mäki M. Coeliac disease in children. (In Russian.). *Pediatrics* 1991;10:79-85.
48. Mäki M, Holm K, Lipsanen V, Hällström O, Viander M, Collin P, Savilahti E, Koskimies S. Serological markers and HLA genes among healthy first-degree relatives of patients with coeliac disease. *Lancet* 1991;338:1350-53.
49. Mäki M, Holm K, Koskimies S, Visakorpi JK. Latent coeliac disease. In: Mearin ML, Mulder CJJ, eds. *Coeliac Disease, 40 years gluten-free*. Kluwer Academic Publishers, Dordrecht, 1991:153-6.
50. Collin P, Korpela M, Hällström O, Viander M, Keyriläinen O, Mäki M. Rheumatic complaints as a presenting symptom in patients with coeliac disease. *Scand J Rheumatol* 1992;21:20-3.
51. Collin P, Mäki M, Keyriläinen O, Hällström O, Reunala T, Pasternack A. Selective IgA-deficiency and coeliac disease. *Scand J Gastroenterol* 1992;27:367-71.
52. Mäki M, Hällström O, Marttinen A, Lipsanen V, Viander M, Holm K, Collin P, Koskimies S. Screening tools for use in coeliac disease. In: Auricchio S, Visakorpi JK, eds. *Common Food Intolerances 1: Epidemiology of Coeliac Disease*. *Dyn Nutr Res*, Basel, Karger 1992;2:93-104.
53. Mäki M. Use of serological antibody tests in celiac disease. In: Branski D, Rozen P, Kagnoff M, eds. *Gluten-Sensitive enteropathy*. *Front Gastrointest Res*. Basel, Karger 1992;19:108-29. (review, includes original material).
54. Ascher H, Holm K, Kristiansson B, Mäki M. Different features of coeliac disease in two neighbouring countries. *Arch Dis Child* 1993;69:375-80.
55. Collin P, Helin H, Mäki M, Hällström O, Karvonen AL. Follow-up of patients positive in reticulin and gliadin antibody tests with normal small-bowel biopsy findings. *Scand J Gastroenterol* 1993;28:595-98.
56. Reunala T, Mäki M. Dermatitis herpetiformis: A genetic disease. *Eur J Dermatol* 1993;3:519-26.

57. Collin P, Mäki M. Associated disorders in coeliac disease. Clinical aspects. *Scand J Gastroenterol* 1994;29:769-75.
58. Visakorpi JK, Mäki M. Changing clinical features of coeliac disease. *Acta Paediatr* 1994;Suppl 395:10-13.
59. Mäki M, Hällström O, Sulkanen S, Marttinen A. Antibodies in coeliac disease. In: Auricchio S, Ferguson A, Troncone R, eds: *Mucosal Immunity and the Gut Epithelium: Interactions in Health and Disease*. Dyn Nutr Res, Basel, Karger 1995;4:115-26.
60. Mäki M, Huupponen T, Holm K, Hällström O. Seroconversion of reticulín autoantibodies predicts coeliac disease in insulin-dependent diabetes mellitus. *Gut* 1995;36:239-42.
61. Mäki M. Keliakian diagnostiikka (Diagnosis of coeliac disease, in Finnish). *Moodi* 1995;19 (1):34-35.
62. Visakorpi JK, Mäki M. Çöliak hastaliginin degisen klinik özellikleri. (Changing clinical features of coeliac disease, in Hungarian.) *Cocuk Sagligi ve Hastaliklari Dergisi* 1995;38:103-109.
63. Mäki M. Keliakian periytyvyys. (Inheritance of coeliac disease, in Finnish.) *Keliakia* 1996;2:5-7.
64. Mäki M. Onko suomalainen kaura vehnätöntä. (Is Finnish oats gluten-free, in Finnish.) *Keliakia* 1996;2:8.
65. Visakorpi JK, Mäki M. Celiakia - zmieniający się obraz kliniczny. (Changing clinical features of coeliac disease, in Polish). *Medycyna Praktyczna* 1996;WS 3:32-5.
66. Collin P, Holm K, Kaukinen K, Laurila K, Mäki M. Latentti keliakia. (Latent coeliac disease, in Finnish.) *Suom lääkl* 1997;52:4279-83.
67. Collin P, Julkunen R, Lehtola J, Mäki M, Rasmussen M, Reunala T, Savilahti E, Uusitupa M, Vuoristo M. Keliakian hoitosuositus. (Diagnosis and treatment recommendations of coeliac disease, in Finnish.) *Duodecim* 1997;113:82-7.
68. Collin P, Mäki M. Celiac disease - even a neurological disorder. *J Pediatr Gastroenterol Nutr* 1997;24:116-7.
69. Collin P, Mäki M. Non-gastrointestinal manifestations of coeliac disease. In: Gobbi G et al, eds. *Epilepsy and other neurological disorders in coeliac disease*. London: John Libbey & Company, 1997:13-18.
70. Collin P, Reunala T, Mäki M. New diagnostic strategy to find coeliac disease. In: Mäki M, Collin P, Visakorpi JK, eds. *Coeliac disease. Proceedings of the Seventh International Symposium on Coeliac Disease*. Coeliac Disease Study Group, University of Tampere. Tampere 1997;47-51.
71. Marttinen A, Sulkanen S, Mäki M. Human fibroblast-derived molecules as antigens in enzyme-linked immunosorbent assay for celiac disease-specific IgA. *Eur J Clin Invest* 1997;27:135-40.
72. Mäki M, Collin P. Coeliac disease. *Lancet* 1997;349:1755-9.
73. Mäki M, Laurila K. Keliakia. *Moodi* 1997;21:247-50.
74. Mäki M. Keliakia ja lehmänmaidon sietämättömyys. (Coeliac disease and cow's milk intolerance, in Finnish). Kirjassa: Simell O, toim. *Neuvolakirja*. Orion, Vammala, 1997:447-53.
75. Mäki M. Keliakian hoito puhuttaa lähitulevaisuudessa - vehnätkkelystä vai ei. Pääkirjoitus. (Leading article, in Finnish). *Keliakia* 1997;4:3.
76. Mäki M. Puheenjohtajan palsta. Pääkirjoitus. (Leading article, in Finnish.) *Keliakia* 1997;1:3.
77. Patinen P, Savilahti E, Hietanen J, Malmström M, Mäki M, Reunala T. Intraepithelial lymphocytes bearing the  $\gamma/\delta$  receptor in the oral and jejunal mucosa in patients with dermatitis herpetiformis. *Eur J Oral Sci* 1997;105:130-5.
78. Collin P, Mäki M. Ohutsuoli. Keliakia. (Coeliac disease, in Finnish.) In: Höckerstedt K, Färkkilä M, Kivilaakso E, Pikkarainen P, eds. *Gastroenterologia*. Duodecim, Jyväskylä 1998;265-74.
79. Jokinen J, Peters U, Mäki M, Miettinen A, Collin P. Coeliac sprue in patients with chronic oral mucosal symptoms. *J Clin Gastroenterol* 1998;26:23-6.
80. Kaukinen K, Collin P, Holm K, Karvonen A-L, Pikkarainen P, Mäki M. Small-bowel mucosal inflammation in reticulín or gliadin antibody-positive patients without villous atrophy. *Scand J Gastroenterol* 1998;33:944-9.
81. Lohiniemi S, Mustalahti K, Collin P, Mäki M. Measuring quality of life in coeliac disease patients. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;73-7.
82. Lähteenoja H, Toivanen A, Viander M, Mäki M, Irjala K, Rähä I, Syrjänen S. Oral mucosal changes in coeliac patients on a gluten-free diet. *Eur J Oral Sci* 1998;106:899-906.
83. Mäki M, Collin P. Coeliac disease. Reply. *Lancet* 1998;351:63.
84. Mäki M, Collin P. Enfermedad Celíaca. (In Spanish). *Tribuna Celíaca* 1998;6-14.
85. Mäki M, Kaukinen K, Holm K, Collin P. Treatment of coeliac patients with oats and wheat starch. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;93-6.
86. Mäki M, Sulkanen S, Collin P. Antibodies in relation to gluten intake. *Digestive Diseases* 1998;16:330-2.
87. Mäki M. Celiac disease. *Pediatrics Review Series. Food Allergy and Gastroenterology* 1998;2:2-5.
88. Mäki M. Changing features of coeliac disease. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;1-6.
89. Mäki M. Coeliac disease in Finland. In: Banchini G, Donadio A, eds. *La Malattia Celiaca (alla ricerca del tempo perduto)*. 8 Incontro Internazionale "Nuove Tecnologie in Pediatria". Divisione di Pediatria Arcispedale S. Maria Nuova, Associazione Italiana Celiachia. Reggio Emilia 1998; 43-8.
90. Mäki M. Keliakia tunnetuksi. Pääkirjoitus (Leading article, in Finnish). *Keliakia* 1998;4:3.
91. Mäki M. Keliakia yleistyy ja muuttaa muotoaan. *Keliakia* 1998;4:12.

92. Mäki M. Keliakiaseulonnan haitat ja edut punnittava. *Keliakia* 1998;4:13.
93. Reunala T, Collin P, Holm K, Pikkarainen P, Miettinen A, Vuolteenaho N, Mäki M. Tolerance to oats in dermatitis herpetiformis. *Gut* 1998;43:490-3.
94. Sulkanen S, Collin P, Laurila K, Mäki M. IgA- and IgG-class antihuman umbilical cord antibody tests in adult coeliac disease. *Scand J Gastroenterol* 1998;33:251-4.
95. Sulkanen S, Halttunen T, Laurila K, Kolho K-L, Korponay-Szabo IR, Sarnesto A, Savilahti E, Collin P, Mäki M. Tissue transglutaminase autoantibody enzyme-linked immunosorbent assay in detecting celiac disease. *Gastroenterology* 1998;115:1322-8.
96. Sulkanen S, Halttunen T, Marttinen A, Leivo E-L, Laurila K, Mäki M. Autoantibodies in celiac disease: Importance of fibroblasts. *J Pediatr Gastroenterol Nutr* 1998;27:206-13.
97. Collin P, Kaukinen K, Mäki M. Clinical features of celiac disease today. *Digestive Diseases* 1999;17:100-6.
98. Iltanen S, Collin P, Korpela M, Holm K, Partanen J, Polvi A, Mäki M. Celiac disease and markers of celiac disease latency in patients with primary Sjögren's syndrome. *Am J Gastroenterol* 1999;94:1042-6.
99. Kaukinen K, Collin P, Holm K, Rantala I, Vuolteenaho N, Reunala T, Mäki M. Wheat starch-containing gluten-free flour products in the treatment of coeliac disease and dermatitis herpetiformis. A long-term follow-up study. *Scand J Gastroenterol*, 1999;34:163-9.
100. Kaukinen K, Collin P, Mykkänen A-H, Partanen J, Mäki M, Salmi J. Celiac disease and autoimmune endocrinologic disorders. *Dig Dis Sci* 1999;44:1428-33.
101. Mäki M. Is coeliac disease a syndrome? In: UEGW Roma'99. ESPGHAN Postgraduate Course "Gastrointestinal uncertainties". 1999, p. 20-22.
102. Mustalahti K, Collin P, Sievänen H, Salmi J, Mäki M. Osteopenia in patients with clinically silent coeliac disease warrants screening. *Lancet* 1999;354:744-5.
103. Collin P, Mäki M. Ohutsuoli. Keliakia. (Coeliac disease, in Finnish.) In: Höckerstedt K, Färkkilä M, Kivilaakso E, Pikkarainen P, eds. *Gastroenterologia. Duodecim, Jyväskylä* 2000;266-75.
104. Janatuinen E, Kemppainen T, Pikkarainen P, Holm K, Kosma V-M, Uusitupa M, Mäki M, Julkunen R. Lack of cellular and humoral immunological responses to oats in adults with coeliac disease. *Gut* 2000;46:327-31.
105. Kaukinen K, Turjanmaa K, Mäki M, Partanen J, Venäläinen R, Reunala T, Collin P. Intolerance to cereals is not specific for coeliac disease. *Scand J Gastroenterol* 2000;35:942-6.
106. Kokkonen J, Holm K, Karttunen TJ, Mäki M. Children with untreated food allergy express a relative increment in the density of duodenal  $\gamma\delta$ + T cells. *Scand J Gastroenterol* 2000;35:1137-42.
107. Lähteenoja H, Mäki M, Viander M, Rähä I, Vilja P, Rantala I, Toivanen A, Syrjänen S. Local challenge on oral mucosa with an  $\alpha$ -gliadin related synthetic peptide in patients with celiac disease. *Am J Gastroenterol* 2000;95:2880-7.
108. Lähteenoja H, Mäki M, Viander M, Toivanen A, Syrjänen S. Local challenge of oral mucosa with gliadin in patients with coeliac disease. *Clin Exp Immunol* 2000;120:38-45.
109. Lähteenoja H, Toivanen A, Viander M, Rähä I, Rantala I, Syrjänen S, Mäki M. Increase in T-cell subsets of oral mucosa: a late immune response in patients with treated coeliac disease? *Scand J Immunol* 2000;52:602-8.
110. Lohiniemi S, Mäki M, Kaukinen K, Laippala P, Collin P. Gastrointestinal symptoms rating scale in coeliac disease patients on wheat starch-based gluten-free diets. *Scand J Gastroenterol* 2000;35:947-9.
111. Mäki M. Keliakian diagnostiikka tänään. *Pharmacia Now*, 4/2000.
112. Stern M, Working Group on Serologic Screening for Celiac Disease (Andersson C, Bürgin-Wolff A, Grodzinsky E, Kapitzka S, Laurila K, Lütke R, Mäki M, Mascart F, Neunhoffer E, Ocmant A, Savilahti E, Volta U, Wäscher I, Wruk M, Zalewski T). Comparative evaluation of serologic tests for celiac disease: a European initiative toward standardization. *J Pediatr Gastroenterol Nutr* 2000;31:513-9.
113. Collin P, Hakanen M, Salmi J, Mäki M, Kaukinen K. Autoimmune hypopituitarism in patients with coeliac disease - symptoms confusingly similar. *Scand J Gastroenterol* 2001;36:558-60.
114. Fabiani E, Catassi C and the International Working Group on Eu-tTG (incl. Sulkanen S, Mäki M). The serum IgA class anti-tissue transglutaminase antibodies in the diagnosis and follow up of coeliac disease. Results of an international multi-centre study. *Eur J Gastroenterol Hepatol* 2001;13:659-65.
115. Kaukinen K, Mäki M, Collin P. Compliance to a stright gluten-free diet : Are dietary transgressions detrimental? *Food Allergy Intolerance* 2001;15:27.
116. Kaukinen K, Mäki M, Collin P. Physiopathology of celiac disease. In: Morteau O, ed. *The intestinal mucosa and dietary antigens/ Oral Tolerance. Landes Bioscience, Georgetown*, 2001;Chapter 10:1-13.
117. Kaukinen K, Mäki M, Partanen J, Sievänen H, Collin P. Coeliac disease without villous atrophy: revision of criteria called for. *Dig Dis Sci* 2001;46:879-87.
118. Kokkonen J, Haapalahti M, Laurila K, Karttunen T, Mäki M. Cow's milk protein-sensitive enteropathy at school age. *J Pediatr* 2001;139:797-803.
119. Luostarinen L, Collin P, Peräaho M, Mäki M, Pirttilä T. Coeliac disease in patients with cerebellar ataxia of unknown origin. *Ann Med* 2001;33:445-9.
120. Luostarinen L, Dastidar P, Collin P, Peraaho M, Maki M, Erila T, Pirttila T. Association between coeliac disease, epilepsy and brain atrophy. *Eur Neurol* 2001;46:187-91.
121. Mäki M. Celiac disease. In: Delvin EE, Lentze MJ, eds. *Gastrointestinal functions. Nestlé Nutrition Workshop Series, Pediatric Program Vol. 46. Vevey* 2001;257-74.

122. Mäki M. Minne menet, keliakia? *Keliakia* 2001;4:17-18.
123. Collin P, Kaukinen K, Mäki M, Vehmainen P, Partanen J, Färkkilä M, Halme L, Höckerstedt K. Discussion on celiac disease in patients with severe liver disease: Gluten-free diet may reverse hepatic failure. Reply. *Gastroenterology* 2002;123:2159-60.
124. Hill I, Bhatnagar S, Cameron DJ, De Rosa S, Mäki M, Russell GJ, Troncone R. Report of the Celiac Disease: Working Group Report of the First World Congress of Pediatric Gastroenterology, Hepatology and Nutrition. *J Pediatr Gastroenterol Nutr* 2002;35,Suppl2:S78-88.
125. Janatuinen EK, Kempainen TA, Julkunen RJK, Kosma V-M, Mäki M, Heikkinen M, Uusitupa MIJ. No harm from five year ingestion of oats in coeliac disease. *Gut* 2002;50:332-5.
126. Kaukinen K, Halme L, Collin P, Färkkilä M, Mäki M, Vehmanen P, Partanen J, Höckerstedt K. Celiac disease in patients with severe liver disease: gluten-free diet may reverse hepatic failure. *Gastroenterology* 2002;122:881-8.
127. Kaukinen K, Sulkanen S, Mäki M, Collin P. IgA-class transglutaminase antibodies in evaluating the efficacy of gluten-free diet in coeliac disease. *Eur J Gastroenterol Hepatol* 2002;14:311-5.
128. Kokkonen J, Ruuska T, Karttunen TJ, Mäki M. Lymphonodular hyperplasia of the terminal ileum associated with colitis shows an increased  $\gamma\delta$ + T-cell density in children. *Am J Gastroenterol* 2002;97:667-72.
129. Mustalahti K, Lohiniemi S, Collin P, Vuolteenaho N, Laippala P, Mäki M. Gluten-free diet and quality of life in patients with screen-detected celiac disease. *Effective Clinical Practice* 2002;5:105-13.
130. Mustalahti K, Sulkanen S, Holopainen P, Laurila K, Collin P, Partanen J, Mäki M. Coeliac disease among healthy members of multiple case coeliac disease families. *Scand J Gastroenterol* 2002;37:161-5.
131. Järvinen TT, Kaukinen K, Laurila K, Kyrönpalo S, Rasmussen M, Mäki M, Korhonen H, Reunala T, Collin P. Intraepithelial lymphocytes in celiac disease. *Am J Gastroenterol* 2003;98:1332-7.
132. Kaukinen K, Collin P, Mäki M, Partanen J, Halme L, Färkkilä M, Höckerstedt K. Screening for celiac disease in patients with chronic liver disease. Reply. *Gastroenterology* 2003;125:1289-90.
133. Mäki M. Strategies for screening for coeliac disease. In: Cerf-Bensussan N, Brousse N, Caillat-Zucman S, Cellier C, Schmitz J, eds. *Coeliac disease. Proceedings of the Xth International Symposium on Coeliac Disease, Paris, June 2-5, 2002.* John Libbey Eurotext. Montrouge 2003;191-8.
134. Peräaho M, Kaukinen K, Paasikivi K, Sievänen H, Lohiniemi S, Mäki M, Collin P. Wheat-starch-based gluten-free products in the treatment of newly detected coeliac disease. Prospective and randomized study. *Aliment Pharmacol Ther* 2003;17:587-94.
135. Collin P, Mäki M, Kaukinen K. Complete small intestinal mucosal recovery is obtainable in the treatment of celiac disease. *Gastrointest Endosc* 2004;59:158-9.
136. Collin P, Mäki M, Kaukinen K. It is the compliance, not milligrams of gluten, that is essential in the treatment of celiac disease. Letter to the Editor. *Nutrition Reviews* 2004;62:490.
137. Collin P, Thorell L, Kaukinen K, Mäki M. The safe threshold for gluten contamination in gluten-free products. Can trace amounts be accepted in the treatment of coeliac disease? *Aliment Pharmacol Ther* 2004;19:1277-83.
138. Järvinen TT, Collin P, Rasmussen M, Kyrönpalo S, Mäki M, Partanen J, Reunala T, Kaukinen K. Villous tip intraepithelial lymphocytes as markers of early-stage coeliac disease. *Scand J Gastroenterol* 2004;39:428-33.
139. Kokkonen J, Holm K, Karttunen TJ, Mäki M. Enhanced local immune response in children with prolonged gastrointestinal symptoms. *Acta Paediatr* 2004;93:1601-7.
140. Lohi O, Mäki M. Vehnää keliakikolle? *Duodecim* 2004;120:645-6.
141. Mäki M, Lohi O. Celiac disease. In: Walker et al, eds. *Pediatric Gastrointestinal Disease. Pathophysiology, diagnosis, management.* 4<sup>th</sup> Edition. BC Decker Inc. Hamilton, 2004;932-43.
142. Peräaho M, Collin P, Kaukinen K, Kekkonen L, Miettinen S, Mäki M. Oats can diversify a gluten-free diet in celiac disease and dermatitis herpetiformis. *J Am Diet Assoc* 2004;104:1148-50.
143. Peräaho M, Kaukinen K, Mustalahti K, Vuolteenaho N, Mäki M, Laippala P, Collin P. Effect of an oats-containing gluten-free diet on symptoms and quality of life in coeliac disease. A randomised study. *Scand J Gastroenterol* 2004;39:27-31.
144. Troncone R, Bhatnagar S, Butzner D, Cameron D, Hill I, Hoffenberg E, Mäki M, Mendez V, de Jimenez MZ. Celiac disease and other immunologically mediated disorders of the gastrointestinal tract: Working Group Report of the Second World Congress of Pediatric Gastroenterology, Hepatology and Nutrition. *J Pediatr Gastroenterol Nutr* 2004;39:S601-10.
145. Collin P, Julkunen R, Lehtola J, Kaukinen K, Mäki M, Rasmussen M, Reunala T, Savilahti E, Uusitupa M, Vuoristo M, Vuorio A. Keliakia. Käypä hoito-suosituksen päivitys. *Duodecim* 2005;121:2705-7.
146. Collin P, Kaukinen K, Vogelsang H, Korponay-Szabó I, Sommer R, Schreier E, Volta U, Granito A, Veronesi L, Mascart F, Ocmant A, Ivarsson A, Lagerqvist C, Bürgin-Wolff A, Hadziselimovic F, Furlano R I, Sidler M A, Mulder C J J, Goerres M S, Mearin ML, Ninaber MK, Gudmand-Hoyer E, Fabiani E, Catassi C, Tidlund H, Alaintalo L, Mäki M. Antiendomysial and antihuman recombinant tissue transglutaminase antibodies in the diagnosis of coeliac disease. A biopsy-proven European multicentre study. *Eur J Gastroenterol Hepatol* 2005;17:85-91.
147. Haapalahti M, Kulmala P, Karttunen TJ, Paajanen L, Laurila K, Mäki M, Mykkänen H, Kokkonen J. Nutritional status in adolescents and young adults with screen-detected celiac disease. *J Pediatr Gastroenterol Nutr* 2005;40:566-70.

148. Hernando A, Garcia E, Llorente M, Mujico JR, Lombardia M, Mäki M, Kaukinen K, Collin P, Mendez E. Measurements of hydrolysed gliadins in malts, breakfast cereals, heated/hydrolysed foods, whiskies and beers by means of a new competitive R5 ELISA. In: Stern M, ed. Proceedings of the 19<sup>th</sup> Meeting, Working Group on Prolamin Analysis and Toxicity. Verl. Wissensch. Scripten, Zwickau 2005;31-7.
149. Kaukinen K, Mäki M, Collin P. Trace amounts of gluten in the treatment of coeliac disease. In: Stern M, ed. Proceedings of the 19<sup>th</sup> Meeting Working Group on Prolamin Analysis and Toxicity. Verl. Wissensch. Scripten. Zwickau 2005;115-6.
150. Kaukinen K, Peräaho M, Collin P, Partanen J, Woolley N, Kaartinen T, Nuutinen T, Halttunen T, Mäki M, Korponay-Szabo I. Small-bowel mucosal transglutaminase 2-specific IgA deposits in coeliac disease without villous atrophy: A prospective and randomized clinical study. *Scand J Gastroenterol* 2005;40:564-72.
151. Korponay-Szabó I, Raivio T, Laurila K, Opre J, Király R, Kovács JB, Kaukinen K, Fésüs L, Mäki M. Celiac disease case finding and diet monitoring by point-of-care testing. *Aliment Pharmacol Ther* 2005;22:729-37.
152. Lähdeaho M-L, Kaukinen K, Collin P, Ruuska T, Partanen J, Haapala A-M, Mäki M. Celiac disease: From inflammation to atrophy: A long-term follow-up study. *J Pediatr Gastroenterol Nutr* 2005;41:44-8.
153. Mujico JR, Hernando A, Lombardia M, Benavides A, Silio V, Mäki M, Kaukinen K, Collin P, Thorell L, Ryöppy P, Mendez E. Quantification of wheat, barley and rye contamination in oat samples by real-time PCR. In: Stern M, ed. Proceedings of the 19<sup>th</sup> Meeting Working Group on Prolamin Analysis and Toxicity. Verl. Wissensch. Scripten. Zwickau 2005;87-94.
154. Peräaho M, Collin P, Kaukinen K, Mäki M. Contaminated oats and other gluten-free foods in the United States. Authors' response. *J Am Diet Assoc* 2005;105:348-9.
155. Ranua J, Luoma K, Auvinen A, Haapala AM, Mäki M, Peltola J, Raitanen J, Isojärvi JI. Antimitochondrial antibodies in patients with epilepsy. *Epilepsy Behav* 2005;7:95-7.
156. Ranua J, Luoma K, Auvinen A, Mäki M, Haapala AM, Peltola J, Raitanen J, Isojärvi J. Celiac disease-related antibodies in an epilepsy cohort and matched reference population. *Epilepsy Behav* 2005;6:388-92.
157. Viljamaa M, Collin P, Huhtala H, Sievänen H, Mäki M, Kaukinen K. Is coeliac disease screening in risk groups justified? A fourteen-year follow-up with special focus on compliance and quality of life. *Aliment Pharmacol Ther* 2005;22:317-24.
158. Collin P, Kaukinen K, Vogelsang H, Korponay-Szabó I, Sommer R, Schreier E, Volta U, Granito A, Veronesi L, Mascart F, Ocmant A, Ivarsson A, Lagerqvist C, Bürgin-Wolff A, Hadziselimovic F, Furlano R I, Sidler M A, Mulder CJJ, Goerres M S, Mearin ML, Ninaber MK, Gudmand-Høyer E, Fabiani E, Catassi C, Tidlund H, Alaintalo L, Mäki M. Antibodies and biopsy in the diagnosis of coeliac disease: Lessons from the Celikey European Multicentre Study. In: McMillan S, Feighery C, Watson P, O'Farrelly C, eds. Coeliac disease. Proceedings of the Eleventh International Symposium on Coeliac Disease, April 28-30, 2004, Belfast, Northern Ireland. Juvela, Semper. UK & Ireland, Nordic Countries 2006;11-4.
159. Hischenhuber C, Crevel R, Jarry B, Mäki M, Moneret-Vautrin DA, Romano A, Troncone R, Ward R. Review article: Safe amounts of gluten for patients with wheat allergy or coeliac disease. *Aliment Pharmacol Ther* 2006;23:559-75.
160. Holm K, Mäki M, Vuolteenaho N, Mustalahti K, Ashorn M, Ruuska T, Kaukinen K. Oats in the treatment of childhood coeliac disease: a 2-year controlled trial and a long-term clinical follow-up study. *Aliment Pharmacol Ther* 2006;23:1463-72.
161. Kaukinen K, Mäki M, Collin P. Immunohistochemical features in antiendomysium positive patients with normal villous architecture. Letter to the Editor. *Am J Gastroenterol* 2006;101:675-6.
162. Kaukinen K, Mäki M, Collin P. Intraepithelial lymphocytes in the diagnosis of coeliac disease. In: McMillan S, Feighery C, Watson P, O'Farrelly C, eds. Coeliac disease. Proceedings of the Eleventh International Symposium on Coeliac Disease, April 28-30, 2004, Belfast, Northern Ireland. Juvela, Semper. UK & Ireland, Nordic Countries 2006;22-5.
163. Kaukinen K, Mäki M, Kekkonen L. Uusi tutkimus etsii vastausta. Kannattaako keliakiaa seuloa väestöstä? *Keliakia* 2006;5:6.
164. Korponay-Szabó I, Raivio T, Nemes É, B. Kovács J, Laurila K, Kaukinen K, Mäki M. Coeliakia autoantitestek helyszíni kimutatása Biocard<sup>TM</sup> gyorstesztel (Detection of coeliac disease antibodies by the Biocard rapid test, in Hungarian). *Gyermekgyógyászat* 2006;57:351-357.
165. Kurppa K, Kaukinen K, Mäki M. Keliakian puhkeamista ja taudinkuvaa selvitetään. Keliakian perinnöllisyystutkimus menossa. *Keliakia* 2006;5:4-5.
166. Kurppa K, Kaukinen K, Mäki M. Uusi tutkimus alkamassa. Keliakian puhkeamiseen vaikuttavia perintötekijöitä etsitään. *Keliakia* 2006;1:8-9.
167. Mäki M. Gluteenitonta turvallisesti. Gluteenin mittaaminen elintarvikkeista. In: Mäki M et al, eds. *Keliakia*. Kustannus Oy Duodecim, Jyväskylä 2006;132-3.
168. Mäki M. Keliakia. In: Lehtola J et al., eds. Suomen Gastroenterologiayhdistys 50 vuotta. Suomen Gastroenterologiayhdistys ry, Tampere 2006:73-5.
169. Mäki M. Kuinka keliakia syntyy? Gluteeni. In: Mäki M et al, eds. *Keliakia*. Kustannus Oy Duodecim, Jyväskylä 2006;18-20.
170. Mäki M. Kuinka keliakia syntyy? Perimä. In: Mäki M et al, eds. *Keliakia*. Kustannus Oy Duodecim, Jyväskylä 2006;20-1.

171. Mäki M. Kuinka keliakia syntyy? Suolen limakalvolla tapahtuu. In: Mäki M et al, eds. Keliakia. Kustannus Oy Duodecim, Jyväskylä 2006;21-34.
172. Mäki M. Minne menet keliakia? In: Suomen Gastroenterologiayhdistys ry 50-vuotisjuhlakokous 7.-8.9.2006. Gastroenterologian saavutuksia ja tulevaisuuden näkymiä. Luennot. Turku 2006;12-14.
173. Mäki M. Minne menet keliakia? Keliakia huomenna. In: Mäki M et al, eds. Keliakia. Kustannus Oy Duodecim, Jyväskylä 2006;167-70.
174. Mäki M. Miten keliakia todetaan? Vasta-ainetestit diagnostiikassa. In: Mäki M et al, eds. Keliakia. Kustannus Oy Duodecim, Jyväskylä 2006;62-6.
175. Raivio T, Kaukinen K, Nemes E, Laurila K, Collin P, Kovacs JB, Mäki M, Korponay-Szabo IR. Self transglutaminase-based rapid coeliac disease antibody detection by a lateral flow method. *Aliment Pharmacol Ther* 2006;24:147-54.
176. Salmi T, Collin P, Korponay-Szabo IR, Laurila K, Partanen J, Huhtala H, Kiraly R, Lorand L, Reunala T, Mäki M, Kaukinen K. Endomysial antibody-negative coeliac disease: clinical characteristics and intestinal autoantibody deposits. *Gut* 2006;55:1746-53.
177. Salmi TT, Collin P, Järvinen O, Haimila K, Partanen J, Laurila K, Korponay-Szabo IR, Huhtala H, Reunala T, Mäki M, Kaukinen K. Immunoglobulin A autoantibodies against transglutaminase 2 in the small intestinal mucosa predict forthcoming coeliac disease. *Aliment Pharmacol Ther* 2006;24:541-52.
178. Collin P, Mäki M, Kaukinen K. Safe gluten threshold for patients with coeliac disease: some patients are more tolerant than others. Letter to the editor. *Am J Clin Nutr* 2007;86:260.
179. Collin P, Mäki M. Ohutsuoli. Keliakia. In: Höckerstedt K, Färkkilä M, Kivilaakso E, Pikkarainen P, eds. Gastroenterologia ja hepatologia. Duodecim, Jyväskylä 2007:378-88.
180. Kaukinen K, Collin P, Laurila K, Kaartinen T, Partanen J, Mäki M. Resurrection of gliadin antibodies in coeliac disease. Deamidated gliadin peptide antibody test provides additional diagnostic benefit. *Scand J Gastroenterol* 2007;42:1428-33.
181. Kaukinen K, Collin P, Mäki M. Latent coeliac disease or coeliac disease beyond villous atrophy? *Gut* 2007;56:1339-40.
182. Kaukinen K, Collin P, Mäki M. Long-term consumption of oats in coeliac disease. In: Stern M, ed. Proceedings of the 21st Meeting, Working Group on Prolamin Analysis and Toxicity. Verl. Wissensch. Scripten. Zwickau 2007:131-5.
183. Kaukinen K, Peräaho M, Lindfors K, Partanen J, Woolley N, Pikkarainen P, Karvonen A-L, Laasanen T, Sievänen H, Mäki M, Collin P. Persistent small bowel mucosal villous atrophy without symptoms in coeliac disease. *Aliment Pharmacol Ther* 2007;25:1237-45.
184. Kempainen T, Janatuinen E, Holm K, Kosma V-M, Heikkinen M, Mäki M, Laurila K, Uusitupa M, Julkunen R. No observed local immunological response at cell level after five years of oats in adult coeliac disease. *Scand J Gastroenterol* 2007;42:54-9.
185. Korponay-Szabó IR, Szabados K, Pusztai J, Uhrin K, Ludmány É, Nemes É, Kaukinen K, Kapitány A, Koskinen L, Sipka S, Imre A, Mäki M. Population screening for coeliac disease in primary care by district nurses using a rapid antibody test: diagnostic accuracy and feasibility study. *Brit Med J* 2007;335:1244-7.
186. Mäki M. Lääkehoito keliakiaan? *Keliakia* 2007;(4):6-7.
187. Raivio T, Korponay-Szabo I, Collin P, Laurila K, Huhtala H, Kaartinen T, Partanen J, Mäki M, Kaukinen K. Performance of a new rapid whole blood coeliac test in adult patients with low prevalence of endomysial antibodies. *Dig Liver Dis* 2007;39:1057-63.
188. Kaukinen K, Collin P, Mäki M. Natural history of coeliac disease. In: Fasano A, Troncone R, Branski D, eds. *Frontiers in Celiac Disease*. *Pediatr Adolesc Med*. Basel, Karger 2008;12:12-7.
189. Kaukinen K, Collin P, Mäki M. Vehnätärkkelyspohjaiset makeutusaineet ovat turvallisia keliakian hoidossa. *Keliakia* 2008;(4):6-7.
190. Kaukinen K, Mäki M. Self-detection of coeliac disease through a transglutaminase-based antibody: from laboratory to onsite testing. In: Stern M, ed. Proceedings of the 22nd Meeting, Working Group on Prolamin Analysis and Toxicity. Verlag Wissenschaftliche Scripten, Zwickau 2008:81-4.
191. Kaukinen K, Salmi T, Collin P, Huhtala H, Kärjä-Lahdensuu T, Mäki M. Clinical trial: gluten microchallenge with wheat-based starch hydrolysates in coeliac disease patients - a randomized, double-blind, placebo-controlled study to evaluate safety. *Aliment Pharmacol Ther* 2008;28:1240-8.
192. Koskinen O, Collin P, Korponay-Szabo I, Salmi T, Iltanen S, Haimila K, Partanen J, Mäki M, Kaukinen K. Gluten-dependent small bowel mucosal transglutaminase 2 -specific IgA deposits in overt and mild enteropathy coeliac disease. *J Pediatr Gastroenterol Nutr* 2008;47:436-42.
193. Kurppa K, Koskinen O, Collin P, Mäki M, Reunala T, Kaukinen K. Changing phenotype of coeliac disease after long-term gluten exposure. *J Pediatr Gastroenterol Nutr* 2008;47:500-3.
194. Kurppa K, Saavalainen P, Mäki M, Kaukinen K. Perinnöllisyystutkimuksessa on löydetty runsaasti keliakiaepäilyjä. *Keliakia* 2008;(4):8-9.
195. Lohi S, Mäki M, Montonen J, Knekt P, Pukkala E, Reunanen A, Kaukinen K. Tunnistamaton keliakia ei lisää yleistä syöpäriskiä. *Duodecim* 2008;124:2484.
196. Mäki M. Celiac disease. In: Kleinman RE, Goulet O\_J, Mieli-Vergani G, Sanderson IR, Sherman P, Shneider BL, eds. *Walker's Pediatric Gastrointestinal Disease*, 5th ed. BC Decker Inc, Hamilton 2008;319-27.

197. Raivio T, Korponay-Szabo IR, Paajanen T, Ashorn M, Iltanen S, Collin P, Laurila K, Nemes E, Kovacs JB, Carrard G, Saramäki M, Mäki M, Kaukinen K. Comparison of a novel whole blood transglutaminase-based ELISA with a whole blood rapid antibody test and established conventional serological coeliac disease assays. *J Pediatr Gastroenterol Nutr* 2008;47:562-7.
198. Troncone R, Ivarsson A, Szajewska H, Mearin ML, the European Multistakeholder Platform on CD (CDEUSSA), incl. Mäki M, Kaukinen K, Lähdeaho ML. Review article: future research on coeliac disease - a position report from the European multistakeholder platform on coeliac disease (CDEUSSA). *Aliment Pharmacol Ther* 2008;27:1030-43.
199. Collin P, Kaukinen K, Mäki M. Duodenal intraepithelial lymphocytosis: Celiac disease or not? Letter to the Editor. *Am J Gastroenterol* 2009;104:1847.
200. Kaukinen K, Collin P, Mäki M. Keliakian nykyhoito: ”vältä viljoja” ei riitä ohjeeksi. *S Lääkäril* 2009;64:277-83.
201. Koskinen O, Villanen M, Korponay-Szabo I, Lindfors K, Mäki M, Kaukinen K. Oats do not induce systemic or mucosal autoantibody response in children with coeliac disease. *J Pediatr Gastroenterol Nutr* 2009;48:559-65.
202. Kurppa K, Collin P, Viljamaa M, Haimila K, Saavalainen P, Partanen J, Laurila K, Huhtala H, Paasikivi K, Mäki M, Kaukinen K. Diagnosing mild enteropathy celiac disease: a randomized, controlled clinical study. *Gastroenterology* 2009;136:816-23. (doi:10.1053/j.gastro.2008.11.040).
203. Kurppa K, Collin P, Viljamaa M, Haimila K, Saavalainen P, Partanen J, Laurila K, Huhtala H, Paasikivi K, Mäki M, Kaukinen K. Keliakia tulee hoitaa riippumatta ohutsuolilöydöksestä. *Duodecim* 2009;125:307.
204. Peltola M, Kaukinen K, Dastidar P, Haimila K, Partanen J, Haapala AM, Mäki M, Keränen T, Peltola J. Hippocampal sclerosis in refractory temporal lobe epilepsy is associated with gluten sensitivity. *J Neurol Neurosurg Psychiatry* 2009;80:626-30 (doi:10.1136/jnnp.2008.148221).
205. Solakivi T, Kaukinen K, Kunnas T, Lehtimäki T, Mäki M, Nikkari ST. Serum fatty acid profile in celiac disease patients before and after a gluten-free diet. *Scand J Gastroenterol* 2009;44:826-30.
206. Lohi S, Lohi O, Vierula M, Mäki M, Toppari J. Coeliac disease autoantibodies in seminal plasma from cases with screen-detected coeliac disease. *Scand J Gastroenterol*, in press (Epub ahead of print).
207. Salmi TT, Collin P, Reunala T, Mäki M, Kaukinen K. Diagnostic methods beyond conventional histology in coeliac disease diagnosis. *Digest Liver Disease*, in press (doi:10.1016/j.dld.2009.04.004).
208. Koskinen O, Collin P, Lindfors K, Laurila K, Mäki M, Kaukinen K. Usefulness of small-bowel mucosal transglutaminase-2 specific autoantibody deposits in the diagnosis and follow-up of celiac disease. *J Clin Gastroenterol*, in press.

## Epidemiology

209. Mäki M, Holm K. Incidence and prevalence of coeliac disease in Tampere. Coeliac disease is not disappearing. *Acta Paediatr Scand* 1990;79:980-82.
210. Greco L, Mäki M, Di Donato F, Visakorpi JK. Epidemiology of coeliac disease in Europe and the mediterranean area. A summary report on the multicentre study by the European Society of Paediatric Gastroenterology and Nutrition. In: Auricchio S, Visakorpi JK, eds. *Common Food Intolerances 1: Epidemiology of Coeliac Disease*. Dyn Nutr Res, Basel, Karger 1992;2:25-44.
211. Mäki M, Holm K, Ascher H, Greco L. Factors affecting clinical presentation of coeliac disease: Role of type and amount of gluten-containing cereals in the diet. In: Auricchio S, Visakorpi JK, eds. *Common Food Intolerances 1: Epidemiology of Coeliac Disease*. Dyn Nutr Res, Basel, Karger 1992;2:76-82.
212. Uibo O, Uibo R, Kleimola V, Jögi T, Mäki M. Serum IgA anti-gliadin antibodies in an adult population sample: high prevalence without celiac disease. *Dig Dis Sci* 1993;38:2034-37.
213. Mäki M. Changing incidence of coeliac disease. *International Symposium on the Coeliac Condition*. Conference proceedings, Nutricia Dietary Products. The Governor's Hall, St. Thomas' Hospital, London, 27 October 1995; pp. 29-34.
214. Collin P, Reunala T, Rasmussen M, Kyrönpalo S, Pehkonen E, Laippala P, Mäki M. High incidence and prevalence of adult celiac disease: Augmented diagnostic approach. *Scand J Gastroenterol*. 1997;32:1129-33.
215. Mäki M. To screen or not to screen for coeliac disease. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;51-3.
216. Mäki M, Mustalahti K, Kokkonen J, Kulmala P, Haapalahti M, Karttunen T, Ilonen J, Laurila K, Dahlbom I, Hansson T, Höpfl P, Knip M. Prevalence of celiac disease among children in Finland. *N Engl J Med* 2003;348:2517-24.
217. Mäki M, Mustalahti K, Knip M. Screening for celiac disease. The author reply. *N Engl J Med* 2003;349:1673-4.
218. Lohi S, Mustalahti K, Kaukinen K, Laurila K, Collin P, Rissanen H, Lohi O, Bravi E, Gasparin M, Reunanen A, Mäki M. Increasing prevalence of coeliac disease over time. *Aliment Pharmacol Ther* 2007;26:1217-25.
219. Kondrashova A, Mustalahti K, Kaukinen K, Viskari H, Volodicheva V, Haapala AM, Ilonen J, Knip M, Mäki M, Hyöty H, and the EPIVIR Study Group. Lower economic status and inferior hygienic environment may protect against celiac disease. *Ann Med* 2008;40:223-31.

220. Vilppula A, Collin P, Mäki M, Valve R, Luostarinen M, Krekelä I, Patrikainen H, Kaukinen K, Luostarinen L. Undetected coeliac disease in the elderly. A biopsy-proven population-based study. *Digest Liver Dis* 2008;40:809-13.
221. Lohi S, Mäki M, Montonen J, Knekt P, Pukkala E, Reunanen A, Kaukinen K. Malignancies in cases with screening-identified evidence of coeliac disease: a long-term population-based cohort study. *Gut* 2009;58:643-7 (doi:10.1136/gut.2007.140970).
222. Vilppula A, Kaukinen K, Luostarinen L, Krekelä I, Patrikainen H, Valve R, Mäki M, Collin P. Increasing prevalence and high incidence of celiac disease in elderly people: A population-based study. *BMC Gastroenterology* 2009;9:49 (doi:10.1186/1471-230X-9-49).
223. Lohi S, Mäki M, Rissanen H, Knekt P, Reunanen A, Kaukinen K. Prognosis of unrecognized coeliac disease as regards mortality: A population-based cohort study. *Ann Med*, in press (2009,1-8,iFirst).

### **Genetics, immunology, pathogenesis and epithelial cell biology**

224. Rantala I, Mäki M, Laasonen A, Visakorpi JK. Periodate lysine-paraformaldehyde as fixative for the study of duodenal mucosa. Morphologic and immunohistochemical results at light and electron microscopic levels. *Acta Path Microb Immunol Scand Sect A* 1985; 93:165-73.
225. Koskimies S, Mäki M, Lipsanen V, Visakorpi J. HLA haplotypes in Finnish coeliac patients and their family members, possible protecting haplotypes. In: Kumar PJ, Walker-Smith JA, eds. *Coeliac Disease: One Hundred Years*. Univ. Printing Service, Univ. Leeds, Leeds, 1991:51-53.
226. Koskimies S, Lipsanen V, Mäki M, Visakorpi JK. HLA and coeliac disease, a Finnish family study. In: Mearin ML, Mulder CJJ, eds. *Coeliac Disease, 40 years gluten-free*. Kluwer Academic Publishers, Dordrecht, 1991:9-13.
227. Mäki M, Holm K, Collin P, Savilahti E. Increase in gamma/delta T cell receptor bearing lymphocytes in normal small bowel mucosa in latent coeliac disease. *Gut* 1991;32:1412-4.
228. Mäki M, Hällström O, Marttinen A. Reaction of human non-collagenous polypeptides with coeliac disease autoantibodies. *Lancet* 1991;338:724-25.
229. Holm K, Mäki M, Savilahti E, Lipsanen V, Laippala P, Koskimies S. Intraepithelial gamma/delta T-cell-receptor lymphocytes and genetic susceptibility to coeliac disease. *Lancet* 1992;339:1500-03.
230. Lähdeaho M-L, Mäki M, Parkkonen P, Kantonen T, Lehtinen M. Determination of antibodies to non-structural and structural adenovirus antigens by ELISA. *Acta Virol* 1992;36:524-32.
231. Savilahti E, Mäki M, Holm K, Koskimies S, Lipsanen V, Reunala T, Klemola T. Immunological markers of latent coeliac disease. In: Auricchio S, Visakorpi JK, eds. *Common Food Intolerances 1: Epidemiology of Coeliac Disease*. Dyn Nutr Res, Basel, Karger 1992;2:128-41.
232. Savilahti E, Reunala T, Mäki M. Increase of lymphocytes bearing the gamma/delta T-cell receptor in the jejunum of patients with dermatitis herpetiformis. *Gut* 1992;33:206-11.
233. Lähdeaho M-L, Lehtinen M, Rissa H-R, Hyöty H, Reunala T, Mäki M. Antipeptide antibodies to adenovirus E1b protein indicate enhanced risk of celiac disease and dermatitis herpetiformis. *Int Arch Allergy Immunol* 1993;101:272-76.
234. Lähdeaho M-L, Parkkonen P, Reunala T, Mäki M, Lehtinen M. Antibodies to E1b protein-derived peptides of enteric adenovirus type 40 are associated with celiac disease and dermatitis herpetiformis. *Clin Immunol Immunopathol* 1993;69:300-5.
235. Mäki M, Marttinen A, Koskimies S. Molecular pathogenesis of coeliac disease. Autoimmune mechanisms operative? In: *Frontiers in Molecular and Clinical Gastroenterology*, Farthing MJG, Grejs GJ, eds. Syllabus of European Association for Gastroenterology and Endoscopy, Postgraduate Course-1993, 1993;4:1-12.
236. Marttinen A, Mäki M. Purification of fibroblast-derived coeliac disease autoantigen molecules. *Pediatr Res* 1993;34:420-23.
237. Partanen J, Milner C, Campbell RD, Mäki M, Lipsanen V, Koskimies S. HLA-linked heat shock protein 70 (HSP70-2) gene polymorphism and coeliac disease. *Tissue Antigens* 1993;41:15-19.
238. Holm K, Savilahti E, Koskimies S, Lipsanen V, Mäki M. Immunohistochemical changes in the jejunum in first degree relatives of patients with coeliac disease and the coeliac disease marker DQ genes. HLA class II antigen expression, interleukin-2 receptor positive cells and dividing crypt cells. *Gut* 1994;35:55-60.
239. Mäki M, Marttinen A, Koskimies S. *Maladie coeliaque*. *Pathologie moléculaire et mécanismes auto-immuns*. *Méd Chir Dig* 1994;23:215-17.
240. Mäki M. Autoantibodies as markers of autoimmunity in coeliac disease pathogenesis. In: Feighery C, O'Farrelly C, eds. *Gastrointestinal immunology and gluten-sensitive disease*. Oak Tree Press, Dublin 1994:246-52.
241. Kerttula TO, Hällström O, Mäki M. Phenotypical characterization of peripheral blood T cells in patients with coeliac disease - elevation of antigen-primed CD45RO+ T lymphocytes. *Immunology* 1995;86:104-9.
242. Lähdeaho ML, Vainio E, Lehtinen M, Parkkonen P, Partanen J, Koskimies S, Mäki M. Activation of coeliac disease immune system by specific  $\alpha$ -gliadin peptides. *Cereal Chem* 1995;72:475-9.
243. Mäki M. The humoral immune system in coeliac disease. In: Howdle PD (ed.), *Coeliac disease*. Baillière's *Clinical Gastroenterology* 1995;9:231-49.

244. Halttunen T, Mäki M. Epiteelisolujen biologia suolen krypta-villus akselilla ja sen tutkiminen kolmiulotteisella soluviljelytekniikalla. *Solubiologi* 1996;14:51-57.
245. Halttunen T, Marttinen A, Rantala I, Kainulainen H, Mäki M. Fibroblasts and transforming growth factor  $\beta$  induce organization and differentiation of T84 human epithelial cells. *Gastroenterology* 1996;111:1252-62.
246. Jalava T, Mäki M, Marttinen A, Partanen J, Koskimies S. The in vitro response to human fibroblast-derived extracellular matrix proteins is restricted by specific HLA class II genes. Relevance for coeliac disease. *Human Immunol*, 1996;49:106-112.
247. Kerttula T, Collin P, Polvi A, Korpela M, Partanen J, Mäki M. Distinct immunological features of Finnish Sjögren's syndrome patients with the alleles HLA-DRB1\*0301, DQA1\*0501, and DQB1\*0201. Alterations in circulating T cell receptor  $\gamma\delta$  subsets. *Arthr Rheum* 1996;39:1733-9.
248. Mäki M, Rantala I. Jejunal epithelial cell stress protein expression in gluten-induced enteropathy (celiac disease). In: van Eden W and Young DB, eds. *Stress proteins in medicine*. Marcel Dekker, New York, 1996;pp. 399-405.
249. Mäki M. Coeliac disease and autoimmunity due to unmasking of cryptic epitopes? *Lancet* 1996;348:1046-47.
250. Polvi A, Eland C, Koskimies S, Mäki M, Partanen J. HLA DQ and DP in Finnish families with celiac disease. *Eur J Immunogenetics* 1996;23:221-34.
251. Mäki M. Autoantibodies and autoantigens in coeliac disease. In: Mäki M, Collin P, Visakorpi JK, eds. *Coeliac disease. Proceedings of the Seventh International Symposium on Coeliac Disease*. Coeliac Disease Study Group, University of Tampere. Tampere 1997;155-62.
252. Mäki M. Immunopathogenesis of coeliac disease. *Int J Immunopath Pharmacol* 1997;10:81-2.
253. Mäki M. Tissue transglutaminase as the autoantigen of coeliac disease. *Gut* 1997;41:565-6.
254. Polvi A, Garden OA, Elwood CM, Sørensen SM, Batt RM, Mäki M, Partanen J. Canine major histocompatibility complex genes DQA and DQB in Irish setter dogs. *Tissue Antigens* 1997;49:236-43.
255. Polvi A, Mäki M, Partanen J. Celiac patients predominantly inherit HLA-DPB1\*0101 positive haplotype from HLA-DQ2 homozygous parent. *Human Immunol* 1997;53:156-8.
256. Savilahti E, Örmälä T, Arato A, Hacsek G, Holm K, Klemola T, Nemeth A, Mäki M, Reunala T. Density of  $\gamma\delta^+$  T cells in the jejunal epithelium of patients with coeliac disease and dermatitis herpetiformis is increased with age. *Clin Exp Immunol* 1997;109:464-7.
257. Kerttula TO, Holm K, Partanen J, Polvi A, Mäki M. Circulating T lymphocyte subsets in coeliac disease (CoD) patients and healthy family members. *Clin Exp Immunol* 1998;111:536-40.
258. Mäki M. Autoimmunità, transglutaminasi tissutale e malattia celiaca. *Medico e Bambino* 1998;17:7.
259. Mäki M. Ohutsuoli. Immunologia. (Immunology, in Finnish). In: Höckerstedt K, Färkkilä M, Kivilaakso E, Pikkarainen P, eds. *Gastroenterologia. Duodecim, Jyväskylä* 1998;263-4.
260. Polvi A, Arranz E, Fernandez-Arquero M, Collin P, Mäki M, Sanz A, Calvo C, Maluenda C, Westman P, de la Concha EG, Partanen J. HLA-DQ2-negative celiac disease in Finland and Spain. *Human Immunol* 1998;59:169-75.
261. Polvi A, Garden OA, Houlston RS, Mäki M, Batt RM, Partanen J. Genetic susceptibility to gluten sensitive enteropathy in Irish setter dogs is not linked to the major histocompatibility complex. *Tissue Antigens* 1998;52:543-9.
262. Polvi A, Mäki M, Collin P, Partanen J. TNF microsatellite alleles  $\alpha 2$  and  $\beta 3$  are not primarily associated with celiac disease in the Finnish population. *Tissue Antigens* 1998;51:553-5.
263. Halttunen T, Mäki M. Serum immunoglobulin A from patients with celiac disease inhibits human T84 intestinal crypt epithelial cell differentiation. *Gastroenterology* 1999;116:566-72.
264. Holopainen P, Arvas M, Sistonen P, Mustalahti K, Collin P, Mäki M, Partanen J. CD28/CTLA4 gene region on chromosome 2q33 confers genetic susceptibility to celiac disease. A linkage and family-based association study. *Tissue Antigens* 1999;53:470-5.
265. Iltanen S, Holm K, Ashorn M, Ruuska T, Laippala P, Mäki M. Changing jejunal  $\gamma\delta$  T cell receptor (TCR)-bearing intraepithelial lymphocyte density in coeliac disease. *Clin Exp Immunol* 1999;117:51-5.
266. Iltanen S, Holm K, Partanen J, Laippala P, Mäki M. Increased density of jejunal  $\gamma\delta^+$  T cells in patients having normal mucosa - marker of operative autoimmune mechanisms? *Autoimmunity* 1999;29:179-87.
267. Iltanen S, Rantala I, Laippala P, Holm K, Partanen J, Mäki M. Expression of HSP-65 in jejunal epithelial cells in patients clinically suspected of coeliac disease. *Autoimmunity* 1999;31:125-32.
268. Kerttula TO, Collin P, Mäki M, Hurme M. Normal T-helper 1/T-helper 2 balance in peripheral blood of coeliac disease patients. *Scand J Immunol* 1999;49:197-202.
269. Mäki M. Autoantibodies and autoantigens in coeliac disease. In: *Eurodoctor . First Joint Scientific Meeting. Autoimmune diseases, recent developments & diagnostic tools*. Palais des Congres, Brussels. April 29-30, 1999. p. 35-7. European Medical Association, 1999.
270. Rantala I, Collin P, Holm K, Kainulainen H, Mustonen J, Mäki M. Small bowel T-cells, HLA-DR antigen, and GroEL stress protein in IgA nephropathy. *Kidney Int* 1999;55:2274-80.
271. Korponay-Szabo I, Sulkanen S, Halttunen T, Maurano F, Rossi M, Mazzarella G, Laurila K, Troncone R, Mäki M. Tissue transglutaminase is the target in both rodent and primate tissues for celiac disease-specific autoantibodies. *J Pediatr Gastroenterol Nutr* 2000;31:520-7.

272. Lindfors K, Halttunen T, Huotari P, Nupponen N, Vihinen M, Visakorpi T, Mäki M, Kainulainen H. Identification of novel transcription factor-like gene from human intestinal cells. EMBL Nucleotide Sequence Database accession number AJ249940. *Biochem Biophys Res Com* 2000;276:660-6.
273. Mäki M. Autoantibodies and autoantigens in coeliac disease. In: Auricchio S, Greco L, Maiuri L, Troncone R, eds. *Proceedings of the Eighth International Symposium on Coeliac Disease, April 21-24, 1999, Naples, Italy.* JGC Editions, Naples, 2000;57-63.
274. Mäki M. Ohutsuoli. Immunologia. (Immunology, in Finnish). In: Höckerstedt K, Färkkilä M, Kivilaakso E, Pikkarainen P, eds. *Gastroenterologia. Duodecim, Jyväskylä 2000*;264-5.
275. Bourgain C, Genin E, Holopainen P, Mustalahti K, Mäki M, Partanen J, Clerget-Darpoux F. Use of closely related affected individuals for the genetic study of complex diseases in founder populations. *Am J Hum Genet*, 2001;68:154-9.
276. Halttunen T, Mäki M. The potential role of disease specific IgA autoantibodies in coeliac disease pathogenesis. In: Mäki M, Tossavainen M, eds. *Proceedings of the Workshop on Transglutaminases, Protein Cross-Linking and Coeliac Disease, September 14-15, 2001, Tampere, Finland.* Coeliac Disease Study Group, University of Tampere. Tampere, 2001;149-56.
277. Holopainen P, Mustalahti K, Uimari P, Collin P, Mäki M, Partanen J. Candidate gene regions and genetic heterogeneity in gluten sensitivity. *Gut* 2001;48:696-701.
278. Korponay-Szabo I, Dahlbom I, Mäki M. Recognition of transglutaminase 2 by coeliac autoantibodies - importance of TG2 conformations and cofactors. In: Mäki M, Tossavainen M, eds. *Proceedings of the Workshop on Transglutaminases, Protein Cross-Linking and Coeliac Disease, September 14-15, 2001, Tampere, Finland.* Coeliac Disease Study Group, University of Tampere. Tampere, 2001;123-33.
279. Lindfors K, Halttunen T, Kainulainen H, Mäki M. Differentially expressed CC3/TIP30 and rab11 along in vivo and in vitro intestinal epithelial cell crypt-villus axis. *Life Sciences* 2001;69:1363-72.
280. Mäki M. Coeliac disease, gluten-induced disease entities, clinical aspects. In: Mäki M, Tossavainen M, eds. *Proceedings of the Workshop on Transglutaminases, Protein Cross-Linking and Coeliac Disease, September 14-15, 2001, Tampere, Finland.* Coeliac Disease Study Group, University of Tampere. Tampere, 2001;1-5.
281. Susi M, Holopainen P, Mustalahti K, Mäki M, Partanen J. Candidate gene region 15q26 and genetic susceptibility to coeliac disease in Finnish families. *Scand J Gastroenterol* 2001;36:372-4.
282. Kainulainen H, Rantala I, Collin P, Ruuska T, Päivärinne H, Halttunen T, Lindfors K, Kaukinen K, Mäki M. Blisters in the small intestinal mucosa of coeliac patients contain T cells positive for cyclooxygenase-2. *Gut* 2002;50:84-9.
283. Karell K, Holopainen P, Mustalahti K, Collin P, Mäki M, Partanen J. Not all HLA DR3 DQ2 haplotypes confer equal susceptibility to coeliac disease: transmission analysis in families. *Scand J Gastroenterol* 2002;37:56-61.
284. Karell K, Korponay-Szabo I, Szalai Zs, Holopainen P, Mustalahti K, Collin P, Mäki M, Partanen J. Genetic dissection between coeliac disease and dermatitis herpetiformis in sib pairs. *Ann Hum Genet* 2002;66:387-92.
285. Kaukinen K, Partanen J, Mäki M, Collin P. HLA-DQ typing in the diagnosis of celiac disease. *Am J Gastroenterol* 2002;97:695-9.
286. Liu J, Joo S-H, Holopainen P, Terwilliger J, Tong X, Grunn A, Brito M, Green P, Mustalahti K, Mäki M, Gilliam TC, Partanen J. Genomewide linkage analysis of celiac disease in Finnish families. *Am J Hum Genet* 2002;70:51-9.
287. Mustalahti K, Holopainen P, Karell K, Mäki M, Partanen J. Genetic dissection between silent and clinically diagnosed symptomatic forms of coeliac disease in multiplex families. *Digest Liver Dis* 2002;34:842-5.
288. Woolley N, Holopainen P, Bourgain C, Mustalahti K, Collin P, Mäki M, Partanen J. CD80 (B7-1) and CD86 (B7-2) genes and genetic susceptibility to coeliac disease. *Eur J Immunogenet* 2002;29:331-3.
289. Woolley N, Holopainen P, Ollikainen V, Mustalahti K, Mäki M, Kere J, Partanen J. A new locus for coeliac disease mapped to chromosome 15 in a population isolate. *Human Genetics* 2002;111:40-5.
290. Babron M-C, Nilsson S, Adamovic S, Torinsson Naluai Å, Wahlström J, Ascher H, Ciclitira PJ, Sollid LM, Partanen J, Greco L, Clerget-Darpoux F, and the European Genetics Cluster on Coeliac Disease (incl. Holopainen P, Karell K, Partanen J, Collin P, Mustalahti K, Mäki M). Meta and pooled analysis of European coeliac disease data. *Eur J Hum Genet* 2003;11:828-34.
291. Bolognesi E, Karell K, Percopo S, Coto I, Greco L, Mantovani V, Suoraniemi E, Partanen J, Mustalahti K, Mäki M, Momigliano-Richiardi P. Additional factor in some HLA DR3/DQ2 haplotypes confers a fourfold increased genetic risk of celiac disease. *Tissue Antigens* 2003;61:308-16.
292. Hausch F, Halttunen T, Mäki M, Khosla C. Design, synthesis and evaluation of gluten peptide analogs as selective inhibitors of human tissue transglutaminase. *Chem Biol* 2003;10:225-31.
293. Heinonen TYK, Pasternack L, Lindfors K, Breton C, Gastinel LN, Mäki M, Kainulainen H. A novel human glycosyltransferase: primary structure and characterisation of the gene and transcripts. *Biochem Biophys Res Com* 2003;309:166-74.
294. Iltanen S, Mäki M, Collin P, Kaukinen K, Partanen J, Hulkkonen J, Hurme M, Aittoniemi J. The association between mannan-binding lectin gene alleles and celiac disease. *Am J Gastroenterol* 2003;98:2808-9.
295. Karell K, Louka AS, Moodie SJ, Ascher H, Clot F, Greco L, Ciclitira PJ, Sollid LM, Partanen J and the members of the European Genetics Cluster on Coeliac Disease (incl. Collin P, Mustalahti K, Mäki M). HLA

- types in coeliac disease patients not carrying the DQA1\*05-DQB1\*02 (DQ2) heterodimer: Results from the European genetics cluster on coeliac disease. *Human Immunol* 2003;64:469-477.
296. Korponay-Szabo IR, Dahlbom I, Laurila K, Koskinen S, Woolley N, Partanen J, Kovacs JB, Mäki M, Hansson T. Elevation of IgG antibodies against tissue transglutaminase as a diagnostic tool for coeliac disease in selective IgA deficiency. *Gut* 2003;52:1567-71.
297. Korponay-Szabó IR, Laurila K, Szondy Z, Halttunen T, Szalai Z, Dahlbom I, Rantala I, Kovács JB, Fésüs L, Mäki M. Missing endomysial and reticulin binding of coeliac antibodies in transglutaminase 2 knockout tissues. *Gut* 2003;52:199-204.
298. Lindfors K, Viiri KM, Niittynen M, Heinonen TYK, Mäki M, Kainulainen H. TGF-beta induces the expression of SAP30L, a novel nuclear protein. *BMC Genomics* 2003;4:53.
299. Louka AS, Moodie SJ, Karell K, Bolognesi E, Ascher H, Greco L, Momigliano-Richiardi P, Partanen J, Ciclitira PJ, Sollid LM, and the European Genetics Cluster on Celiac Disease (incl. Holopainen P, Karell K, Partanen J, Collin P, Mustalahti K, Mäki M). A collaborative European search for non-DQA1\*05-DQB1\*02 celiac disease loci on HLA-DR3 haplotypes: analysis of transmission from homozygous parents. *Human Immunol* 2003;64:350-8.
300. Haimila K, Smedberg T, Mustalahti K, Mäki M, Partanen J, Holopainen P. Genetic association of coeliac disease susceptibility to polymorphisms in the ICOS gene on chromosome 2q33. *Genes Immun* 2004;5:85-92.
301. Holopainen P, Naluai ÁT, Moodie S, Percopo S, Coto I, Clot F, Ascher H, Sollid L, Ciclitira P, Greco L, Clerget-Darpoux F, Partanen J, and the members of the European Genetics Cluster on Coeliac Disease (incl. Collin P, Mustalahti K, Mäki M). Candidate gene region 2q33 in European families with coeliac disease. *Tissue Antigens* 2004;63:212-22.
302. Juuti-Uusitalo K, Mäki M, Kaukinen K, Collin P, Visakorpi T, Vihinen M, Kainulainen H. cDNA microarray analysis of gene expression in coeliac disease jejunal biopsy samples. *Journal of Autoimmunity* 2004;22:249-65.
303. Korponay-Szabo IR, Halttunen T, Szalai Z, Laurila K, Király R, Kovács JB, Fésüs L, Mäki M. *In vivo* targeting of intestinal and extraintestinal transglutaminase 2 by coeliac autoantibodies. *Gut* 2004;53:641-8.
304. Woolley N, Mustalahti K, Mäki M, Partanen J. Cytokine gene polymorphisms and genetic association with coeliac disease in the Finnish population. *Scand J Immunol* 2005;61:51-6.
305. Hadjivassiliou M, Mäki M, Sanders DS, Williamson C, Grünewald RA, Woodroffe N, Korponay-Szabó IR. Autoantibody targeting of brain and intestinal transglutaminase in gluten ataxia. *Neurology* 2006;66:373-7.
306. Heinonen TYK, Pelto-Huikko M, Pasternack L, Mäki M, Kainulainen H. Murine ortholog of the novel glycosyltransferase, B3GTL: primary structure, characterization of the gene and transcripts, and expression in tissues. *DNA Cell Biol* 2006;25:465-74.
307. Juuti-Uusitalo K, Kaukinen K, Mäki M, Kainulainen H. Gene expression in TGFbeta induced epithelial cell differentiation in a three-dimensional intestinal epithelial cell differentiation model. *BMC Genomics* 2006;7:279.
308. Metzger M-H, Heier M, Mäki M, Bravi E, Schneider A, Loewel H, Illig T, Schuppan D, Wichmann H-E. Mortality excess in individuals with elevated IgA anti-transglutaminase antibodies: The KORA/MONICA Augsburg cohort study 1989-1998. *Eur J Epidemiol* 2006;21:359-65.
309. Viiri KM, Korkeamäki H, Kukkonen MK, Nieminen LK, Lindfors K, Peterson P, Mäki M, Kainulainen H, Lohi O. SAP30L interacts with members of the Sin3A corepressor complex and targets Sin3A to the nucleolus. *Nucleic Acids Research* 2006;34:3288-98.
310. Garcia-Horsman JA, Venäläinen JI, Lohi O, Auriola IS, Korponay-Szabo IR, Kaukinen K, Mäki M, Männistö PT. Deficient activity of prolyl oligopeptidase on the immunoactive peptide digestion in coeliac disease. *Scand J Gastroenterol* 2007;42:562-71.
311. Juuti-Uusitalo K, Mäki M, Kainulainen H, Isola J, Kaukinen K. Gluten affects epithelial differentiation-associated genes in small intestinal mucosa of coeliac patients. *Clin Exp Immunol* 2007;150:294-305.
312. Ashorn S, Raukola H, Välineva T, Ashorn M, Wei B, Braun J, Rantala I, Kaukinen K, Luukkaala T, Collin P, Mäki M, Iltanen S. Elevated serum anti-Saccharomyces cerevisiae, anti-I2 and anti-OmpW antibody levels in patients with suspicion of celiac disease. *J Clin Immunol* 2008;28:486-94.
313. Haveri H, Westerholm-Ormio M, Lindfors K, Mäki M, Andersson LC, Heikinheimo M. Transcription factors GATA-4 and GATA-6 in normal and neoplastic human gastrointestinal mucosa. *BMC Gastroenterology* 2008;8:9.
314. Korkeamäki H, Viiri K, Kukkonen MK, Mäki M, Lohi O. Alternative mRNA splicing of SAP30L regulates its transcriptional repression activity. *FEBS Lett* 2008;582:379-84.
315. Korponay-Szabo IR, Vecsei Z, Kiraly R, Dahlbom I, Chirido R, Nemes E, Fesus L, Mäki M. Deamidated gliadin peptides form epitopes that transglutaminase antibodies recognize. *J Pediatr Gastroenterol Nutr* 2008;46:253-61.
316. Koskinen LLE, Korponay-Szabo IR, Viiri K, Juuti-Uusitalo K, Kaukinen K, Lindfors K, Mustalahti K, Kurppa K, Adany R, Pocsai Z, Szeles G, Einarsdottir E, Wijmenga C, Mäki M, Partanen J, Kere J, Saavalainen P. Myosin IXB gene region and gluten intolerance: linkage to coeliac disease and a putative dermatitis herpetiformis association. *J Med Genet* 2008;45:222-7.
317. Lindfors K, Blomqvist T, Juuti-Uusitalo K, Stenman S, Venäläinen J, Mäki M, Kaukinen K. Live probiotic *Bifidobacterium lactis* bacteria inhibit the toxic effects induced by wheat gliadin in epithelial cell culture. *Clin Exp Immunol* 2008;152:552-8.

318. Myrsky E, Kaukinen K, Syrjänen M, Korponay-Szabo IR, Mäki M, Lindfors K. Celiac disease-specific autoantibodies targeted against transglutaminase 2 disturb angiogenesis. *Clin Exp Immunol* 2008;152:111-9.
319. Oikarinen M, Tauriainen S, Honkanen T, Oikarinen S, Vuori K, Kaukinen K, Rantala I, Mäki M, Hyöty H. Detection of enteroviruses in the intestine of type 1 diabetic patients. *Clin Exp Immunol* 2008;151:71-5.
320. Stenman S, Lindfors K, Korponay-Szabo IR, Lohi O, Saavalainen P, Partanen J, Haimila K, Wieser H, Mäki M, Kaukinen K: Secretion of celiac disease autoantibodies after *in vitro* gliadin challenge is dependent on small-bowel mucosal transglutaminase 2-specific IgA deposits. *BMC Immunology* 2008;9:6..
321. Venäläinen J, Männistö PT, Kaukovirta-Norja A, Loponen J, Salovaara H, Stenman S, Lindfors K, Kaukinen K, Mäki M. Viljojen omista entsyymeistä apua keliakiaan? *Dosis* 2008;24:67-75.
322. Viiri LE, Viiri KM, Ilveskoski E, Huhtala H, Mäki M, Tienari PJ, Perola M, Lehtimäki T, Karhunen PJ. Interactions of functional apolipoprotein E gene promoter polymorphisms with smoking on aortic atherosclerosis. *Circ Cardiovasc Genet* 2008;1:107-16. (doi:10.1161/Circgenetics.108.791764).
323. Ashorn S, Välineva T, Kaukinen K, Ashorn M, Braun J, Raukola H, Rantala I, Collin P, Mäki M, Luukkaala T, Iltanen S. Serological responses to microbial antigens in celiac disease patients during a gluten-free diet. *J Clin Immunol* 2009;29:190-5.
324. Einarsdóttir E, Koskinen LLE, Dukes E, Kainu K, Suomela S, Lappalainen M, Zibera F, Korponay-Szabo IR, Kurppa K, Kaukinen K, Adany R, Pocsai Z, Széles G, Färkkilä M, Turunen U, Halme L, Paavola-Sakki P, Not T, Vatta S, Ventura A, Löfberg R, Torkvist P, Bresso F, Halfvarson J, Mäki M, Kontula K, Saarialho-Kere U, Kere J, D'Amato M, Saavalainen P. IL23R in the Swedish, Finnish, Hungarian and Italian populations: association with IBD and psoriasis, and linkage to coeliac disease. *BMC Medical Genetics* 2009;10:8. (doi:10.1186/1471-2350-10-8.)
325. Haimila K, Einarsdóttir E, de Kauwe A, Koskinen LLE, Pan-Hammarström Q, Kaartinen T, Kurppa K, Zibera F, Not T, Vatta S, Ventura A, Korponay-Szabo IR, Ádány R, Pocsai Z, Széles G, Dukes E, Kaukinen K, Mäki M, Koskinen S, Partanen J, Hammarström L, Saavalainen P. The shared CTLA4-ICOS risk locus in celiac disease, IgA deficiency and common variable immunodeficiency. *Genes Immun* 2009;10:151-61. (doi:10.1038/gene.2008.89).
326. Heinonen TYK, Mäki M. Peters' plus syndrome is a congenital disorder of glycosylation caused by a defect in the  $\beta$ 1,3-glucosyltransferase that modifies thrombospondin type 1 repeats. *Ann Med* 2009;41:2-10. (DOI: 10.1080/07853890802301975).
327. Juuti-Uusitalo K, Lindfors K, Mäki M, Patrikainen M, Isola J, Kaukinen K. Inhibition of epithelial growth factor receptor signalling does not preserve epithelial barrier function after *in vitro* gliadin insult. *Scand J Gastroenterol* 2009;44:820-5.
328. Koskinen L, Romanos J, Kaukinen K, Mustalahti K, Korponay-Szabo I, Barisani D, Bardella MT, Zibera F, Vatta S, Széles G, Pocsai Z, Karell K, Haimila K, Ádány R, Not T, Ventura A, Mäki M, Partanen J, Wijmenga C, Saavalainen P. Cost-effective HLA-typing with tagging SNPs predicts celiac disease risk haplotypes in the Finnish, Hungarian and Italian populations. *Immunogenetics* 2009;61:247-56 (DOI 10.1007/s00251-009-0361-3).
329. Koskinen LLE, Einarsdóttir E, Dukes E, Heap GAR, Dubois P, Korponay-Szabo IR, Kaukinen K, Kurppa K, Zibera F, Vatta S, Not T, Ventura A, Sistonen P, Ádány R, Pocsai Z, Széles G, Mäki M, Kere J, Wijmenga C, van Heel DA, Saavalainen P. Association study of the IL18RAP locus in three European populations with coeliac disease. *Human Molec Genet* 2009;18:1148-55.
330. Lindfors K, Kaukinen K, Mäki M. A role for anti-transglutaminase 2 autoantibodies in the pathogenesis of coeliac disease? *Amino Acids* 2009;36:685-91 (DOI 10.1007/s00726-008-0127-5).
331. Myrsky E, Syrjänen M, Korponay-Szabo IR, Mäki M, Kaukinen K, Lindfors K. Altered small-bowel mucosal vasculature network in untreated coeliac disease. *Scand J Gastroenterol* 2009;44:162-7.
332. Sareneva I, Koskinen LLE, Korponay-Szabo IR, Kaukinen K, Kurppa K, Zibera F, Vatta S, Not T, Ventura A, Ádány R, Pocsai Z, Széles G, Mäki M, Saavalainen P, Einarsdóttir E. Linkage and association study of FCγR polymorphisms in celiac disease. *Tissue Antigens* 2009;73:54-8.
333. Stenman SM, Venäläinen JI, Lindfors K, Auriola S, Mauriala T, Kaukovirta-Norja A, Jantunen A, Laurila K, Quiao S-W, Sollid LM, Männistö PT, Kaukinen K, Mäki M. Enzymatic detoxification of gluten by germinating wheat proteases: Implications for new treatment of celiac disease. *Ann Med* 2009;41:390-400.
334. Viiri KM, Jänis J, Siggers T, Heinonen TYK, Valjakka J, Bulyk ML, Mäki M, Lohi O. DNA-binding and -bending activities of SAP30L and SAP30 are mediated by a zinc-dependent module and monophosphoinositides. *Molec Cell Biol* 2009;29:342-56.
335. Viiri KM, Heinonen TYK, Mäki M, Lohi O. Phylogenetic analysis of the SAP30 family of transcriptional regulators reveals functional divergence in the domain that binds the nuclear matrix. *BMC Evol Biol* 2009;9:149. (doi:10.1186/1471-2148-9-149).
336. Koskinen LLE, Einarsdóttir E, Korponay-Szabo IR, Kurppa K, Kaukinen K, Sistonen P, Pocsai Z, Széles G, Ádány R, Mäki M, Kere J, Saavalainen P. Finemapping of the *CELIAC2* locus on chromosome 5q31-q33 in the Finnish and Hungarian populations. *Tissue Antigens*, in press.
337. Myrsky E, Caja S, Simon-Vecsei Z, Korponay-Szabo IR, Nadalutti C, Collighan R, Mongeot A, Griffin M, Mäki M, Kaukinen K, Lindfors K. Coeliac disease IgA modulates vascular permeability *in vitro* through the activity of transglutaminase 2 and RhoA. *Cell Mol Life Sci*, in press (DOI 10.1007/s00018-009-0116-1).

## Inflammatory bowel disease

338. Kangas E, Matikainen M, Auvinen O, Harju E, Inkovaara J, Mäki M. Clinical features of Crohn's disease: a clinical study of one hundred patients found in an unselected population. *Int Surg* 1986; 71:256-9.
339. Mäki M, Vaajalahti P, Arajärvi P. Upper gastrointestinal endoscopic findings in children with ulcerative colitis and Crohn's disease. *Scand J Gastroenterol* 1989; 24 (Suppl 158):93.
340. Ruuska T, Vaajalahti P, Arajärvi P, Mäki M. Prospective evaluation of upper gastrointestinal mucosal lesions in children with ulcerative colitis and Crohn's disease. *J Pediatr Gastroenterol Nutr* 1994;19:181-6.
341. Ruuska T, Savilahti E, Mäki M, Örmälä T, Visakorpi JK. Exclusive whole protein enteral diet versus prednisolone in the treatment of acute Crohn's disease in children. *J Pediatr Gastroenterol Nutr* 1994;19:175-80.

## Teeth and gastrointestinal diseases

342. Aine L, Mäki M, Collin P, Keyriläinen O. Dental enamel defects in coeliac disease. *J Oral Pathol Med* 1990;19:241-45.
343. Mäki M, Aine L, Lipsanen V, Koskimies S. Dental enamel defects in first-degree relatives of coeliac disease patients. *Lancet* 1991; 337:763-4.
344. Aine L, Reunala T, Mäki M. Dental enamel defects in children with dermatitis herpetiformis. *J Pediatr* 1991;118:572-4.
345. Aine L, Mäki M, Reunala T. Coeliac-type dental enamel defects in patients with dermatitis herpetiformis. *Acta Derm Venereol* 1992;72:25-27.
346. Aine L, Baer M, Mäki M. Dental erosions caused by gastroesophageal reflux disease in children. *J Dent Child* 1993;60:210-14.
347. Hölttä P, Aine L, Mäki M, Ruuska T, Vuento R, Ashorn M, Alaluusua S. Mutans streptococcal serotypes in children with gastroesophageal reflux disease. *J Dent Child* 1997;201-204.
348. Aine L, Backström MC, Mäki R, Kuusela A-L, Koivisto A-M, Ikonen RS, Mäki M. Enamel defects in primary and permanent teeth of children born prematurely. *J Oral Pathol Med* 2000;29:403-9.
349. Backström MC, Aine L, Mäki R, Kuusela AL, Sievänen H, Koivisto AM, Ikonen RS and Mäki M. Maturation of primary and permanent teeth in preterm infants. *Arch Dis Child Fetal Neonatal* 2000;83:F104-8.

## Helicobacter pylori, dyspepsia and gastroesophageal reflux

350. Baer M, Mäki M, Nurminen J, Turjanmaa V, Pukander J, Vesikari T. Esophagitis and findings of long-term esophageal pH recording in children with repeated lower respiratory tract symptoms. *J Pediatr Gastroenterol Nutr* 1986; 5:187-90.
351. Mäki M. Gastroesophageal reflux och organiska orsaker till recidiverande buksmärtor hos barn. In: Lindberg T, Stintzing G, eds. *Födoämnesreaktioner hos barn. Utbildningspärm, Bristol-Myers Nutrition* 1992;8:1-8:6.
352. Ashorn M, Mäki M, Ruuska T, Karikoski-Leo R, Hällström M, Kokki M, Miettinen A, Visakorpi JK. Upper gastrointestinal endoscopy in recurrent abdominal pain of childhood. *J Pediatr Gastroenterol Nutr* 1993;16:273-7.
353. Ashorn M, Ruuska T, Karikoski R, Välipakka J, Mäki M. Gastric mucosal cell densities in Helicobacter pylori positive and negative dyspeptic children and healthy controls. *J Pediatr Gastroenterol Nutr* 1994;18:146-51.
354. Ashorn M, Ruuska T, Karikoski R, Miettinen A, Mäki M. Helicobacter pylori gastritis in dyspeptic children. A long-term follow-up after treatment with colloidal bismuth subcitrate and tinidazole. *Scand J Gastroenterol* 1994;29:203-8.
355. Ashorn M, Karikoski-Leo R, Ruuska T, Miettinen A, Mäki M. Upper gastrointestinal endoscopy and the prevalence of Helicobacter pylori infection in recurrent abdominal pain of childhood. In: Gasbarrini G and Pretolani S (eds), *Basic and clinical aspects of Helicobacter pylori infection*, 1994, pp. 19-22. Springer-Verlag, Berlin.
356. Ashorn M, Mäki M, Hällström M, Uhari M, Åkerblom HK, Viikari J, Miettinen A. Helicobacter pylori infection in Finnish children and adolescents: A serologic cross-sectional and follow-up study. *Scand J Gastroenterol* 1995;30:876-9.
357. Ashorn M, Miettinen A, Ruuska T, Laippala P, Mäki M. Seroepidemiological study on Helicobacter pylori infection in infancy. *Arch Dis Child* 1996; 74:F141-2.
358. Mäki M: Gastroesophageal reflux hos spädbarn. In: Lindberg T, Stintzing G, eds. *Födoämnesreaktioner hos barn. Andra upplagan, Bristol-Myers Squibb*. 1998;11:1-5.
359. Mäki M: Recidiverande buksmärtor hos barn. In: Lindberg T, Stintzing G, eds. *Födoämnesreaktioner hos barn. Andra upplagan, Bristol-Myers Squibb*. 1998;12:1-2.
360. Oksanen K, Kainulainen H, Ruuska T, Mäki M, Ashorn M. Reverse transcription - polymerase chain reaction in the diagnosis of Helicobacter pylori infection in Finnish children. *J Pediatr Gastroenterol Nutr* 1999;28:252-6.

## Neonates in intensive care

361. Mäki M, Ruuska T, Kuusela A-L, Karikoski-Leo R, Ikonen RS. High prevalence of asymptomatic esophageal and gastric lesions in preterm infants in intensive care. *Critical Care Med* 1993;21:1863-67.

362. Kuusela AL, Ruuska T, Karikoski R, Laippala P, Ikonen RS, Janas M, Mäki M. A randomized, controlled study of prophylactic ranitidine in preventing stress-induced gastric mucosal lesions in neonatal intensive care unit patients. *Critical Care Med* 1997;25:346-51.
363. Sievänen H, Backström MC, Kuusela A-L, Ikonen RS, Mäki M. Dual energy X-ray absorptiometry of the forearm in preterm and term infants: evaluation of the methodology. *Pediatr Res* 1999;45:100-105.
364. Backström M, Mäki R, Kuusela A-L, Sievänen H, Koivisto A-M, Ikonen RS, Kouri T, Mäki M. Randomised controlled trial of vitamin D supplementation on bone density and biochemical indices in preterm infants. *Arch Dis Child Fetal Neonatal Ed* 1999;80:F161-6.
365. Backström MC, Mäki R, Kuusela A-L, Sievänen H, Koivisto A-M, Koskinen M, Ikonen RS, Mäki M. The long-term effect of early mineral, vitamin D and breast milk intake on bone mineral status in 9- to 11-year-old children born prematurely. *J Pediatr Gastroenterol Nutr* 1999;29:575-82.
366. Backström MC, Kouri T, Kuusela A-L, Sievänen H, Koivisto A-M, Ikonen RS, Mäki M. Bone isoenzyme of serum alkaline phosphatase and serum inorganic phosphate in metabolic bone disease of prematurity. *Acta Paediatr* 2000;89:867-73.
367. Kuusela A-L, Mäki M, Ruuska T, Laippala P. Stress-induced gastric findings in critically ill newborn infants: frequency and risk factors. *Intensive Care Med* 2000;26:1501-6.
368. Backström MC, Mahonen A, Ala-Houhala M, Sievänen H, Mäenpää P, Koivisto A M, Olkku A, Mäki R, Mäki M. Genetic determinants of bone mineral content in premature infants. *Arch Dis Child Fetal Neonatal Ed* 2001;85:F214-6.

## Miscellaneous

369. Mäki M, Sorto A, Hällström O, Visakorpi JK. Hepatic dysfunction and dysgammaglobulinaemia in Shwachman-Diamond syndrome. *Arch Dis Child* 1978; 53:693-4.
370. Hietanen E, Ståhlberg M-R, Mäki M. Development of intestinal drug-metabolizing enzymes in man. *Progress in Pharmacology and Clinical Pharmacology* 1989; 7/2:267-74.
371. Rantala I, Mäki M, Harmoinen A. Immunoelectron microscopy of duodenal epithelium in patients with milk-induced iron deficiency anaemia. In: *Inst Phys Conf Ser No 98*. IOP Publishing Ltd 1990:699-702
372. Lehtinen M, Lähdeaho ML, Parkkonen P, Mäki M, Paavonen J. The chlamydial groEL-like stress protein HypB contains both evolutionarily conserved and chlamydia specific epitopes. 2nd Scand Chlamydia Meeting 1991;1-4
373. Lehtinen M, Lähdeaho ML, Parkkonen P, Hyöty H, Mäki M, Paavonen J. Evolutionarily conserved and species-specific B- cell epitopes in chlamydial 60kD groEL-like stress protein HypB. *Proceedings of the European Society for Chlamydia Research and 2nd Int. Symp. of Uppsala University Centre for Std Research* 1992;87-91.
374. Paavonen J, Lähdeaho ML, Puolakkainen M, Mäki M, Parkkonen P, Lehtinen M. Antibody response to B cell epitopes of *Chlamydia trachomatis* 60-kDa heat-shock protein and corresponding mycobacterial and human peptides in infants with chlamydial pneumonitis. *J Infect Dis* 1994;169:908-11.
375. Savilahti E, Mäki M. Lasten toistuvat vatsakivut. *Duodecim* 1997;113:589-93.
376. Ashorn M, Mäki M, Ruuska T. Lasten toistuvat vatsavaivat. *Ferring Lääkkeet Oy: Espoo*; 1998:1-23.
377. Ruuska T, Ashorn M, Mäki M. Suoliston, maksan ja haiman sairaudet. In: *Raivio K, Siimes MA. Lastentaudit. Duodecim: Jyväskylä* 1998:343-53.
378. Korhonen R, Korpela R, Saxelin M, Mäki M, Kankaanranta H, Moilanen E. Induction of nitric oxide synthesis by probiotic *Lactobacillus rhamnosus* GG in J774 macrophages and human T84 intestinal epithelial cells. *Inflammation* 2001;25:223-32.
379. Ashorn P, Korpela P, Käär M-L, Lautala P, Viitanen T, Salo M, Csonka P, Eriksson K, Mäki M. Keskussairaaloiden ja yliopistosairaalan yhteistyö lastenlääkärien koulutuksessa. *S Lääkäril* 2004;59:2723-6.
380. Mäki M. Jarmo Visakorpi vuoden 2004 kunniapalkinnon saaja. *Suomalainen Tiedekatemia, vuosikirja* 2004. Vammala 2005;25-6.
381. Mäki M. Jarmo Visakorpi recipient of the Academy Award 2004. *Academia Scientiarum Fennica, Year book* 2004. Vammala 2005;27-8.
382. Korppi M, Kuusela A-L, Holm T, Lautala P, Tomminen A, Viitanen T, Mäki M. Lastentautien erikoislääkärinkoulutus keskussairaalassa. Koulutuksen auditoinnin toinen vaihe. *S Lääkäril* 2009;64:1467-71.

## Films

383. Mäki M. Small bowel biopsy in children. Rapid technique and handling of the biopsy specimen. Videofilm 17 min., Alfa-Video 1984, Tampere, Finland.
384. Mäki M. Gastro-oesophageal reflux disease in children (in Finnish). *Med-News, Terveystieteiden tutkimuskeskuksen ajankohtaiskasetti*. November 5/92.

## Patents

385. Mäki M, Korponay-Szabo I. USA States Patent Number 7,361,480 - USA, Methods and Means for Detecting Gluten-Induced Diseases, Patent Granted 22.4.2008.

386. Mäki M, Korponay-Szabo I. European Patent No. 1390753. Methods and Means for Detecting Gluten-Induced Diseases, European Patent Office 22.10.2008.
387. Viiri K, Mäki M, Lohi O. Patent Application no 20086014-Finland. 24.10.2008.

## Abstracts

1. Mäki M, Grönroos P, Vesikari T, Visakorpi JK. The invasiveness of *Yersinia enterocolitica* in acute diarrhoea. *Acta Paediatr Belg* 1978; 31:108-9.
2. Vesikari T, Mäki M, Grönroos P. *Yersinia enterocolitica* is invasive for human epithelial cells. XII International Congress of Microbiology, München 1978. Abstract I 21.
3. Mäki M, Vesikari T. Etiologisk utredning av diarré hos barn. (Aetiologic evaluation of childhood diarrhoea, in Swedish). XIX Nordiska Pediatriska Kongressen, Göteborg 1979. Abstract 37.
4. Vesikari T, Mäki M. Studies on pathogenic mechanisms in bacterial gastroenteritis. The Medical Research Council 1977-1979. The Academy of Finland, Helsinki 1980. Abstract 176.
5. Mäki M, Vesikari T, Visakorpi JK. Etiology of acute gastroenteritis in children. The Medical Research Council 1977-1979. The Academy of Finland, Helsinki 1980. Abstract 105.
6. Mäki M. Akuutti ripuli. (Acute diarrhoea, in Finnish). *Lääketiede* 80 *Medicin*, 6. - 11.1.80, Otaniemi. Luentolyhennelmät s. 59-60.
7. Janas M, Harmoinen A, Mäki R, Mäki M. Tissue damage as a cause of elevated C-reactive protein in newborns. 7th European Congress of Perinatal Medicine, September 2-5, 1980, Barcelona, Spain. Abstract I 4.
8. Mäki M, Vesikari T, Rantala I, Grönroos P. Childhood diarrhoea caused by *Yersinia enterocolitica*. XVI International Congress of Pediatrics, September 8-13, 1980, Barcelona, Spain. Abstract DH 2.
9. Vesikari T, Mäki M, Mäki R. Antibody response in human campylobacteriosis studied with enzyme linked immunosorbent assay (ELISA). Scottish-Scandinavian Conference on Infectious Diseases, St. Andrews, Sept. 1-5, 1980. Abstract 36.
10. Mäki M, Vesikari T, Sarkkinen HK. Clinical and laboratory features of rotavirus diarrhoea in young children. Scottish-Scandinavian Conference on Infectious Diseases, St. Andrews, Sept. 1-5, 1980. Abstract 37.
11. Visakorpi JK, Mäki M, Vesikari T. La *Yersinia enterocolitica* como causa de diarrea aguda. (*Yersinia enterocolitica* as the cause of acute diarrhoea, in Spanish). VI Reunion de la seccion de gastroenterologia infantil y nutricion de la asociacion Espanola de pediatria, Isla de La Toja, 19-21 Marzo, 1981. Abstract p. 23-4.
12. Mäki M. Akuutin ripulitaudin diagnoosi ja hoito. (Diagnosis and treatment of acute diarrhoea, in Finnish). *Lääketiede* 81 *Medicin*, 4.-9.1.1981, Otaniemi. Luentolyhennelmät s. 46-7.
13. Mäki M. *Yersinia*-infektiot. (Infections caused by *Yersinia*, in Finnish). Luennot, Tampereen Lääkäripäivät 1981. Abstrakti s. 97-8.
14. Vesikari T, Mäki M, Visakorpi JK. Plasmid-mediated adherence to epithelial cells and subsequent invasiveness as pathogenic mechanisms of diarrhoea and colitis in yersiniosis. *Ped Res* 1981; 15:1195.
15. Vesikari T, Mäki M, Sundqvist C, Nurmi T, Skurnik M. Plasmid associated pathogenic properties of *Yersinia enterocolitica*. International Symposium on Recent Advances in Enteric Infections, Brugge, Belgium, September 8-11, 1981. Abstract p. 33.
16. Turjanmaa V, Baer M, Arajärvi P, Mäki M, Ahola T, Karhumäki L, Uusitalo A. Gastro-oesophageal reflux (GER) during sleep detected by long-term monitoring of oesophageal pH (pH-LTM) in children with recurrent respiratory diseases (RRD). *Clin Phys* 1981; 1:6:613.
17. Mäki M. Lasten akuutit ripulitaudit. (Acute diarrhoea in children, in Finnish). *Lääketiede* 82 *Medicin*, 3. - 8.1.1982, Otaniemi. Luentolyhennelmät s. 182-3.
18. Mäki M, Vesikari T. Förändrad bild av spädbarnsdiarré i Finland. (Changing pattern of childhood diarrhoea in Finland, in Swedish). XX Nordiske Pediatriska Kongress, Oslo 2. - 5. juni 1982. Abstract 67.
19. Vesikari T, Mäki M, Rantala I, Visakorpi JK. Studies on the pathogenesis of *Yersinia* infection in guinea pigs - Serény test revisited. *Ped Res* 1982; 16:1042.
20. Mäki M. Lasten akuutti vatsa-pediatrin kannalta. (Paediatric aspects in acute abdominal crises, in Finnish). Luennot, Tampereen Lääkäripäivät 1983. Abstrakti s. 189-91.
21. Mäki M. Ulosteen mikrobiitutkimukset. (Microbial stool examinations, in Finnish). *Lääketiede* 83 *Medicin*, 9. - 14.1.1983, Otaniemi. Luentolyhennelmät s. 262-3.
22. Mäki M, Vesikari T, Isolauri E, Grönroos P, Visakorpi JK. *Clostridium difficile* in stools of young children with and without diarrhoea: Association with antibiotic usage. *Ped Res* 1983; 17:429.
23. Huupponen T, Mäki M, Hällström O, Reunala T, Vesikari T, Visakorpi JK. Reticulin antibodies as an indicator of coeliac disease in diabetic children. Scandinavian Society for the study of diabetes. 18th Annual Meeting, June 1-3, 1983, Kuopio, Finland. Abstract 30.
24. Hietanen E, Heinonen T, Ståhlberg M-R, Mäki M. Metabolism of xenobiotics in the small intestinal mucosa of children. Extrahepatic drug metabolism and chemical carcinogenesis. Univ. Stockholm and the Royal Swedish Academy of Sciences, IUB Symposium nr 125, Stockholm, Sweden, May 17-20, 1983. Abstract A 38.
25. Baer M, Pukander J, Arajärvi P, Mäki M, Turjanmaa V, Karhumäki L, Visakorpi JK. Gastroesophageal reflux (GER) and chronic bronchitis in childhood. IVth Congress of European Pediatric Respiratory Society, Cracow, Poland, September 22-24th, 1983. Abstract 15.
26. Isolauri E, Vesikari T, Mäki M. Clinical and laboratory features of *Yersinia*, *Campylobacter* and *Salmonella* enteritis in children. 8th Scottish-Scandinavian Conference on Infectious Diseases, Tampere, Finland, June 8-11, 1983. Abstract 49.

27. Parviainen T, Baer M, Ahola T, Karhumäki L, Mäki M, Turjanmaa V, Uusitalo A. Long-term monitoring of oesophageal pH (pH-LTM) during sleep in children with recurrent respiratory diseases (RRD) to detect gastro-oesophageal reflux (GER). *Clin Physiol* 1983; 3(suppl 2):54.
28. Mäki M. Lasten akuutti ripulitauti Suomessa. (Acute diarrhoea in children in Finland, in Finnish). *Lääketiede 84 Medicin*, 8.-13.1.1984, Otaniemi. Luentolyhennelmät s. 402-3.
29. Mäki M. Evaluation of a serum IgA class reticulín antibody test for the detection of childhood coeliac disease. 6. Symposium der Arbeitsgemeinschaft Gastroenterologie, Ernährung und Stoffwechsel der Gesellschaft für Pädiatrie der DDR. Oberhof 21.-23.5.1984, DDR. Abstract.
30. Baer M, Mäki M, Vesikari T, Turjanmaa V, Nurminen J, Visakorpi JK. Oesophagitis and gastro-oesophageal reflux during sleep in children with repeated lower respiratory tract symptoms. *Ped Res* 1984; 18:1052.
31. Mäki M, Hällström O, Vesikari T, Visakorpi JK. Serum IgA reticulín antibody detects childhood coeliac disease. *Ped Res* 1984; 18:1054.
32. Mäki M, Isolauri E, Vesikari T. Efficacy of cholestyramine at an early stage of acute diarrhoea in infants. *Ped Res* 1984; 18:1061.
33. Huupponen T, Lenko HL, Mäki M. Diabetes and coexisting chronic disease in children. *Diabetes in the Young. Bulletin of International Study Group of Diabetes in Children and Adolescents* 1984; 11:66. (The 6th International Beilinson Symposium on Future Trends in Juvenile Diabetes, Israel, October 1984).
34. Mäki M, Vaajalahti P, Arajärvi P, Visakorpi JK. Upper gastrointestinal endoscopic findings in childhood Crohn's disease. *Acta Paediatr Scand* 1985; Suppl 322:30.
35. Mäki M, Alfa-Video Tampere. Small bowel biopsy in children. Rapid technique and handling of the biopsy specimen. Teaching videotape, 17 minutes. *Acta Paediatr Scand Suppl* 1985; Suppl 322:33.
36. Mäki M, Vaajalahti P, Arajärvi P, Visakorpi JK. Upper gastrointestinal endoscopic findings in childhood Crohn's disease. The Third International Symposium on Infant Nutrition and Gastrointestinal Disease. August 20-25, 1985, Brussels, Belgium. Abstract 88.
37. Aine L, Mäki M, Aine E, Visakorpi JK. Enamel hypoplasia (EH) of permanent teeth in coeliac disease (CD). *Ped Res* 1986; 20:696.
38. Ståhlberg M-R, Hietanen E, Mäki M. Detoxication enzymes in small intestinal mucosa in children. *Ped Res* 1986; 20:697.
39. Mäki R, Ala-Houhala M, Koskinen M, Koskinen T, Janas M, Mäki M, Visakorpi JK. Bone mineralization of premature infants depends on calcium and phosphorus intake and not on the dose of vitamin D supplementation. *Ped Res* 1987; 22:110.
40. Mäki M, Hällström O, Verronen P, Visakorpi JK. Arthritis in childhood coeliac disease (CD). *Ped Res* 1987; 22:101.
41. Mäki M, Ståhlberg M-R, Hällström O, Viander M, Visakorpi JK. Predictability of small bowel mucosal damage by reticulín (ARA) and gliadin antibodies (AGA). *Ped Res* 1987; 22:103.
42. Mäki M, Hällström O, Visakorpi JK. Gluten challenge in postpubertal children with coeliac disease (CD): Detection of mucosal relapse with IgA-class reticulín antibody (ARA). *Ped Res* 1987; 22:103.
43. Verronen P, Mäki M, Vesikari T. Inflammatory gut changes in patients with JCA of recent onset. IIIrd international symposium on paediatric rheumatology, 16-19 October 1988, Prague, Czechoslovakia. Abstract p. 128.
44. Baer M, Mäki M. Barrett's esophagus and recurrent respiratory disease. First International Symposium on Gastroesophageal Reflux and Respiratory Disorders. Bruxelles 29 - 30 April 1988. Abstract.
45. Mäki M, Hällström O, Polanco I, Visakorpi JK. Comparison of reticulín (ARA) and endomysium (EMA) antibodies in coeliac disease. *Ped Res* 1988; 24:406.
46. Mäki M, Visakorpi JK. Normal small bowel histology does not exclude coeliac disease (CD). *Ped Res* 1988; 24:411.
47. Koskimies S, Mäki M, Lipsanen V, Visakorpi JK. HLA haplotypes in Finnish coeliac patients and their family members, possible protecting haplotypes. Samuel Gee 1888 - 1988 Coeliac Symposium, St. Bartholomew's Hospital, 4th - 6th September, 1988. Abstract T10.
48. Mäki M, Koskimies S, Visakorpi JK. Latent coeliac disease. Samuel Gee 1888 - 1988 Coeliac Symposium, St. Bartholomew's Hospital, 4th - 6th September, 1988. Abstract T31.
49. Mäki M. Milloin on syytä epäillä keliakiaa? (When to suspect coeliac disease, in Finnish). Tampereen Lääkäripäivät 1988, Luennot IX, s. 74.
50. Mäki M. Occurrence of coeliac disease in Finland. Tampereen Lääkäripäivät -89. Luennot X, p. 143.
51. Koskimies S, Lipsanen V, Mäki M, Visakorpi JK. Restriction fragment length polymorphism of HLA class II genes in coeliac patients, their family members and controls. *Ped Res* 1989;26:274.
52. Lähdeaho M-L, Lehtinen M, Mäki M, Parkkonen P, Rissa H-R, Hällström O, Visakorpi JK. Adenovirus antibodies in childhood coeliac disease, anti-gliadin antibodies in adenovirus diseased children - what is the chain of events? *Ped Res* 1989;26:274.
53. Mäki M, Ruuska T, Vaajalahti P. Prospective study of upper gastrointestinal endoscopic findings in children with ulcerative colitis and Crohn's disease. *Ped Res* 1990;27:534.
54. Mäki M, Koskimies S, Visakorpi JK. Latent coeliac disease. In: Kumar PJ, Walker-Smith JA, eds. *Coeliac Disease: One Hundred Years*. Univ. Printing Service, Univ. Leeds, Leeds, 1991:245.

55. Holm K, Mäki M, Hällström O, Koskimies S, Visakorpi JK. Healthy individuals with or without enteropathy but positive for serum reticulin antibodies (RA) are genetically similar to coeliac disease (CD) patients. *J Pediatr Gastroenterol Nutr* 1991;13:321.
56. Ruuska T, Kuusela A-L, Mäki M. Upper gastrointestinal endoscopy (UGE) of newborn infants in intensive care unit. *J Pediatr Gastroenterol Nutr* 1991;13:326.
57. Savilahti E, Reunala T, Mäki M. Increase of gamma/delta T-cell receptor bearing lymphocytes in the jejunum of patients with dermatitis herpetiformis. *J Pediatr Gastroenterol Nutr* 1991;13:323.
58. Mäki M, Hällström O, Marttinen A. Purification of human protein molecules - autoantigens - reacting with coeliac disease (CD) patient sera IgA. *J Pediatr Gastroenterol Nutr* 1991;13:321.
59. Lähdeaho M-L, Mäki M, Parkkonen P, Lehtinen M. Antibodies to a structural and a non-structural adenovirus antigen in adenovirus diseased children and children with coeliac disease. 12th Nordic Virus Symposium, June 13-15, 1991, Espoo, Finland. Abstract.
60. Hällström M, Miettinen A, Hyöty H, Ashorn M, Ruuska T, Mäki M. The age-dependent incidence of serum IgG antibodies to *Helicobacter pylori* in healthy Finnish children. *Ital J Gastroenterol* 1991;23(suppl.2):11.
61. Ashorn M, Ruuska T, Karikoski-Leo R, Hällström M, Mäki M. Upper gastrointestinal endoscopy and the prevalence of *Helicobacter pylori* infection in recurrent abdominal pain of childhood. *Ital J Gastroenterol* 1991;23(suppl.2):70-71.
62. Ascher H, Holm K, Kristiansson B, Mäki M. Different clinical and epidemiological features of childhood coeliac disease in two neighbouring countries, Sweden and Finland. *Epidemiology of Coeliac Disease*, Capri 11-12 October 1991. Abstract p. 34.
63. Mäki M, Savilahti E, Holm K, Lipsanen V, Koskimies S. Association between genetic markers and the density of intraepithelial (IEL) gamma/delta and alpha/beta T cells in healthy first-degree relatives of coeliac disease (CD) patients. Eleventh Annual Bristol Meyers Squibb/Mead Johnson Nutrition Research Symposium "Immunophysiology of the Gut", Boston, August 26-28, 1991, Abstract No. 5.
64. Ascher H, Kristiansson B, Holm K, Mäki M. Vad betyder glutenkonsumtionsmönstret för skillnaderna mella celiaki i Sverige och Finland? XLVIII Svenska Läkaresällskapets Riksstämman, Stockholm 27-29 november 1991. *Svenska Läkaresällskapets Handlingar Hygiea*, 1991;Band 100, Häfte 2:297.
65. Savilahti E, Mäki M, Holm K, Reunala T, Klemola T, Koskimies S. Intraepithelial gamma/delta T-cells receptor bearing cells as a marker of latent coeliac disease. Workshop on Epidemiology of Coeliac Disease. Capri, 11-12 October 1991. Abstract p. 28.
66. Mäki M. Screening tools to be used. Workshop on Epidemiology of Coeliac Disease. Capri, 11-12 October 1991. Abstract p. 26.
67. Mäki M. Effect of the type and amount of cereals in the diet. Workshop on Epidemiology of Coeliac Disease. Capri, 11-12 October 1991. Abstract p. 23.
68. Collin P, Korpela M, Keyriläinen O, Hällström O, Viander M, Mäki M. Rheumatic complaints as a presenting symptom in patients with coeliac disease. *Scand J Rheumatol* 1991;20:222.
69. Mäki M. Oksenteleva imeväinen: diagnostinen selvittely ja kotihoito. 13. Tampereen Lääkäripäivät -92. *Luennot XIII*, s. 232.
70. Ascher H, Holm K, Kristiansson B, Mäki M. Quality and quantity of gluten intake in infancy in Sweden and Finland: One factor of importance for the clinical expression of coeliac disease? *J Pediatr Gastroenterol Nutr* 1992;15:347.
71. Holm K, Mäki M, Savilahti E, Lipsanen V, Koskimies S, Visakorpi JK. Association between DR, DQA and DQB genes and the density of jejunal intraepithelial gamma/delta T cells. *J Pediatr Gastroenterol Nutr* 1992;15:352.
72. Holm K, Savilahti E, Mäki M, Koskimies S, Lipsanen V. HLA class II antigens in the intestine of first degree relatives of coeliac patients. *J Pediatr Gastroenterol Nutr* 1992;15:352.
73. Holm K, Savilahti E, Koskimies S, Lipsanen V, Mäki M. Coeliac disease family study: immunohistochemical changes in the jejunum of first-degree relatives. *International Coeliac Symposium, Dublin 22-25 July, 1992*. Abstract no. 36, p. 36.
74. Marttinen A, Mäki M. Human fibroblasts synthesise and secrete coeliac disease autoantigen molecules. *International Coeliac Symposium, Dublin 22-25 July, 1992*. Abstract no. 47, p. 39.
75. Reunala T, Mäki M. Gluten-free diet treatment in dermatitis herpetiformis: A 5-year study. *International Coeliac Symposium, Dublin 22-25 July, 1992*. Abstract no. 54, p. 41.
76. Ascher H, Holm K, Kristiansson B, Mäki M. Is gluten intake in infancy a factor of importance for the clinical expression of coeliac disease? *International Coeliac Symposium, Dublin 22-25 July, 1992*. Abstract no 6, p. 29.
77. Mäki M, Huupponen T, Holm K, Hällström O. Onset of diabetes initiates coeliac disease autoantibody production and development of enteropathy. *International Coeliac Symposium, Dublin 22-25 July, 1992*. Abstract no. 44, p. 38.
78. Mäki M. The role of antibody in coeliac disease pathogenesis. *International Coeliac Symposium, Dublin 22-25 July, 1992*. Abstract p. 24.
79. Reunala T, Aine L, Hällström O, Laurila K, Mäki M. Celiac-type dental enamel defects and circulating IgA-class antibodies in children with dermatitis herpetiformis. *Ped Dermatol* 1992;9:180.
80. Reunala T, Aine L, Mäki M. Coeliac-type dental enamel defects in patients with dermatitis herpetiformis. 18th World Congress of Dermatology, New York City, June 12-18, 1992. Abstract 189 A.

81. Ashorn M, Ruuska T, Välipakka J, Karikoski-Leo R, Mäki M. Gastric mucosal cell densities in children. *Irish J Med Sci* 1992;161(suppl.10):86.
82. Ashorn M, Ruuska T, Karikoski R, Välipakka J, Mäki M. Gastric inflammatory cell densities in children with dyspepsia. *Scand J Gastroenterol* 1993;28(suppl. 197):47.
83. Ascher H, Holm K, Kristiansson B, Mäki M. Different trends in epidemiology and clinical expression of childhood coeliac disease in Sweden and Finland. Is gluten intake in infancy a factor of importance? *Scand J Gastroenterol* 1993;28(suppl. 197):53.
84. Collin P, Helin H, Mäki M, Hällström O, Karvonen AL. Latent coeliac disease in patients positive in reticulin and gliadin antibody tests. *Scand J Gastroenterol* 1993;28(suppl. 197):53.
85. Marttinen A, Hakala S, Mäki M. Reticulin and endomysium antibodies recognize fibroblast-derived molecules. *Scand J Gastroenterol* 1993;28(suppl. 197):54.
86. Holm K, Rantala I, Hällström O, Ivanyi J, Mäki M. Gamma/delta T cell density and heat-shock protein expression in jejunal epithelium in children suspected for coeliac disease. *Scand J Gastroenterol* 1993;28(suppl. 197):54.
87. Ruuska T, Kuusela AL, Ikonen RS, Mäki M, Visakorpi JK. Four-day prophylactic H2-blocker treatment prevents preterm and term newborns from stress-induced gastric mucosal lesions. *J Pediatr Gastroenterol Nutr* 1993;17:460.
88. Hakala S, Mäki M, Marttinen A. Coeliac disease (CD) patient sera IgA recognizes human fibroblast-derived molecules. *J Pediatr Gastroenterol Nutr* 1993;17:462.
89. Lähdeaho ML, Vainio E, Mäki M, Lehtinen M. Identification of a coeliac disease specific B-cell epitope in A-gliadin. *J Pediatr Gastroenterol Nutr* 1993;17:462.
90. Ashorn M, Mäki M, Ruuska T, Karikoski L, Visakorpi JK. A morphometric study of gastric mucosa in *Helicobacter pylori* positive and negative subjects. *J Pediatr Gastroenterol Nutr* 1993;17:474.
91. Holm K, Rantala I, Hällström O, Ivanyi J, Mäki M. Enhanced expression of the human groEL stress-protein homologue (HSP65) in jejunal enterocytes in children with reticulin autoantibodies with or without mucosal atrophy. *J Pediatr Gastroenterol Nutr* 1993;17:471.
92. Ruuska T, Kuusela AL, Ikonen RS, Mäki M. Prophylactic ranitidine for prevention of gastric mucosal damage in preterms in intensive care. 7th Annual Meeting of the North American Society for Pediatric Gastroenterology and Nutrition. November 5-6, 1993, Chicago, USA. Abstract 75.
93. Mäki M, Marttinen A. Human fibroblast in vitro produced protein complex contains targets for reticulin and endomysium autoantibodies. 7th Annual Meeting of the North American Society for Pediatric Gastroenterology and Nutrition. November 5-6, 1993, Chicago, USA. Abstract 27.
94. Ashorn M, Miettinen A, Mäki M. Serologic findings for *Helicobacter pylori* in Finnish children in a 9 year follow-up study. *Acta Gastroenterologica Belgica* 1993;56(suppl.):59.
95. Mäki M, Huupponen T, Holm K, Hällström O. Serum reticulin autoantibody production and development of coeliac disease is triggered by onset of IDDM. In: Greene S (ed.) *Diabetes in the young*. XIXth Annual Meeting Aegean Sea, Greece, 1993, p. 68. Medicom Europe BV, The Netherlands.
96. Kuusela AL, Ruuska T, Janas M, Mäki M. Ranitidine treatment of stress-induced gastric lesions in newborns. *Ped Res* 1994;35:266.
97. Mäki M, Marttinen A. Fibroblast-derived human proteins are targets for genetically determined reticulin and endomysium autoantibodies. *Ped Res* 1994;35:266.
98. Ashorn M, Miettinen A, Mäki M. Serum immune response to *Helicobacter pylori* (Hp) in healthy Finnish children and in children with dyspeptic symptoms. *Mucosal Immunity and the gut epithelium. Interactions in health and disease*. Capri 22-23 April 1994, p. 44.
99. Mäki M. Keliakian epidemiologia ja patogeneesi. XV Tampereen Lääkäripäivät 17.-19.3.1994, Luennot XV, s. 31.
100. Mäki M. Antibodies in coeliac disease. *Mucosal Immunity and the gut epithelium. Interactions in health and disease*. Capri 22-23 April 1994, p. 25.
101. Holm K, Mäki M, Eland C, Partanen J, Kainulainen H, Koskimies S, Ruuska T, Rantala I. Jejunal mucosal heat-shock protein 65 expression in coeliac disease. *Mucosal Immunity and the gut epithelium. Interactions in health and disease*. Capri 22-23 April 1994, Abstract 43, p. 77.
102. Halttunen T, Marttinen A, Kainulainen H, Mäki M. Three-dimensional growth and fibroblast-supported differentiation of a gut epithelial cell line (T84). *Experimental Biology* 94, Anaheim, California, April 24-28, 1994. The FASEB Journal, Abstracts Part II 1994;8:A1007.
103. Kuusela A-L, Ruuska T, Lautamatti V, Mäki M, Ikonen RS. Towards optimal ranitidine dosage. Longterm gastric pH monitoring of newborns treated in intensive care (NICU). 14th European Congress of Perinatal Medicine, June 5-8, 1994, Helsinki, Finland. Abstract 563.
104. Kuusela A-L, Janas M, Ruuska T, Mäki M, Ikonen RS. The prevalence of symptomatic gastric haemorrhage in newborns in intensive care unit (NICU). 14th European Congress of Perinatal Medicine, June 5-8, 1994, Helsinki, Finland. Abstract 562.
105. Jalava T, Marttinen A, Mäki M, Koskimies S. Immune response in vitro to coeliac disease associated autoantigen, reticulin, is MHC restricted. 12th European Immunology Meeting, 14-17 June 1994, Barcelona, Spain. Abstract p. 341.

106. Rantala I, Holm K, Mustonen J, Collin P, Kainulainen H, Mäki M. Immunohistochemical quantitation of jejunal T cells, HLA-DR antigen, and human groEL stress-protein homologue in IgA nephropathy. *IgA Nephropathy International Symposium 28.2. - 1.3.1994, Adelaide, Australia. Abstract.*
107. Sulkanen S, Marttinen A, Mäki M. Human fibroblast-derived molecules as antigen in ELISA for celiac disease-specific IgA. *J Pediatr Gastroenterol Nutr* 1994;19:332.
108. Marttinen A, Jalava T, Koskimies S, Mäki M. HLA restricted immune response in vitro to human fibroblast-derived celiac disease-associated autoantigen. *J Pediatr Gastroenterol Nutr* 1994;19:363.
109. Kerttula TO, Hällström O, Mäki M. Analysis of peripheral blood T cells in patients with untreated coeliac disease (CoD) - elevation of CD45RP+ T cells. 12th European Immunology Meeting, 14-17 June 1994, Barcelona, Spain. Abstract p. 343.
110. Halttunen T, Marttinen A, Kainulainen H, Mäki M. Fibroblast-induced differentiation and three-dimensional organization of a gut epithelial cell line (T84). *FEBS Special Meetings, Biological Membranes, Helsinki, Finland, June 26 - July 1, 1994. Abstract 390.*
111. Ruuska T, Savilahti E, Mäki M, Örmälä T, Visakorpi JK. Whole protein enteral diet vs. prednisolone in the treatment of Crohn's disease. *J Pediatr Gastroenterol Nutr* 1994;19:356.
112. Sulkanen S, Marttinen A, Hällström O, Laurila K, Halttunen T, Mäki M. Human umbilical cord autoantibodies (HUC-ab). An ideal screening test for coeliac disease. *J Pediatr Gastroenterol Nutr* 1995;20:445.
113. Hacesek G, Savilahti E, Arato A, Reunala T, Mäki M, Nemeth A. Effect of age on jejunal intraepithelial lymphocyte subsets in patients with gluten sensitive enteropathy and in healthy controls. *J Pediatr Gastroenterol Nutr* 1995;20:474.
114. Polvi A, Koskimies S, Mäki M, Partanen J. A gene dosage effect of DQA1\*0501 and DQB1\*0201 alleles in human coeliac disease. *Scand J Immunol* 1995;41:636.
115. Halttunen T, Marttinen A, Mäki M. Fibroblast-induced differentiation of the epithelial cell line T84 is due to TGF- $\beta$ . *Clin Immunol Immunopathol* 1995;76:S3.
116. Koskimies S, Jalava T, Marttinen A, Partanen J, Mäki M. Immune response in vitro to coeliac disease associated auto-antigen is DR and DQ restricted. *Clin Immunol Immunopathol* 1995;76:S72.
117. Mäki M, Sulkanen S, Laurila K, Hällström O, Halttunen T, Marttinen A. Human umbilical cord autoantibodies (IgA-HUC-AB) are sensitive and specific for coeliac disease. *Clin Immunol Immunopathol* 1995;76:S73.
118. Marttinen A, Sulkanen S, Mäki M. ELISA for celiac disease-specific IgA with human fibroblast-synthesised molecules as the antigen. *Clin Immunol Immunopathol* 1995;76:S73.
119. Savilahti E, Hacesek G, Arato A, Reunala T, Mäki M, Nemeth A. Age and the density of intraepithelial gamma/delta T cell receptor positive cells in patients with coeliac disease. *Clin Immunol Immunopathol* 1995;76:S73.
120. Ashorn M, Mäki M, Ruuska T, Miettinen A. H. pylori infection in infancy. A seroepidemiological study. *Gut* 1995;37 (Suppl 1):A24.
121. Collin P, Nurmikko T, Himanen S-L, Erilä T, Mäki M. Brain atrophy and coeliac disease. International meeting on epilepsy and other neurological disorders in coeliac disease. San Marino 10-12 April 1995.
122. Ashorn M, Mäki M. A serological study on the incidence of H. pylori infection in infancy. *Digestive Disease Week, May 14-17, San Diego 1995; Abstract A-308.*
123. Collin P, Mäki M, Rasmussen M, Kyrönpalo S, Pehkonen E, Reunala T. The prevalence of adult coeliac disease in Finland. *Scand J Gastroenterol* 1995;30, Suppl 209:99. Presented at the Twenty-seventh Scandinavian congress of gastroenterology, Uppsala, Sweden; 29-31 May 1995.
124. Halttunen T, Mäki M. Purified coeliac disease patient serum IgA inhibits T84 epithelial cell differentiation in vitro. *J Pediatr Gastroenterol Nutr* 1996;22:435.
125. Halttunen T, Marttinen A, Rantala I, Kainulainen H, Mäki M. TGF- $\beta$  is responsible for the fibroblast-induced differentiation of the intestinal epithelial cell line T84. *J Pediatr Gastroenterol Nutr* 1996;22:436.
126. Kaukinen K, Collin P, Holm K, Karvonen A-L, Pikkarainen P, Mäki M. Markers of coeliac disease latency in patients with positive reticulin and gliadin antibodies. In: Collin P, Mäki M, eds. *All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 4.*
127. Iltanen S, Holm K, Eland C, Partanen J, Mäki M. Immunohistochemical changes, serological markers and HLA genes in patients with coeliac disease and in patients suspected for coeliac disease, but having normal jejunal mucosa. In: Collin P, Mäki M, eds. *All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 29.*
128. Kaukinen K, Collin P, Holm K, Mäki M, Salmi J. Coeliac disease and endocrinologic disorders. In: Collin P, Mäki M, eds. *All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 38.*
129. Jokinen J, Peters U, Mäki M, Miettinen A, Collin P. Coeliac disease in patients with oral mucosal symptoms. In: Collin P, Mäki M, eds. *All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 44.*
130. Lähteenoja H, Toivanen A, Irjala K, Viander M, Mäki M, Syrjänen S. Oral mucosal changes in patients with coeliac disease. In: Collin P, Mäki M, eds. *All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 44.*

131. Patinen P, Savilahti E, Hietanen J, Malmström M, Mäki M, Reunala T. Intraepithelial lymphocytes bearing gamma/delta receptor in the oral and jejunal mucosa of patients with dermatitis herpetiformis. In: Collin P, Mäki M, eds. All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 45.
132. Sulkanen S, Marttinen A, Laurila K, Halttunen T, Mäki M. Endomysium autoantibodies on human umbilical cord in coeliac disease. In: Collin P, Mäki M, eds. All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 54.
133. Sulkanen S, Halttunen T, Leivo EL, Laurila K, Mäki M. Human umbilical cord-derived fibroblasts as antigen in whole cell Elisa for coeliac disease antibodies. In: Collin P, Mäki M, eds. All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 54.
134. Reunala T, Collin P, Puolakka T, Vuolteenaho N, Miettinen A, Pikkarainen P, Mäki M. Oats for patients with dermatitis herpetiformis. In: Collin P, Mäki M, eds. All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 66.
135. Halttunen T, Mäki M. Purified coeliac disease patient serum IgA inhibits fibroblast and TGF- $\beta$  induced epithelial cell (T84) differentiation in vitro. In: Collin P, Mäki M, eds. All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 72.
136. Jalava T, Polvi A, Partanen J, Marttinen A, Mäki M, Koskimies S. Immune response in vitro to human fibroblast-derived coeliac disease autoantigen is HLA DR/DQ restricted. In: Collin P, Mäki M, eds. All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 88.
137. Kerttula TO, Holm K, Polvi A, Partanen J, Mäki M. Circulating CD45RO+ TcR $\gamma\delta$  cells in coeliac disease (CoD) patients and their healthy family members. In: Collin P, Mäki M, eds. All on Coeliac Disease. Free Paper Abstracts. Seventh International Symposium on Coeliac Disease. September 5-7, 1996, Tampere, Finland, p. 88.
138. Halttunen T, Mäki M. Fibroblasts induce differentiation of the crypt-like epithelial cell line T84 in 3-dimensional co-culture. *Mol Biol Cell* 1996;7:113a.
139. Iltanen S, Holm K, Eland C, Partanen J, Mäki M. Markers of coeliac disease latency in children having normal jejunal morphology. *J Pediatr Gastroenterol Nutr* 1997;24:453.
140. Sulkanen S, Halttunen T, Leivo E-L, Laurila K, Mäki M. Whole cell ELISA for coeliac disease antibodies using human umbilical cord-derived fibroblasts as antigen. *J Pediatr Gastroenterol Nutr* 1997;24:463.
141. Backström MC, Mäki R, Kuusela A-L, Sievänen H, Ikonen RS, Koivisto A-M, Mäki M. The effect of individual low vitamin D intake on bone mineral density in preterm infants. *J Pediatr Gastroenterol Nutr* 1997;24:493.
142. Iltanen S, Collin P, Korpela M, Holm K, Partanen J, Polvi A, Mäki M. Celiac disease and primary Sjögren's syndrome. American Gastroenterological Association, Washington, USA, May 12, 1997. Abstract 960.
143. Mäki M. Nutrition, small bowel mucosa, immunology and coeliac disease. *Nordiske barnelegedager*, Trondheim, June 11-14, 1997. Abstract I 23.
144. Halttunen T, Lindfors K, Mäki M. Coeliac disease patient serum IgA reduces T84 epithelial cell differentiation rate in vitro. FASEB Summer Research Conference 1997, Copper Mountain, Colorado, USA, August 3-8, 1997. Abstract 1:8.
145. Reunala T, Collin P, Mäki M. Gluten-free diet in dermatitis herpetiformis: With or without oats? 6th Congress of the European Academy of Dermatology and Venereology September 11-15, 1997, Dublin, Ireland. *J EADV* 1997;9;(Suppl.1):7.
146. Ruuska T, Ashorn M, Koivu H, Aine L, Karikoski R, Mäki M. Long-term follow-up of gastroesophageal reflux disease in children. *Gastroenterology* 1998;114:A272.
147. Halttunen T, Mäki M. Celiac disease autoantibodies and tissue transglutaminase antibodies interfere similarly fibroblast-induced epithelial cell (T84) differentiation. *Gastroenterology* 1998;114:A377.
148. Kaukinen K, Collin P, Holm K, Rantala I, Vuolteenaho N, Reunala T, Mäki M. Wheat starch containing gluten-free products in the treatment of gluten intolerance. *Gastroenterology* 1998;114:A384-5.
149. Sulkanen S, Halttunen T, Korponay-Szabo I, Laurila K, Collin P, Mäki M. Tissue transglutaminase autoantibody enzyme-linked immunosorbent assay in detecting celiac disease. *Gastroenterology* 1998;114:A420.
150. Lindfors K, Halttunen T, Kainulainen H, Mäki M. Transforming growth factor- $\beta$ 1 downregulates rab11 expression in differentiating T84 intestinal epithelial cells. *Gastroenterology* 1998;114:A892.
151. Holm K, Vuolteenaho N, Mäki M. No harm of oats in the diet of children with newly or previously diagnosed coeliac disease (CD). *J Pediatr Gastroenterol Nutr* 1998;26:549.
152. Korponay-Szabo IR, Sulkanen S, Halttunen T, Laurila K, Maurano F, Rossi M, Troncone R, Mäki M. Complete overlapping of tissue staining patterns of anti-transglutaminase antibodies and coeliac disease-specific IgA antibodies. *J Pediatr Gastroenterol Nutr* 1998;26:553.
153. Lindfors K, Halttunen T, Kainulainen H, Mäki M. rab11 expression is suppressed by TGF- $\beta$ 1 during differentiation of deep crypt T84 intestinal epithelial cells. *J Pediatr Gastroenterol Nutr* 1998;26:554.
154. Santtila S, Partanen J, Mäki M, Hurme M. Polymorphisms of the genes of the pro-inflammatory cytokines in coeliac disease: Association of the IL1A2 and TNF2 alleles. *J Pediatr Gastroenterol Nutr* 1998;26:564.
155. Sulkanen S, Laurila K, Kolho K-L, Sarnesto A, Halttunen T, Korponay-Szabo I, Savilahti E, Collin P, Mäki M. Immunological detection of tissue transglutaminase autoantibodies in coeliac disease. *J Pediatr Gastroenterol Nutr* 1998;26:566.

156. Lohiniemi S, Mäki M, Hallert C, Collin P. Quality of life in coeliac disease patients treated with gluten-free diet for 10 years. *Scand J Gastroenterol* 1998;33 Suppl 227:F2
157. Holm K, Vuolteenaho N, Mäki M. No harm of oats in the diet of children with newly or previously diagnosed coeliac disease (CD). In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;116 (A3).
158. Kaukinen K, Collin P, Holm K, Rantala I, Vuolteenaho N, Reunala T, Mäki M. Wheat starch containing gluten-free products are well tolerated in the treatment of coeliac disease. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;117 (A5).
159. Mustalahti K, Sulkanen S, Holopainen P, Laurila K, Collin P, Partanen J, Mäki M. Undetected coeliac disease in multiple case families. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;125 (A22).
160. Lohiniemi S, Mustalahti K, Collin P, Vuolteenaho N, Rostila I, Mäki M. Quality of life in Finnish coeliac disease patients. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;139 (A50)
161. Iltanen S, Holm K, Ashorn M, Ruuska T, Mäki M. The density of jejunal intraepithelial  $\gamma\delta$ + T cells increases during development of coeliac disease and decreases on gluten-free diet. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;142 (A55).
162. Kaukinen K, Salmi J, Lahtela J, Oksa H, Siljamäki-Ojansuu U, Mäki M, Collin P. Gluten-free diet does not change the metabolic control of diabetes in patients suffering from diabetes mellitus and coeliac disease. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;149 (A70).
163. Korponay-Szabo IR, Sulkanen S, Halttunen T, Laurila K, Mäki M. Coeliac disease-specific autoantibodies recognize epitopes involving the transamidase active site of the calcium-activated tissue transglutaminase. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;157 (A86).
164. Korponay-Szabo IR, Laurila K, Szalai Zs, Török E, Sulkanen S, Kovacs J, Mäki M. Comparison of transglutaminase ELISA and endomysial antibody results in children with dermatitis herpetiformis. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;158 (A87).
165. Sulkanen S, Halttunen T, Laurila K, Kolho K-L, Korponay-Szabo IR, Sarnesto A, Savilahti E, Collin P, Mäki M. Tissue transglutaminase as antigen in enzyme-linked immunosorbent assay for coeliac disease autoantibodies. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;159 (A89)
166. Polvi A, Arranz E, Fernandez-Arquero M, Collin P, Mäki M, Sanz A, Calvo C, Maluenda C, Westman P, de la Concha EG, Partanen J. HLA-DQ2-negative celiac disease in Finland and Spain. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;161 (A93).
167. Arvas M, Mustalahti K, Holopainen P, Sistonen P, Polvi A, Collin P, Mäki M, Partanen J. Evaluation of a major non-MHC candidate susceptibility locus (6p23) for celiac disease in the Finnish population. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;161 (A94).
168. Holopainen P, Mustalahti K, Collin P, Mäki M, Partanen J. HLA DQ in families with both full and clinically silent (ARA/EMA and biopsy proven) celiac disease. In: Lohiniemi S, Collin P, Mäki M, eds. *Changing Features of Coeliac Disease*. The Finnish Coeliac Society. Tampere 1998;162 (A95).
169. Backström MC, Mäki M, Kuusela A-M, Sievänen H, Koivisto A-M, Koskinen M, Ikonen RS, Mäki M. The long-term effect of early mineral, Vitamin D and breast milk intake on bone mineral status in 9-11 years old children born prematurely. *J Pediatr Gastroenterol Nutr* 1999;28:586.
170. Halttunen T, Mäki M. Purified coeliac disease patient serum IgA interferes with fibroblast - extracellular matrix interactions. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.47.
171. Holopainen P, Arvas M, Sistonen P, Mustalahti K, Collin P, Mäki M, Partanen J. CD28/CTLA4 gene region on chromosome 2q33 confers genetic susceptibility to celiac disease. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.53.
172. Holopainen P, Arvas M, Sistonen P, Mustalahti K, Collin P, Mäki M, Partanen J. CD28/CTLA4 gene region on chromosome 2q33 confers genetic susceptibility to celiac disease. *J Pediatr Gastroenterol Nutr* 1999;28:558.
173. Holopainen P, Arvas M, Sistonen P, Mustalahti K, Polvi A, Collin P, Mäki M, Partanen J. Linkage and family-based association study of candidate regions 11qTER, 5qTER, 6p23 and 6p21.3 in celiac disease in Finland. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.10.
174. Holopainen P, Ollikainen V, Klinger N, Mustalahti K, Mäki M, Kere J, Partanen J. Evaluation of celiac disease risk markers in a founder population. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.56.
175. Holopainen P, Ollikainen V, Klinger N, Mustalahti K, Mäki M, Kere J, Partanen J. Evaluation of celiac disease risk markers in a founder population. *J Pediatr Gastroenterol Nutr* 1999;28:557.

176. Iltanen S, Collin P, Korpela M, Holm K, Partanen J, Polvi A, Mäki M. Association between coeliac disease and Sjögren's syndrome. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.105
177. Iltanen S, Holm K, Ashorn M, Ruuska T, Mäki M. Jejunal intraepithelial  $\gamma\delta$ + T cell density increases during coeliac disease development and decreases on gluten-free diet. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.66.
178. Kainulainen H, Rantala I, Halttunen T, Lindfors K, Collin P, Mäki M. Cyclooxygenase-2 (COX-2) is expressed in mucosal lesions of untreated coeliac patients. *Immunology Letters* 1999;69:101.
179. Kaukinen K, Mäki M, Turjanmaa K, Reunala T, Karvonen A-L, Partanen J, Collin P. Abdominal complaints after ingestion of cereals. Coeliac disease or food allergy? Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.106.
180. Korponay-Szabo IR, Halttunen T, Laurila K, Szalai Zs., Török E, Sulkanen S, Szabo T, Kovacs J-B, Mäki M. Transglutaminase ELISA with human red blood cells. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.201.
181. Korponay-Szabo IR, Halttunen T, Sulkanen S, Szalai Zs, Laurila K, Török E, Kovacs JB, Mäki M. Coeliac disease-specific and coeliac disease non-specific transglutaminase autoantibodies. 7<sup>th</sup> UEGW Oral presentations. *Gut* 1999;45 (Suppl V): A27.
182. Korponay-Szabo IR, Sulkanen S, Halttunen T, Maurano F, Rossi M, Laurila K, Troncone R, Mäki M. Tissue transglutaminase is the target in rodent and primates tissues for reticulin/endomysial antibodies. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.73.
183. Lähteenoja H, Toivanen A, Viander M, Rähä I, Vilja P, Mäki M, Syrjänen S. Local challenge on oral mucosa with gliadin and a gliadin derived peptide: is it helpful as second challenge method in CD? Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.286.
184. Lähteenoja H, Viander M, Rähä I, Toivanen A, Syrjänen S, Mäki M. T cell subsets of oral mucosa in patients with coeliac disease: increase of T cell subpopulations despite treatment. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.285.
185. Lindfors K, Halttunen T, Kainulainen H, Mäki M. The expression of rab11 and TIP30/CC3 is modulated during intestinal epithelial cell differentiation. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.122.
186. Lohiniemi S, Mustalahti K, Collin P, Vuolteenaho N, Rostila I, Mäki M. Quality of life in Finnish coeliac disease patients - a national cross sectional survey. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.266.
187. Mäki M. Antibody testing: new diagnostic strategies. Scientific Conference on coeliac disease. Frankfurt/Main, 1999. Abstracts.
188. Mäki M. Autoantibodies and autoantigens in coeliac disease. 7<sup>th</sup> UEGW Oral presentations. *Gut* 1999;45 (Suppl V): A27.
189. Mäki M. Autoantibodies and autoantigens in coeliac disease. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.31.
190. Mäki M. Autoantibodies, autoantigens and coeliac disease. *It J Anat Embr* 1999;104 (Suppl 1):422.
191. Mustalahti K, Collin P, Sievänen H, Salmi J, Mäki M. Bone mineral density in clinically silent celiac disease. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.256.
192. Mustalahti K, Lohiniemi S, Collin P, Mäki M. Effect of gluten-free diet on quality of life of silent celiac disease (CD) patients. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.269.
193. Mustalahti K, Sulkanen S, Holopainen P, Laurila K, Collin P, Partanen J, Mäki M. Undetected coeliac disease in Finnish multiple case families. Eighth International Symposium on Coeliac Disease, Castel Sant'Elmo, Naples, Italy, April 21-24, 1999. Abstract book, p.133.
194. Mustalahti K, Sulkanen S, Holopainen P, Laurila K, Collin P, Partanen J, Mäki M. Strategy of identifying undetected coeliac disease in Finnish multiple case families. *J Pediatr Gastroenterol Nutr* 1999;28:562.
195. Halttunen T, Mäki M. Disrupted fibroblast - extracellular matrix interaction in celiac disease? World Congress of Pediatric Gastroenterology, Hepatology, and Nutrition. Abstracts. Boston, Massachusetts, USA, August 5-9, 2000. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl 2):S59.
196. Halttunen T, Mäki M. Disrupted fibroblast extracellular matrix interaction in celiac disease? Ninth International Symposium on Coeliac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S20.
197. Holm K, Vuolteenaho N, Mäki M. Two years follow up of children with celiac disease (CD) using oats in their diet. World Congress of Pediatric Gastroenterology, Hepatology, and Nutrition. Abstracts. Boston, Massachusetts, USA, August 5-9, 2000. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl 2):S284.
198. Holm K, Vuolteenaho N, Mäki M. Two years follow up of children with celiac disease (CD) using oats in their diet. Ninth International Symposium on Coeliac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S23.

199. Holopainen P, Ollikainen V, Mustalahti K, Collin P, Mäki M, Partanen J. CTLA4/CD28 gene region in Finnish celiac disease families. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S13.
200. Iltanen S and The Coeliac Disease Study Group of Tampere. Markers of latent coeliac disease. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S23.
201. Iltanen S, Rantala I, Laippala P, Holm K, Partanen J, Mäki M. Enhanced HSP-65 expression as a marker of coeliac disease. World Congress of Pediatric Gastroenterology, Hepatology, and Nutrition. Abstracts. Boston, Massachusetts, USA, August 5-9, 2000. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl 2):S65.
202. Kainulainen H, Rantala I, Collin P, Ruuska T, Päivärinne H, Halttunen T, Lindfors K, Kaukinen K, Mäki M. COX-2 positive T-cells are present in blisters of small bowel mucosa of celiac patients. World Congress of Pediatric Gastroenterology, Hepatology, and Nutrition. Abstracts. Boston, Massachusetts, USA, August 5-9, 2000. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl 2):S9-10.
203. Kainulainen H, Rantala I, Collin P, Ruuska T, Päivärinne H, Halttunen T, Lindfors K, Kaukinen K, Mäki M. Cyclooxygenase-2 and celiac disease. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S8.
204. Karell K, Holopainen P, Mustalahti K, Collin P, Mäki M, Partanen J. Not all HLA DQ2 haplotypes confer equal susceptibility to coeliac disease: a transmission analysis in families. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S18.
205. Kaukinen K, Mäki M, Peräaho M, Collin P. A changing pattern of refractory sprue. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S20.
206. Kaukinen K, Peräaho M, Paasikivi K, Sievänen H, Mäki M, Collin P. Wheat starch-containing gluten-free products in the treatment of newly-detected coeliac disease - prospective and randomised study. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S20.
207. Kokkonen J, Haapalahti M, Laurila K, Karttunen TJ, Mäki M. Food allergy at school age - blind challenge and immuno-histochemical findings obtained by endoscopy. World Congress of Pediatric Gastroenterology, Hepatology, and Nutrition. Abstracts. Boston, Massachusetts, USA, August 5-9, 2000. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl 2):S61.
208. Korponay-Szabo I, Halttunen T, Laurila K, Szalai Z, B.Kovacs J, Collin P, Mäki M. Threshold of human transglutaminase autoantibody positivity in known coeliacs. 8th United European Gastroenterology Week, Brussels, 2000. *Gut* 2000;47, Suppl III, 198.
209. Korponay-Szabo IR, Halttunen T, Laurila K, Sulkanen S, Lörincz M, B-Kovacs J, Mäki M. Transglutaminase ELISA for IgA-deficient subjects. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S9.
210. Korponay-Szabo IR, Halttunen T, Laurila K, Sulkanen S, Lörincz M, B.Kovacs J, Mäki M. Detection of non-IgA class tissue transglutaminase autoantibodies in IgA deficient celiac patients. 6th International Symposium on Transglutaminases and Protein Cross-Linking, Lyon, France, September 16-19, 2000.
211. Korponay-Szabo IR, Halttunen T, Laurila K, Sulkanen S, Szalai Z, Dahlbom I, Kovacs JB, Mäki M. Calreticulin-bound tissue transglutaminase is less antigenic for celiac autoantibodies. World Congress of Pediatric Gastroenterology, Hepatology, and Nutrition. Abstracts. Boston, Massachusetts, USA, August 5-9, 2000. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl 2):S10.
212. Korponay-Szabo IR, Halttunen T, Laurila K, Sulkanen S, Szalai Zs, Dahlbom I, B-Kovacs J, Mäki M. Calreticulin-bound tissue transglutaminase is less antigenic for celiac autoantibodies. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S9.
213. Lähteenoja H, Viander M, Vilja P, Syrjänen S, Mäki M. Oral mucosa responds by recruitment and non-proliferative activation of T cells after submucosal challenge with a gliadin derived peptide. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S7.
214. Lindfors K, Halttunen T, Huotari P, Nupponen N, Vihinen M, Visakorpi T, Mäki M, Kainulainen H. Identification of DED, an evolutionary conserved novel putative transcription factor. World Congress of Pediatric Gastroenterology, Hepatology, and Nutrition. Abstracts. Boston, Massachusetts, USA, August 5-9, 2000. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl 2):S280-1.
215. Lindfors K, Halttunen T, Huotari P, Nupponen N, Vihinen M, Visakorpi T, Mäki M. Identification of DED, an evolutionary conserved novel putative transcription factor. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S22.
216. Lindfors K, Halttunen T, Kainulainen H, Mäki M. CC3/TIP30 and rab11 - novel markers for in vitro and in vivo intestinal epithelial cell differentiation. World Congress of Pediatric Gastroenterology, Hepatology, and Nutrition. Abstracts. Boston, Massachusetts, USA, August 5-9, 2000. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl 2):S75.

217. Lindfors K, Halttunen T, Kainulainen H, Mäki M. CC3/TIP30 and rab11: novel markers for in vitro and in vivo intestinal epithelial cell differentiation. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S21.
218. Liu J, Juo S-H H, Holopainen P, Tong X, Grunn A, Brito ME, Mustalahti K, Mäki M, Gilliam CT, Partanen J. A genome-wide mapping on celiac disease in 60 Finnish families. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S5.
219. Luostarinen L, Dastidar P, Collin P, Peräaho M, Erilä T, Mäki M, Pirttilä T. Is coeliac disease associated with epilepsy and intracerebral calcifications? Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S20.
220. Mustalahti K, Lohiniemi S, Collin P, Mäki M. Gluten-free diet improves the quality of life of silent celiac disease patients. World Congress of Pediatric Gastroenterology, Hepatology, and Nutrition. Abstracts. Boston, Massachusetts, USA, August 5-9, 2000. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl 2):S142.
221. Mustalahti K, Lohiniemi S, Collin P, Mäki M. Improving quality of life of silent celiac disease (CD) patients during gluten-free diet warrants screening. Ninth International Symposium on Celiac Disease, Hunt Valley, Maryland, USA, August 10-13, 2000. Abstract Book. *J Pediatr Gastroenterol Nutr* 2000;31(Suppl3):S7.
222. Huotari P, Lindfors K, Halttunen T, Mäki M, Kainulainen H. Is human AATF a multifunctional regulator of epithelial cell differentiation? ESPGHAN 34th Annual Meeting, Geneva, Switzerland 9-12 May 2001. *J Pediatr Gastroenterol Nutr* 2001;32:358.
223. Kaukinen K, Halme L, Collin P, Färkkilä M, Partanen J, Mäki M, Höckerstedt K. Coeliac disease in patients with severe liver failure. ESPGHAN 34th Annual Meeting, Geneva, Switzerland 9-12 May 2001. *J Pediatr Gastroenterol Nutr* 2001;32:385.
224. Korponay-Szabo IR, Halttunen T, Laurila K, Szalai Zs, Kovacs JB, Mäki M. Capillary blood test for coeliac disease. ESPGHAN 34th Annual Meeting, Geneva, Switzerland 9-12 May 2001. *J Pediatr Gastroenterol Nutr* 2001;32:361.
225. Korponay-Szabo IR, Halttunen T, Laurila K, Szalai Zs., Kovacs JB, Mäki M. Coeliac disease autoantibody detection with patient erythrocyte transglutaminase in capillary blood. In: Mäki M, Tossavainen M, eds. Proceedings of the Workshop on Transglutaminases, Protein Cross-Linking and Coeliac Disease, September 14-15, 2001, Tampere, Finland. Coeliac Disease Study Group, University of Tampere. Tampere, 2001;173.
226. Korponay-Szabo IR, Szondy Zs, Laurila K, Halttunen T, Sarang Zs, Kovacs JB, Fesus L, Mäki M. Coeliac autoantibody testing on tissue transglutaminase knockout mouse tissues. ESPGHAN 34th Annual Meeting, Geneva, Switzerland 9-12 May 2001. *J Pediatr Gastroenterol Nutr* 2001;32:361.
227. Lähdeaho M-L, Peltonen S, Collin P, Kaukinen K, Ruuska T, Haapala A-M, Mäki M. The development of small bowel lesions in coeliac disease. ESPGHAN 34th Annual Meeting, Geneva, Switzerland 9-12 May 2001. *J Pediatr Gastroenterol Nutr* 2001;32:362.
228. Mustalahti K, Rannila T, Kuorilehto M, Mäki M. Finnish founder population with coeliac disease. In: Mäki M, Tossavainen M, eds. Proceedings of the Workshop on Transglutaminases, Protein Cross-Linking and Coeliac Disease, September 14-15, 2001, Tampere, Finland. Coeliac Disease Study Group, University of Tampere. Tampere, 2001;177.
229. Mustalahti K, Rannila T, Kuorilehto M, Mäki M. Finnish founder population with coeliac disease. Abstract book of Annual Meeting of NAPGAN; Helsingör, Denmark. 30.9.-1.10. 2001.
230. Gadhavi PL, Korponay-Szabo I, Halttunen T, Mäki M. Preliminary characterisation of the IgG fraction in IgA-deficient coeliacs disease patient sera. International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P76.
231. Haimila K, Holopainen P, Mustalahti K, Mäki M, Partanen J. Candidate genes CD28, CTLA4 and ICOS in celiac disease families. 10th International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002;O1.
232. Halttunen T, Huotari , Mäki M. Mouse TG-2 antibodies and recombinant coeliac disease (CD) IgA increase fibroblast proliferation but 'natural' CD-IgA has no effect. International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P78.
233. Halttunen T, Huotari P, Mäki M. Mouse TG-2 antibodies and recombinant coeliac disease (CD) IgA increase fibroblast proliferation but "natural" CD-IgA has no effect. 7th International Conference on Transglutaminases and Protein Crosslinking Reactions. *Minerva Biotechnologica* 2002;14:226.
234. Halttunen T, Marzari R, Sblattero D, Mäki M. Biological functions of coeliac disease autoantibodies. In: Conrad K et al, eds. From Proteomics to Molecular Epidemiology: Relevance of Autoantibodies. Report on the 6th Dresden Symposium on Autoantibodies held in Dresden on September 4 - 7, 2002. *Autoimmunity, autoantigens, autoantibodies* 2002;3:P502.
235. Heinonen T, Lindfors K, Pasternack L, Korponay-Szabo I, Halttunen T, Kainulainen H, Mäki M. A novel glycosyltransferase isolated from differentiating T84 intestinal epithelial cells. 10th International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P45.
236. Järvinen T, Kaukinen K, Hakanen M, Mäki M, Laurila K, Collin P. Intraepithelial lymphocytes in coeliac disease. International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P125.

237. Juuti-Uusitalo K, Kainulainen H, Halttunen T, Collin P, Kaukinen K, Peräaho M, Vihinen M, Ollila J, Mäki M. Gene expression studies with cDNA microarray in jejunal biopsies and in experimental crypta-villus axis model. 10th International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002;O2.
238. Korponay-Szabo I, Halttunen T, Szalai Z, Kovacs J, Maki M. In vivo targeting of jejunal and extrajejunal transglutaminase 2. International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P85.
239. Korponay-Szabo I, Halttunen T, Szalai Zs, Kovacs J, Mäki M. In vivo autoantibody targeting of transglutaminase 2 in coeliac disease. 7th International Conference on Transglutaminases and Protein Crosslinking Reactions. *Minerva Biotechnologica* 2002;14:226.
240. Korponay-Szabo I, Laurila K, Szondy Z, Dahlbom I, B.Kovacs J, Fésüs L, Maki M. Is endomysial antibody (EMA) positivity in patients without villous atrophy due to antibodies against transglutaminase (TG2)? International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P84.
241. Korponay-Szabo I, Laurila K, Szondy Zs, Szalai Zs, Dahlbom I, Fesus L, Mäki M. Specificity of the endomysial antibody reaction (EMA) for transglutaminase 2. 7th International Conference on Transglutaminases and Protein Crosslinking Reactions. *Minerva Biotechnologica* 2002;14:226.
242. Korponay-Szabó IR, Halttunen T, Szalai Z, Kovács JB, Mäki M. In vivo targeting of jejunal and extrajejunal tissue transglutaminase by coeliac IgA autoantibodies. ESPGHAN 35th Annual Meeting, Taormina, Italy, June 5-8, 2002. *J Pediatr Gastroenterol Nutr* 2002;34:449.
243. Lindfors K, Halttunen T, Kainulainen H, Mäki M. Wanted: AATF transcriptional regulators. 10th International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P47.
244. Mäki M, Mustalahti K, Kokkonen J, Kulmala P, Haapalahti M, Karttunen T, Ilonen J, Laurila K, Dahlbom I, Hansson T, Höpfl P, Knip M. Coeliac disease among Finnish Schoolchildren. 10th International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P21.
245. Mustalahti K, Kokkonen J, Kulmala P, Haapalahti M, Ilonen J, Karttunen T, Laurila K, Dahlbom I, Hansson T, Knip M, Mäki M. Coeliac disease among Finnish schoolchildren. ESPGHAN 35th Annual Meeting, Taormina, Italy, June 5-8, 2002. *J Pediatr Gastroenterol Nutr* 2002;34:463
246. Niittynen M, Lindfors K, Mäki M, Kainulainen H. 3G3T, a novel transcript expressed in differentiated T84 cells. International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P49.
247. Peräaho M, Kaukinen K, Vuolteenaho N, Mäki M, Collin P. Oats and quality of life. International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P151.
248. Woolley N, Holopainen P, Ollikainen V, Mustalahti K, Mäki M, Kere J, Partanen J. Genome scan for coeliac disease in a Finnish founder family. 10th International Symposium on Coeliac Disease, Paris, June 2-5, 2002. Programme and abstract book 2002:P30.
249. Juuti-Uusitalo Kati, Markku Mäki, Katri Kaukinen, Markku Peräaho, Juha Ollila, Pekka Collin, Mauno Vihinen, Heikki Kainulainen. cDNA MICROARRAY METHOD IN STUDYING THE COELIAC DISEASE PATHOGENESIS. IMT Research Day, Tuesday 16 December 2003, Tampere.
250. Korponay-Szabo I, B.Kovács J, Opre J, Maki M. Point-of-care test for coeliac disease. *Gut* 2003, 52; Suppl II. A66. 11th United European Gastroenterology Week, Madrid, 2003.
251. Korponay-Szabó I, Dahlbom I, Laurila K, Koskinen S, Karell K, Partanen J, Kovács J B, Mäki M, Hansson T. IgG transzglutamináz autoantitest vizsgálat diagnosztikus szerepe szelektív IgA hiányban. Hungarian Paediatric Gastroenterology Meeting, Szolnok, 2003.
252. Korponay-Szabó I, Opre J, Laurila K, B.Kovács J, Maki M. Helyszíni (point-of-care) coeliakia szerológiai teszt. Hungarian Paediatric Gastroenterology Meeting, Szolnok, 2003.10.11.
253. Korponay-Szabó I. T.Halttunen, K Laurila, I. Dahlbom, L. Fésüs, M. Mäki Transglutaminase 2 antibodies: identity and in vivo action. Tampere University Hospital Research Day, 2003.01.29.
254. Korponay-Szabó Ilma, Ingrid Dahlbom, Kaija Laurila, Sinikka Koskinen, Kati Karrell, Jukka Partanen, B. Kovács Judit, Markku Mäki, Tony Hansson. IgG transzglutamináz autoantitest vizsgálat diagnosztikus szerepe szelektív IgA hiányban. Hungarian Paediatric Gastroenterology Meeting, Szolnok, 2003.10.11.
255. Korponay-Szabo IR, Dahlbom I, Laurila K, Koskinen S, Karell K, Partanen J, Kovacs JB, Maki M, Hansson T. Elevation of IgG antibodies against tissue transglutaminase as a diagnostic tool for coeliac disease in selective IgA deficiency. *J Pediatr Gastroenterol Nutr* 2003;34:Suppl 36th Annual Meeting of the ESPGHAN, Prague, 2003.
256. Viiri KM, Lindfors K, Laiti M, Niittynen M, Heinonen TYK, Mäki M & Kainulainen H. WHAT'SAP? mSin3A associated protein 30-like (SAP30L) - a new member in mSin3A-HDAC-corepressor complex. IMT Research Day, Tuesday 16 December 2003, Tampere.
257. Collin P, Kaukinen K, Thorell L, Mäki M. The safe threshold for gluten contamination in gluten-free products. Abstracts (P172). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
258. Collin P, Kaukinen K, Vogelsang H, Korponay-Szabó I, Sommer R, Schreier E, Volta U, Granito A, Veronesi L, Mascart F, Ocmant A, Ivarsson A, Lagerqvist C, Bürgin-Wolff A, Hadziselimovic F, Furlano R I, Sidler MA, Mulder C.J.J., Goerres MS, Mearin M.L., Ninaber M.K, Gudmand-Hoyer E, Fabiani E, Catassi C, Tidlund H, Alaintalo L, Mäki M. Small bowel biopsy, anti-tissue-transglutaminase and anti-endomysial antibodies in the diagnosis of coeliac disease. A European multicentre study. Abstracts (P13). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.

259. Jarvinen TT, Collin P, Rasmussen M, Kyronpalo S, Maki M, Partanen J, Reunala T, Kaukinen K. Villous tip intraepithelial lymphocyte count in detecting coeliac disease without villous atrophy. Abstracts (P6). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
260. Kaukinen K, Peräaho M, Collin P, Partanen J, Halttunen T, Mäki M, Korponay-Szabo I. Detection of coeliac disease without villous atrophy. Prospective and randomized study. Abstracts (P4). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
261. Korponay-Szabó IR, Halttunen T, Kovács J B, Mäki M. IgM autoantibody deposition on jejunal transglutaminase in IgA deficient coeliac patients. Abstracts (P94). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
262. Korponay-Szabó IR, Laurila K, Opre J, Kovács J B, Mäki M. Diet compliance evaluation of coeliac patients in point of care fashion. Abstracts (P15). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
263. Lähdeaho M-L, Kaukinen K, Collin P, Hartikainen M, Peltonen S, Partanen J, Haapala A-M, Mäki M. The development of small bowel lesion in coeliac disease: A long term follow-up study. Abstracts (P5). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
264. Lohi S, Mustalahti K, Laurila K, Lohi O, Rissanen H, Reunanen A, Mäki M. Prevalence and mortality of undetected coeliac disease. Abstracts (P65). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
265. Mustalahti K, Reunanen A, Heuer M, Metzger M-H, Fabiani E, Catassi C, Murray L, McMillan S, Caradonna M, Bravi E, Mäki M, and the members of the Coeliac EU Cluster, Epidemiology. Tampere, Finland, Freiburg, Germany, Ancona and Trieste, Italy and Belfast, Northern Ireland. Prevalence of coeliac disease in four European countries. Abstracts (P60). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
266. Peräaho M, Collin P, Kaukinen K, Kekkonen L, Miettinen S, Mäki M. Oats forms an essential part of gluten-free diet in coeliac disease and dermatitis herpetiformis. Abstracts (P173). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
267. Raivio T, Kaukinen K, Korponay-Szabó I, Laurila K, Collin P, Mäki M. Rapid point of care test for coeliac disease. Abstracts (P3). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
268. Siekkinen H, Heinonen T, Mäki M, Kainulainen H. A novel glycosyltransferase, B3GTL, isolated from differentiated T84 intestinal epithelial cells. 7th annual GLIBS meeting 1-3.9.2004, Fiskebäckskil.
269. Siekkinen H, Heinonen T, Mäki M, Kainulainen H. A novel glycosyltransferase, B3GTL, isolated from differentiated T84 intestinal epithelial cells. 5th International EMBL PhD Students Symposium 2.-4.12.2004, Heidelberg.
270. Viiri K, Laiti M, Lindfors K, Lohi O, Peterson P, Mäki M, Kainulainen H. mSin3A-associated protein 30-like (SAP30L) – a new member in mSin3A-HDAC corepressor complex. 5th International EMBL PhD Students Symposium 2.-4.12.2004, Heidelberg.
271. Myrsky E, Lindfors K, Kainulainen H, Kaukinen K, Mäki M: Study on small intestinal epithelial stem cells in normal and coeliac disease affected small intestinal mucosa. II Tissue Engineering Symposium 12.4.2005, Tampere.
272. Vuolteenaho N, Kekkonen L, Aspegren M, Kaukinen K, Mäki M, Collin P. How to detect coeliac disease: population screening or clinical work-up? Abstracts (P57). 11th International Symposium on Coeliac Disease. Belfast, 28-30 April 2004.
273. Korponay-Szabo IR, Hadjivassiliou M, Williamson CA, Laurila K, Haimila K, Woodroffe N, Sanders DS, Mäki M. Gluten ataxia: intestinal and brain targeting by tissue transglutaminase antibodies. Abstracts. 38<sup>th</sup> Annual Meeting of the European Society for Pediatric Gastroenterology, Hepatology and Nutrition. Porto, Portugal, June 1-4, 2005. *J Pediatr Gastroenterol Nutr* 2005;40:616 (OP1-02).
274. Kaukinen K, Holm K, Villanen M, Korponay-Szabo I, Mäki M. Small bowel mucosal transglutaminase autoantibodies during oats and gluten challenge in coeliac disease. Abstracts. 38<sup>th</sup> Annual Meeting of the European Society for Pediatric Gastroenterology, Hepatology and Nutrition. Porto, Portugal, June 1-4, 2005. *J Pediatr Gastroenterol Nutr* 2005;40:618 (OP2-01).
275. Korponay-Szabo IR, Raivio TM, Laurila K, Kovacs JB, Nemes E, Kaukinen K, Mäki M. Rapid detection of coeliac autoantibodies in the office. Abstracts. 38<sup>th</sup> Annual Meeting of the European Society for Pediatric Gastroenterology, Hepatology and Nutrition. Porto, Portugal, June 1-4, 2005. *J Pediatr Gastroenterol Nutr* 2005;40:619 (OP2-03).
276. Kaukinen K, Viljamaa M, Huhtala H, Mäki M, Collin P. Is celiac disease screening in risk groups justified? Abstracts. 38<sup>th</sup> Annual Meeting of the European Society for Pediatric Gastroenterology, Hepatology and Nutrition. Porto, Portugal, June 1-4, 2005. *J Pediatr Gastroenterol Nutr* 2005;40:637 (PG1-14).
277. Myrsky E, Lindfors K, Kainulainen H, Mäki M. A cell culture model for small intestinal stem cell function and differentiation. Abstracts. 38<sup>th</sup> Annual Meeting of the European Society for Pediatric Gastroenterology, Hepatology and Nutrition. Porto, Portugal, June 1-4, 2005. *J Pediatr Gastroenterol Nutr* 2005;40:661 (PG4-19).
278. Viiri KM, Laiti M, Kokeamäki H, Nieminen LK, Lindfors K, Peterson P, Mäki M, Kainulainen H, Lohi O. SIN3A-associated protein 30-like (SAP30L) is degraded in the nucleolus and enhances the repression capability of the SIN3A-HDAC corepressor complex. 1st EMBL Monterotondo International PhD Students Minisymposium, Animal Models, Tips and Tricks from Nature. 12. - 14.5.2005, Rome, Italy.

279. Siekkinen H, Heinonen T, Mäki M, Kainulainen H. A novel glycosyltransferase, B3GTL, isolated from differentiated T84 intestinal epithelial cells. 1<sup>st</sup> International EMBL Monterotondo International PhD Students Minisymposium, Animal Models, Tips and Tricks from Nature. 12.-14.5.2005, Rome, Italy.
280. Myrsky E, Lindfors K, Kainulainen H, Mäki M. Study on small intestinal epithelial stem cells in normal and coeliac disease affected small intestinal mucosa. Kudosteknologiasymposium 12.4.2005, Tampere.
281. Raivio T, Kaukinen K, Korponay-Szabó I, Laurila K, Mäki M. Innovative rapid point of care test in coeliac disease antibody detection. Valtakunnalliset lääkäripäivät, Helsinki 9.-13.1.2005.
282. Raivio T, Kaukinen K, Korponay-Szabó I, Laurila K, Mäki M. Innovative rapid point of care test in coeliac disease antibody detection. Gastropäivät 17.-18.2005, Jyväskylä.
283. Raivio T, Kaukinen K, Korponay-Szabó I, Laurila K, Mäki M. Innovative rapid point of care test in coeliac disease antibody detection. Tampereen lääkäripäivät 17.-19.3.2005, Tampere.
284. Raivio T, Kaukinen K, Laurila K, Collin P, Kovacs J, Mäki M, Korponay-Szabó IR. Self transglutaminase-based rapid coeliac antibody detection in the office. Biannual meeting of NAPGHAN, Nordic Association for Paediatric Gastroenterology, Hepatology and Nutrition. 24-25 November 2005, Helsinki.
285. Juuti-Uusitalo K, Kaukinen K, Mäki M, Kainulainen H. Gene expression profiling in TGF $\beta$ -mediated T84 epithelial cell differentiation in the three-dimensional crypt-villus axis model. Murikka Summer School, August 21st-28<sup>th</sup>, 2005, Tampere.
286. Juuti-Uusitalo K, Kaukinen K, Mäki M, Kainulainen H. Gene expression profiling in TGF $\beta$ -mediated T84 epithelial cell differentiation in the three-dimensional crypt-villus axis model. XX Paulo Symposium, Targeting and activation of TGF $\beta$  family growth and differentiation factors, 2nd -3rd September 2005, Helsinki.
287. Siekkinen H, Heinonen T, Mäki M, Kainulainen H. A glycosyltransferase, B3GTL, isolated from differentiated T84 intestinal epithelial cells. Murikka Summer School, August 21-28, 2005, Tampere.
288. Siekkinen H, Heinonen T, Mäki M, Kainulainen H. A glycosyltransferase, B3GTL, isolated from differentiated T84 intestinal epithelial cells. Finbionet National PhD Student Symposium, 6-7.10.2005, Vammala.
289. Siekkinen H, Heinonen T, Mäki M, Kainulainen H. A glycosyltransferase, B3GTL, isolated from differentiated T84 intestinal epithelial cells. IMT Research Day, 14.12.2005, Tampere.
290. Lindfors K, Myrsky E, Kainulainen H, Mäki M. A cell culture system modelling the entire small intestinal crypt villus axis. Digestive Disease Week 2006, May 20-25, 2006, Los Angeles.
291. Lindfors K, Myrsky E, Syrjänen M, Korponay-Szabó I, Mäki M, Kaukinen K. Celiac disease IGA class autoantibodies disturb normal capillary formation in angiogenesis. Digestive Disease Week 2006, May 20-25, 2006, Los Angeles.
292. Lohi S, Mustalahti K, Kaukinen K, Laurila K, Collin P, Rissanen H, Lohi O, Bravi E, Reunanen A, Mäki M. The prevalence of celiac disease is increasing in process of time. Digestive Disease Week 2006, May 20-25, 2006, Los Angeles.
293. Myrsky E, Syrjänen M, Kaukinen K, Korponay-Szabó I, Mäki M, Lindfors K. Celiac disease IgA class autoantibodies inhibit the recruitment of vascular smooth muscle cells to capillary walls. The 39th Annual Meeting of ESPGHAN, June 7-10, 2006, Dresden.
294. Stenman S, Lindfors K, Korponay-Szabó I, Mäki M, Kaukinen K. Production of coeliac disease autoantibodies after in vitro gluten challenge is dependent on small-bowel mucosal transglutaminase 2-specific IgA deposits. The 39th Annual Meeting of ESPGHAN, June 7-10, 2006, Dresden.
295. Lohi S, Mustalahti K, Kaukinen K, Laurila K, Collin P, Rissanen H, Lohi O, Bravi E, Gasparin M, Reunanen A, Mäki M. Increasing prevalence of celiac disease in process of time. 8. Gastropäivät, 16.-17.2.2006, Helsinki.
296. Vilppula A, Collin P, Mäki M, Valve R, Luostarinen M, Krekelä I, Kaukinen K, Luostarinen L. Keliakian esiintyvyys ja taudinkuva ikääntyvässä väestössä. 8. Gastropäivät, 16.-17.2.2006, Helsinki.
297. Stenman SM, Lindfors K, Korponay-Szabó IR, Mäki M, Kaukinen K. Production of coeliac disease autoantibodies after in vitro gluten challenge is dependent on small-bowel mucosal transglutaminase 2-specific IgA deposits. 8. Gastropäivät, 16.-17.2.2006, Helsinki.
298. Myrsky E, Kaukinen K, Mäki M, Lindfors K. A cell culture system modelling the entire small intestinal crypt villus axis. 8. Gastropäivät, 16.-17.2.2006, Helsinki.
299. Salmi TT, Collin P, Järvinen O, Haimila K, Partanen J, Laurila K, Korponay-Szabó IR, Huhtala H, Reunala T, Mäki M, Kaukinen K. IgA autoantibodies against transglutaminase 2 in the small intestinal mucosa predict forthcoming coeliac disease. 8. Gastropäivät, 16.-17.2.2006, Helsinki.
300. Koskinen L, Korponay-Szabó I, Mustalahti K, Lindfors K, Kaukinen K, Kurppa K, Juuti-Uusitalo K, Kere J, Wijmenga C, Mäki M, Partanen J, Holopainen P. Linkage and association study of MYO9B variants in Finnish and Hungarian populations supports the role of CELIAC4 locus (19p13) in celiac disease. HGM2006 Workshop Abstracts. Human Genome Meeting, Helsinki 31.5.-3.6.2006.
301. Myrsky E, Lindfors K, Kainulainen H, Kaukinen K, Mäki M. A cell culture system modelling the entire small intestinal crypt-villus axis. 20th Anniversary Symposium of the Finnish Cancer Institute: Stem cells and cancer. Biomedicum Helsinki, 1 Aug. 31-Sept. 1, 2006.
302. Myrsky E, Kaukinen K, Syrjänen M, Korponay-Szabó IR, Mäki M, Lindfors K. Celiac disease autoantibodies disturb normal capillary formation in angiogenesis. 1st Nordic Meeting on Genetics and Pathogenesis of Immunological Diseases. 4-5 September 2006, Helsinki.

303. Juuti-Uusitalo K, Mäki M, Kainulainen H, Isola J, Kaukinen K. Common set of genes regulated in coeliac disease and in three-dimensional crypt-villus axis model. Program and Abstract Book. 1st Nordic Meeting on Genetics and Pathogenesis of Immunological Diseases. 4-5 September 2006, Helsinki.
304. Koskinen L, Viiri K, Korponay-Szabo I, Mustalahti K, Kurppa K, Kaukinen K, Wijmenga C, Kere J, Mäki M, Partanen J, Holopainen P. Linkage and association study of MYO9B variants in Finnish and Hungarian populations supports the role of celiac4 locus (19P13) in celiac disease. Program and Abstract Book. 1st Nordic Meeting on Genetics and Pathogenesis of Immunological Diseases. 4-5 September 2006, Helsinki.
305. Kondrashova A, Mustalahti K, Kaukinen K, Viskari H, Volodicheva V, Haapala A-M, Ilonen J, Knip M, Mäki M, Hyöty H, the EPIVIR Study Group. Conspicuous difference between Russian Karelia and Finland in the prevalence of celiac disease among children. Program and Abstract Book. 1st Nordic Meeting on Genetics and Pathogenesis of Immunological Diseases. 4-5 September 2006, Helsinki.
306. Myrsky E, Kaukinen K, Syrjänen M, Korponay-Szabo IR, Mäki M, Lindfors K. Celiac disease specific IgA class autoantibodies disturb normal angiogenesis. Lastenlääkäripäivät 26.-27.10.2006, Tampere.
307. Mäki M, Hernando A, Mena MC, Lombardia M, Lehtonen P, Pöyri S, Soininen-Tengvall P, Pajunen E, Mendez E. Barley malt beer rendered gluten free. XII International Celiac Disease Symposium 2006, November 9-11, 2006, New York.
308. Mustalahti K, Rask S, Mattila A, Mäki M, Joukamaa M. Depression and alexithymia in screen detected coeliac disease patients. XII International Celiac Disease Symposium 2006, November 9-11, 2006, New York.
309. Juuti-Uusitalo K, Mäki M, Kainulainen H, Isola J, Kaukinen K. A common set of genes regulated in epithelial cell differentiation and celiac disease. IMT Research Day, 19 December 2006, Tampere.
310. Myrsky E, Kaukinen K, Syrjänen M, Korponay-Szabo IR, Mäki M, Lindfors K. Coeliac disease IgA class autoantibodies disturb normal angiogenesis. IMT Research Day, 19 December 2006, Tampere.
311. Stenman S, Lindfors K, Korponay-Szabo IR, Wieser H, Mäki M, Kaukinen K. Production of celiac disease autoantibodies after in vitro gliadin challenge is dependent on small-bowel mucosal transglutaminase 2-specific IgA deposits. IMT Research Day, 19 December 2006, Tampere.
312. Viiri KM, Nieminen LK, Korkeamäki H, Kouvonen P, Lahesmaa R, Kainulainen H, Mäki M, Lohi O. SIN3A corepressor act on 60S ribosome biogenesis and is physically linked to the 60S pre-ribosomes by SAP30 and SAP30L proteins and this recruitment is necessary for the cell cycle progression. IMT Research Day, 19 December 2006, Tampere.
313. Einarsdottir E, Koskinen L, Korponay-Szabo I, Mustalahti K, Kurppa K, Kere J, Mäki M, Partanen J, Holopainen P. A whole-genome scan reveals significant linkage of celiac disease to 3p14-p13 and 2q14-q23 in extended pedigrees from Hungary and Finland. 40th Annual Meeting of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition, 9-12 May 2007, Barcelona, Spain. *J Pediatr Gastroenterol Nutr* 2007;44:e52.
314. Einarsdottir E, Koskinen L, Korponay-Szabo I, Mustalahti K, Kurppa K, Partanen J, Mäki M, Kere J, Holopainen P. A whole-genome scan reveals linkage of celiac disease to 6q21-22 and 22q13 in extended pedigrees from Hungary and Finland. ASHG, San Diego.
315. Einarsdottir E, Koskinen L, Korponay-Szabo I, Mustalahti K, Kurppa K, Partanen J, Mäki M, Kere J, Holopainen P. A whole-genome scan reveals linkage of celiac disease to 6q21-22 and 22q13 in extended pedigrees from Hungary and Finland. ESHG Annual Meeting, 16.-19.6.2007, Nice, France.
316. Iltanen S, Ashorn S, Raukola H, Välineva T, Kaukinen K, Rantala I, Wei B, Braun J, Collin P, Mäki M, Luukkaala T, Ashorn M. Serological responses to bacterial and autoantigens in coeliac disease patients during a gluten-free diet. In: Dolinsek J, Ornik T, eds. Proceedings of the International Coeliac Disease Meeting, Maribor 13-16 September 2007, University Medical Center, Maribor 2007:161.
317. Juuti-Uusitalo K, Mäki M, Lindfors K, Venäläinen J, Isola J, Kaukinen K. Gliadin affects on epithelial cell junctions via ErbB. IMT Research Day, University of Tampere, December 18, 2007, Tampere.
318. Juuti-Uusitalo K, Mäki M, Kainulainen H, Isola J, Kaukinen K. Genes regulated in an in vitro epithelial cell differentiation model operate also in gluten-induced small intestinal lesion in coeliac disease. Suomen Gastroenterologiayhdistys ry, 9. Gastropäivät 15.-16.2.2007, Helsinki.
319. Juuti-Uusitalo K, Mäki M, Kainulainen H, Isola J, Kaukinen K. Genes regulated in an in vitro epithelial cell differentiation model operate also in gluten-induced small intestinal lesion in coeliac disease. 40th Annual Meeting of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition, 9-12 May 2007, Barcelona, Spain. *J Pediatr Gastroenterol Nutr* 2007;44:e218.
320. Juuti-Uusitalo K, Mäki M, Kainulainen H, Isola J, Kaukinen K. Genes regulated in an in vitro epithelial cell differentiation model operate also in gluten-induced small intestinal lesion in coeliac disease. The XXXVIII Nordic Meeting of Gastroenterology, Reykjavik, Iceland 6-9 June 2007. *Scand J Gastroenterol* 2007;42(Suppl244):26.
321. Korponay-Szabo IR, Kovacs J, Kiraly R, Nemes E, Mäki M. High efficiency of gluten-dependent transglutaminase-specific intestinal IgA deposits as candidate diagnostic criteria in coeliac disease. 40th Annual Meeting of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition, 9-12 May 2007, Barcelona, Spain. *J Pediatr Gastroenterol Nutr* 2007;44:e50.
322. Korponay-Szabo IR, Kovacs J, Nemes E, Dahlbom I, Mäki M. Coeliac antibody testing with deamidated gliadin peptides in difficult patient samples. 40th Annual Meeting of the European Society for Paediatric

- Gastroenterology, Hepatology and Nutrition, 9-12 May 2007, Barcelona, Spain. *J Pediatr Gastroenterol Nutr* 2007;44:e226.
323. Korponay-Szabo IR, Kovacs J, Kiraly R, Nemes E, Mäki M. High efficiency of gluten-dependent transglutaminase-specific intestinal IgA deposition as candidate diagnostic criteria in coeliac disease. "15th United European Gastroenterology week" Paris, France 27-31 October 2007. *Gut* 2007;56:(SupplIII) A109.
  324. Korponay-Szabo IR, Kovacs J, Nemes E, Dahlbom I, Mäki M. Coeliac antibody testing with deamidated gliadin peptides in difficult patient samples. In: Dolinsek J, Ornik T, eds. Proceedings of the International Coeliac Disease Meeting, Maribor 13-16 September 2007, University Medical Center, Maribor 2007:179.
  325. Korponay-Szabo IR, Kovacs J, Nemes E, Dahlbom I, Mäki M. Coeliac antibody testing with deamidated gliadin peptides in difficult patient samples. "15th United European Gastroenterology week" Paris, France 27-31 October 2007. *Gut* 2007;56:(SupplIII) A109.
  326. Korponay-Szabo IR, Vecsei Z, Kiraly R, Dahlbom I, Mäki M. Homology of deamidated gliadin peptides and tissue transglutaminase. 40th Annual Meeting of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition, 9-12 May 2007, Barcelona, Spain. *J Pediatr Gastroenterol Nutr* 2007;44:e227.
  327. Korponay-Szabo IR, Vecsei Z, Kiraly R, Nemes E, Dahlbom I, Fesus L, Mäki M. Homology of deamidated gliadin peptides and tissue transglutaminase. In: Dolinsek J, Ornik T, eds. Proceedings of the International Coeliac Disease Meeting, Maribor 13-16 September 2007, University Medical Center, Maribor 2007:163.
  328. Korponay-Szabo IR, Vecsei Z, Kiraly R, Nemes E, Dahlbom I, Fesus L, Mäki M. Homology of deamidated gliadin peptides and tissue transglutaminase. "15th United European Gastroenterology week" Paris, France 27-31 October 2007. *Gut* 2007;56:(SupplIII) A109.
  329. Koskinen L, Korponay-Szabo I, Mustalahti K, Kurppa K, Kaukinen K, Einarsdottir E, Mäki M, Partanen J, Kere J, Holopainen P. Linkage and association study of the CELIAC2 locus (5q31-33) in Hungarian and Finnish families with celiac disease. The European Society of Human Genetics (ESHG) Annual Meeting, 16.-19.6.2007, Nice, France.
  330. Koskinen L, Korponay-Szabo I, Viiri K, Juuti-Uusitalo K, Kaukinen K, Lindfors K, Mustalahti K, Kurppa K, Adany R, Pocsai Z, Szeles G, Wijmenga C, Mäki M, Partanen J, Kere J, Holopainen P. MYO9B gene at 19P13 (CELIAC4) shows genetic linkage and association with dermatitis herpetiformis, the skin manifestation of celiac disease. Suomen Gastroenterologiyhdistys ry, 9. Gastropäivät 15.-16.2.2007, Helsinki.
  331. Kurppa K, Collin P, Viljamaa M, Haimila K, Partanen J, Saavalainen P, Laurila K, Mäki M, Kaukinen K. Treatment of endomysial antibody-positive patients without villous atrophy: are current diagnostic criteria for celiac disease valid? In: Dolinsek J, Ornik T, eds. Proceedings of the International Coeliac Disease Meeting, Maribor 13-16 September 2007, University Medical Center, Maribor 2007:141.
  332. Kurppa K, Mäki M, Collin P, Reunala T, Kaukinen K. A phenotype of celiac disease may change from classical disorder to dermatitis herpetiformis after a long asymptomatic period with gluten challenge. 40th Annual Meeting of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition, 9-12 May 2007, Barcelona, Spain. *J Pediatr Gastroenterol Nutr* 2007;44:e74.
  333. Lindfors K, Blomqvist T, Stenman S, Kaukinen K, Mäki M. Live bacteria of a probiotic bacterial strain inhibit gliadin-induced damage. 40th Annual Meeting of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition, 9-12 May 2007, Barcelona, Spain. *J Pediatr Gastroenterol Nutr* 2007;44:e67.
  334. Lohi S, Mäki M, Laurila K, Montonen J, Bravi E, Gasparin M, Knekt P, Reunanen A, Kaukinen K. Malignancy in unrecognized celiac disease: population based cohort study. Suomen Gastroenterologiyhdistys ry, 9. Gastropäivät 15.-16.2.2007, Helsinki.
  335. Lohi S, Mäki M, Laurila K, Montonen J, Bravi E, Gasparin M, Knekt P, Reunanen A, Kaukinen K. Malignancy in unrecognized celiac disease: population based cohort study. 40th Annual Meeting of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition, 9-12 May 2007, Barcelona, Spain. *J Pediatr Gastroenterol Nutr* 2007;44:e22.
  336. Mustalahti K, Hällström M, Mäki M. Characteristics of silent coeliac disease. In: Dolinsek J, Ornik T, eds. Proceedings of the International Coeliac Disease Meeting, Maribor 13-16 September 2007, University Medical Center, Maribor 2007:168.
  337. Myrsky E, Kaukinen K, Syrjänen M, Korponay-Szabo IR, Mäki M, Lindfors K. Celiac disease-specific IgA class autoantibodies disturb angiogenesis. Suomen Gastroenterologiyhdistys ry, 9. Gastropäivät 15.-16.2.2007, Helsinki.
  338. Myrsky E, Kaukinen K, Syrjänen M, Korponay-Szabo IR, Mäki M, Lindfors K. Coeliac disease-specific iGA class autoantibodies disturb angiogenesis. 9th International Conference of Transglutaminases and Protein Cross-linking, Marrakech September 1-5, 2007.
  339. Myrsky E, Kaukinen K, Syrjänen M, Korponay-Szabo IR, Mäki M, Lindfors K. Coeliac disease-specific IgA class autoantibodies disturb angiogenesis. 8th Dresden symposium on autoantibodies, September 12-15th, 2007, Dresden, Germany.
  340. Myrsky E, Nadalutti C, Kaukinen K, Korponay-Szabo I, Mäki M, Lindfors K. Coeliac disease-specific autoantibodies targeted against transglutaminase 2 increase endothelial permeability in vitro. IMT Research Day, Tampere, December 18, 2007.

341. Nadalutti C, Myrsky E, Viiri K, Kaukinen K, Lindfors K, Mäki M. Is the down-regulation of TG2 the mechanism of coeliac disease IgA autoantibody action? IMT Research Day, December 18th, 2007, Tampere, Finland.
342. Rauhavirta T, Mäki M, Kaukinen K, Lindfors K. Coeliac disease IgA - a wolf in sheep's clothing? IMT Research Day Tampere, December 18, 2007.
343. Saavalainen P, Einarsdottir E, Koskinen L, Korponay-Szabo I, Mustalahti K, Kurppa K, Partanen J, Mäki M, Kere J. A whole-genome scan reveals significant linkage of celiac disease to 6q21-22 and 22q13 in extended pedigrees from Hungary and Finland. In: Dolinsek J, Ornik T, eds. Proceedings of the International Coeliac Disease Meeting, Maribor 13-16 September 2007, University Medical Center, Maribor 2007:132.
344. Saavalainen P, Koskinen L, Korponay-Szabo I, Mustalahti K, Viiri K, Juuti-Uusitalo K, Lindfors K, Kurppa K, Kaukinen K, Adany R, Pocsai Z, Szeles G, Wijmenga C, Mäki M, Partanen J, Kere J. Linkage study of Myosin IXB variants in Hungarian and Finnish populations supports the role of CELIAC4 locus (19p13) in celiac disease. In: Dolinsek J, Ornik T, eds. Proceedings of the International Coeliac Disease Meeting, Maribor 13-16 September 2007, University Medical Center, Maribor 2007:133
345. Stenman S, Lindfors K, Korponay-Szabo IR, Lohi O, Wieser H, Mäki M, Kaukinen K. Production of celiac disease autoantibodies after in vitro gliadin challenge is dependent on small-bowel mucosal transglutaminase 2-specific IgA deposits. 8th Dresden symposium on autoantibodies, September 12-15th, 2007, Dresden, Germany
346. Stenman S, Männistö P, Kaukovirta-Norja A, Lindfors K, Kaukinen K, Mäki M, Venäläinen J. Idätetyistä viljoista eristetyt entsyymit pilkkovat vehnän gliadiinin keliakikoille vaarattomampaan muotoon in vitro. Suomen Gastroenterologiyhdistys ry, 9. Gastropäivät 15.-16.2.2007, Helsinki.
347. Stenman S, Venäläinen J, Männistö P, Lindfors K, Kaukinen K, Mäki M. Proteases from germinating cereals decrease the toxic effects of PT-gliadin in coeliac disease in vitro. 40th Annual Meeting of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition, 9-12 May 2007, Barcelona, Spain. *J Pediatr Gastroenterol Nutr* 2007;44:e48.
348. Stenman S., Venäläinen J., Männistö P., Lindfors K., Kaukinen K. and Mäki M. Proteases from germinating cereals decrease the toxic effects of PT-gliadin in coeliac disease in vitro. IMT Research Day, December 18th, 2007, Tampere, Finland.
349. Viiri K, Siggers T, Jänis J, Berger M, Valjakka J, Mäki M, Bulyk M, Lohi O. PtdInsP modulate DNA-binding of a novel zinc-coordinating module in SAP30L. IMT Research Day, Tampere, December 18, 2007.
350. Zibera F, Braicu O, Kaukinen K, Mäki M, Collin P, Holopainen P. Serum sHLA-G in celiac disease and co-associated autoimmunity. Abstracts, 40th Annual Meeting of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition, 9-12 May 2007, Barcelona, Spain. *J Pediatr Gastroenterol Nutr* 2007;44:e54.
351. Zibera F, Braicu OL, Collin P, Mäki M, Kaukinen K, Saavalainen P. Serum sHLA-G in celiac disease and co-associated autoimmunity. In: Dolinsek J, Ornik T, eds. Proceedings of the International Coeliac Disease Meeting, Maribor 13-16 September 2007, University Medical Center, Maribor 2007:131.
352. Ashorn SM, Iltanen S, Honkanen T, Raukola H, Välineva T, Kaukinen K, Rantala I, Wei B, Braun J, Mäki M, Ashorn M. Serological responses to microbial and autoantigens in children with gut inflammation. 16<sup>th</sup> United European Gastroenterology Week, Vienna, Austria, 18-22 October 2008. *Gut* 2008;57(Suppl II):A125.
353. Caja S, Martucciello S, Lauvric M, Sblattero D, Marzari R, Mäki M, Kaukinen K, Lindfors K. Coeliac disease-specific autoantibodies targeted against transglutaminase 2 disturb vessel morphogenesis in vitro and in vivo through RhoB. IMT Research day 2008, December 17, Tampere.
354. Dahlbom I, Korponay-Szabo IR, Kovacs JB, Szalai Z, Mäki M. Prediction of clinical and mucosal severity of celiac disease by quantification of IgA/IgG serum antibodies to tissue transglutaminase. 16<sup>th</sup> United European Gastroenterology Week, Vienna, Austria, 18-22 October 2008. *Gut* 2008;57(Suppl II):A5.
355. Einarsdottir E, Dukes E, de Kauwe A, Koskinen L, Zibera F, Korponay-Szabo IR, Kurppa K, Mustalahti K, Not T, Vatta S, Ventura A, Adány R, Pocsai Z, Széles G, Mäki M, Kere J, Anderson R, Kaukinen K, Saavalainen P. The IBD risk gene IL23R is linked to and associated with celiac disease and points to shared Th17 pathways in intestinal inflammatory diseases. World Immune Regulation Meeting II, March 17-20, 2008. Poster Abstracts WIRM II, Davos 2008:108.
356. Einarsdottir E, Koskinen L, Dukes E, Kainu K, Zibera F, Korponay-Szabo IR, Kurppa K, Kaukinen K, Adány R, Not T, Ventura A, Löfberg R, Torkvist L, Bresso F, Halfvarson J, Mäki M, Saarialho-Kere U, Kere J, D'Amato M, Saavalainen P. IL23R in the Swedish, Finnish, Hungarian and Italian populations: Association with inflammatory bowel disease and psoriasis, and linkage to celiac disease. 58th Annual Meeting of the American Society of Human Genetics; November 11-15, 2008, Philadelphia, USA.
357. Hamilton G, Mäki M, Lähdeaho M, Bekker P, Schall T. A randomized, double-blind, placebo-controlled, phase II study testing CCX282-B (Traficet-EN) in the treatment of celiac disease. Digestive Disease Week and the 109<sup>th</sup> Annual Meeting of the AGA Institute May 17-22, 2008, San Diego, CA, USA.
358. Kataja L, Kaukinen K, Mäki M, Lindfors K. Oats and celiac disease – study of oats-derived compounds in vitro. IMT Research day 2008, December 17, Tampere.
359. Kivistö R, Rauhavirta T, Mongeot A, Collighan R, Griffin M, Mäki M, Lindfors K. Transglutaminase 2 inhibitors – protecting by preventing? IMT Research day 2008, December 17, Tampere.

360. Korponay-Szabo IR, Nemes E, Vecsei Z, Kiraly R, Mäki M. Rapid bedside detection of humoral IgA deficiency and celiac disease antibodies from capillary blood. 3<sup>rd</sup> World Congress of Pediatric Gastroenterology, Hepatology and Nutrition. 16-20 August 2008, Iguassu Falls, Brazil.
361. Korponay-Szabo IR, Nemes E, Vecsei Z, Mäki M. Rapid bedside detection of humoral IgA deficiency and celiac disease antibodies from capillary blood. 16<sup>th</sup> United European Gastroenterology Week, Vienna, Austria, 18-22 October 2008. *Gut* 2008;57(Suppl II):A4.
362. Korponay-Szabó I, Myrsky E, Caja S, Nemes É, Lindfors K, Mäki M. Az intracelluláris transzglutamináz vizsgálata coeliakiában (Investigation of intracellular transglutaminase in coeliac disease). Annual Paediatric Gastroenterology Meeting, 10-11 October, 2008, Budapest, Hungary.
363. Koskinen LLE, Einarsdottir E, Dukes E, Heap GAR, Dubois P, Korponay-Szabo IR, Kaukinen K, Kurppa K, Ziberna F, Vatta S, Not T, Ventura A, Ádány R, Pocsai Z, Széles G, Mäki M, Kere J, Wijmenga C, van Heel DA, Saavalainen P. Association study of IL18RAP in three European populations with celiac disease. 58th Annual Meeting of the American Society of Human Genetics; November 11-15, 2008, Philadelphia, USA.
364. Kurppa K, Collin P, Huhtala H, Haimila K, Saavalainen P, Partanen J, Laurila K, Mäki M, Kaukinen K. Keliakian diagnostiset kriteerit tulee uudistaa. Yleislääketieteen päivät 2008 Tampereella 23.-24.10.2008.
365. Kurppa K, Collin P, Sievänen H, Huhtala H, Mäki M, Kaukinen K. Reduced quality of life and bone mineral density in mild enteropathy celiac disease warrants new diagnostic criteria. 16<sup>th</sup> United European Gastroenterology Week, Vienna, Austria, 18-22 October 2008. *Gut* 2008;57(Suppl II):A4.
366. Kurppa K, Collin P, Viljamaa M, Haimila K, Saavalainen P, Partanen J, Laurila K, Huhtala H, Paasikivi K, Mäki M, Kaukinen K. Gluteeniherkkyys alkavassa keliakiassa: satunnaistettu, kontrolloitu kliininen tutkimus. XXIX Tampereen Lääkäripäivät Tampereella 13.-15.3.2008, luennot. Tampereen lääkäripäivät 2008 järjestelytoimikunta. Lege Artis Oy, Vammala 2008:263.
367. Kurppa K, Collin P, Viljamaa M, Holopainen P, Haimila K, Laurila K, Paasikivi K, Mäki M, Kaukinen K. Gluten sensitivity of endomysial antibody-positive adults without villous atrophy. Should the current diagnostic criteria of celiac disease be revisited? A prospective, randomized and controlled clinical trial. Digestive Disease Week and the 109<sup>th</sup> Annual Meeting of the AGA Institute May 17-22, 2008, San Diego, CA, USA. *Gastroenterology* 2008;134 (Suppl 1):A-80.
368. Martucciello C, Caja S, Mäki M, Kaukinen K, Lindfors K. The autoantibodies anti-TG2 can modulate gene expression related with angiogenesis in endothelial cells. IMT Research day 2008, December 17, Tampere.
369. Myrsky E, Caja S, Korponay-Szabo I, Collighan R, Mongeot A, Griffin M, Mäki M, Kaukinen K, Lindfors K. Coeliac disease autoantibodies modulate vascular permeability in vitro. IMT Research day 2008, December 17, Tampere.
370. Myrsky E, Rauhavirta T, Kaukinen K, Korponay-Szabo IR, Mäki M, Lindfors K. Celiac disease autoantibodies targeted against transglutaminase 2 increase endothelial permeability. Helsinki Biomedical Graduate School: Cell Structure and Organelles. May 8-9, 2008, Helsinki.
371. Myrsky E, Rauhavirta T, Kaukinen K, Korponay-Szabo IR, Mäki M, Lindfors K. Celiac disease autoantibodies targeted against transglutaminase 2 increase endothelial permeability. 5th International EDHF Symposium, June 24-27, 2008. Tampere, Finland.
372. Myrsky E, Rauhavirta T, Korponay-Szabo I, Kaukinen K, Mäki M, Lindfors K. Celiac disease-specific autoantibodies targeted against transglutaminase 2 increase endothelial permeability in vitro. Digestive Disease Week and the 109<sup>th</sup> Annual Meeting of the AGA Institute May 17-22, 2008, San Diego, CA, USA. *Gastroenterology* 2008;134 (Suppl 1):A-519.
373. Nadalutti C, Viiri K, Kaukinen K, Mäki M, Lindfors K. Downregulation of tissue transglutaminase and its implications for coeliac disease. IMT Research day 2008, December 17, Tampere.
374. Nadalutti C, Viiri K, Kaukinen K, Mäki M, Lindfors K. G1 cell cycle arrest and anoikis induction by siRNA tissue transglutaminase in HUVECs. Murikka Summer School in Molecular Biology 2008, Cell signalling and its transcriptional readout. Tampere, August 12-19, 2008.
375. Rauhavirta T, Kaukinen K, Mäki M, Lindfors K. Coeliac disease specific IgA enhances the transepithelial passage of gliadin peptides. 16<sup>th</sup> United European Gastroenterology Week, Vienna, Austria, 18-22 October 2008. *Gut* 2008;57(Suppl II):A4.
376. Rauhavirta T, Kaukinen K, Korponay-Szabó IR, Mongeot A, Collighan R, Griffin M, Mäki M, Lindfors K. The transepithelial passage of gliadin peptides is enhanced in the presence of coeliac disease-specific IgA. IMT Research day 2008, December 17, Tampere.
377. Rauhavirta T, Mäki M, Kaukinen K, Lindfors K. Coeliac disease specific IgA enhances the transepithelial passage of gliadin peptides. Murikka Minisymposium Cell signalling and its transcriptional readout. Murikka Summer School in Molecular Biology 2008, Cell signalling and its transcriptional readout. Tampere, August 12-19, 2008.
378. Rauhavirta T, Mäki M, Kaukinen K, Lindfors K. Coeliac disease specific IgA enhances the transepithelial passage of gliadin peptides. Helsinki Biomedical Graduate School, Student Council Symposium: Cell Structure and Organelles, May 8-9, 2008, Helsinki.
379. Stenman S, Hautala A, Venäläinen J, Männistö P, Kaukovirta-Norja A, Lindfors K, Mäki M, Kaukinen K. Reducing the toxic effects of rye secalin by germinating cereal enzymes. 16<sup>th</sup> United European Gastroenterology Week, Vienna, Austria, 18-22 October 2008. *Gut* 2008;57(Suppl II):A5.

380. Stenman S, Lindfors K, Venäläinen J, Männistö PT, Kaukinen K, Mäki M. Proteases from germinating wheat decrease the toxic effects of celiac disease-inducing wheat gliadin. Digestive Disease Week and the 109<sup>th</sup> Annual Meeting of the AGA Institute May 17-22, 2008, San Diego, CA, USA. *Gastroenterology* 2008;134 (Suppl 1):A80.
381. Stenman S, Hautala A, Venäläinen J, Männistö P, Kaukovirta-Norja A, Lindfors K, Mäki M, Kaukinen K. Reducing the toxic effects of rye secalin by germinating cereal enzymes. IMT Research day 2008, December 17, Tampere.
382. Ukkola A, Mäki M, Kurppa K, Collin P, Kekkonen L, Kaukinen K. Relevance of phenotypic characteristics of celiac disease in health-related quality of life: a prospective 1-year follow-up study. 13th International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book 2009:45-6(O1.2).
383. Kurppa K, Collin P, Sievänen H, Saavalainen P, Huhtala H, Mäki M, Kaukinen K. Increased morbidity in mild enteropathy coeliac disease. A prospective, randomized study with special focus on symptoms, quality of life and bone mineral density. 13th International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book 2009:54-5(O3.5).
384. Koskinen LLE, Einarsdottir E, Dukes E, Heap GAR, Dubois P, Korponay-Szabo IR, Kaukinen K, Kurppa K, Zibera F, Vatta S, Not T, Ventura A, Sistonen P, Adany R, Pocsai Z, Szeles G, Mäki M, Kere J, Wijmenga C, Van Heel DA, Saavalainen P. Association study of the IL18RAP locus in three European populations with coeliac disease. 13th International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book 2009:59(O5.1).
385. Hadjivassiliou M, Aeschlimann P, Sanders S, Korponay-Szabo I, Mäki M, Kaukinen K, Woodroffe N, Aeschlimann D. Antibodies against TG6 as the only serological marker of gluten ataxia. 13th International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book 2009:75(O9.1).
386. Lohi S, Mäki M, Rissanen H, Knekt P, Reunanen A, Kaukinen K. Prognosis of unrecognized celiac disease as regards mortality. 13th International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book 2009:96(P-035).
387. Ruuskanen AT, Kaukinen K, Collin P, Huhtala H, Valve R, Mäki M, Luostarinen L. Clinical relevance of positive serum antigliadin antibodies in elderly population. 13th International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book 2009:98(P-039).
388. Vilppula R-L, Kaukinen K, Luostarinen L, Krekelä I, Patrikainen H, Mäki M, Collin P. The incidence and prevalence of coeliac disease in elderly people. 13th International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book 2009:100(P-045).
389. Kurppa K, Mäki M, Saavalainen P, Partanen J, Collin P, Kaukinen K. Deamidated antigliadin antibodies are accurate markers for mild enteropathy coeliac disease. 13th International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book 2009:105(P-057).
390. Koskinen O, Collin P, Laurila K, Lindfors K, Mäki M, Kaukinen K. Usefulness of small-bowel mucosal transglutaminase-2 specific IgA autoantibody deposits in coeliac disease diagnosis and follow-up. 13th International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book 2009:106-7(P-060).
391. Bhatnagar S, Duttagupta S, Wadhwa N, Sodhi J, Bhatia V, Saini S, Sengupta S, Tandon N, Ranbir A, Jain V, Makharia G, Mäki M, Phillips AD, Not T. Point of contact testing for celiac disease in India. 13th International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book 2009:117(P-087).
392. Einarsdottir E, Koskinen LE, Dukes E, Kainu K, Suomela S, Lappalainen M, Zibera F, Korponay-Szabo IR, Kurppa K, Kaukinen K, Adany R, Pocsai Z, Szeles G, Färkkilä M, Turunen U, Halme L, Paavola-Sakki P, Not T, Vatta S, Ventura A, Löfberg R, Torkvist L, Bresso F, Halfvarson J, Mäki M, Kontula K, Saarialho-Kere U, Kere J, D'Amato M, Saavalainen PM. IL23R in the Swedish, Finnish, Hungarian and Italian populations: association with IBD and psoriasis, and linkage to coeliac disease. 13th International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book 2009:132-3(P-125).
393. Iltanen S, Ashorn S, Kurppa K, Ashorn M, Collin P, Mäki M, Wei B, Braun J, Kaukinen K. Serological responses to microbial antigens in patients with mild enteropathy coeliac disease. 13th International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book, 2009:141(P-146).
394. Caja Galan S, Martucciello S, Lavric M, Sblattero D, Marzari R, Mäki M, Kaukinen K, Lindfors K. Coeliac disease-specific autoantibodies targeted against transglutaminase 2 disturb vessel morphogenesis in vitro and in vivo through RhoB. 13th International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book 2009:143(P-150).
395. Nadalutti C, Viiri K, Kaukinen K, Mäki M, Lindfors K. Downregulation of transglutaminase 2 and its implications for coeliac disease. 13th International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book 2009:143(P-151).
396. Martucciello S, Caja S, Mäki M, Kaukinen K, Lindfors K. The autoantibodies anti-TG2 can modulate gene expression related with angiogenesis in endothelial cells. 13th International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book 2009:143-4(P-152).
397. Rauhavirta T, Kaukinen K, Korponay-Szano IR, Mongeot A, Collighan R, Griffin M, Mäki M. The transepithelial passage of gliadin peptides is enhanced in the presence of coeliac disease-specific IgA. 13th

- International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book 2009:145(P-155).
398. Myrsky E, Caja S, Korponay-Szabo IR, Collighan R, Mongeot A, Griffin M, Mäki M, Kaukinen K, Lindfors K. Coeliac disease-specific autoantibodies modulate vascular permeability in vitro. 13th International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book 2009:146(P-158).
399. Korponay-Szabo I, Kiraly R, Nemes E, Toth B, Vecsei Z, Juuti-Uusitalo K, Csipo A, Fesus L, Mäki M. Materno-fetal transmission of celiac autoimmunity. 13th International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book 2009:149-50(P-168).
400. Pusa E, Kaukinen K, Mäki M, Wacklin P, Partanen J, Mättö J. Composition of the mucosa-associated microbiota in the small intestine of coeliac disease patients and controls. 13th International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book 2009:171(P-224).
401. Korponay-Szabo I, Tumpek J, Nemes E, Papp M, Kovacs JB, Szabados K, Mäki M. Gluten-dependency of thyroid autoimmunity in celiac disease. 13th International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book 2009:185(P-259).
402. Stenman M, Lindfors K, Hautala A, Venäläinen J, Männistö P, Kaukovirta-Norja A, Mäki M, Kaukinen K. Reducing the toxic effects of rye secalin by germinating cereal enzymes in vitro-implications for future therapeutics in coeliac disease. 13th International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book 2009:193(P-278).
403. De Kauwe L, Anderson R, Kaukinen K, Tye-Din J, Kurppa K, Stewart J, Koskinen L, Mäki M, Saavalainen P. Short-term gluten challenge in Finnish patients with DQ2+ coeliac disease. 13th International Coeliac Disease Symposium 2009. April 6-8, 2009, Amsterdam. Programme & Abstract Book 2009:197(P-288).